

This is the html version of the file [ftp://ftp.sheridan.com/pub/content_services/ADA/DiabetesCare/Piccinni_dc102412_70dcoo.pdf](http://ftp.sheridan.com/pub/content_services/ADA/DiabetesCare/Piccinni_dc102412_70dcoo.pdf).
 Google automatically generates html versions of documents as we crawl the web.

Epidemiology/Health Services Research
 BRIEF REPORT

Assessing the Association of Pioglitazone Use and Bladder Cancer Through Drug Adverse Event Reporting

CARLO PICCINNI, PHD¹
 DOMENICO MOTOLA, PHD¹

GIULIO MARCHESINI, MD²
 ELISABETTA POLUZZI, PHD¹

OBJECTIVE—To analyze the association between pioglitazone use and bladder cancer through a spontaneous adverse event reporting system for medications.

RESEARCH DESIGN AND METHODS—Case/noncase bladder cancer reports associated with antidiabetic drug use were retrieved from the U.S. Food and Drug Administration (FDA) Adverse Event Reporting System (AERS) between 2004 and 2009 and analyzed by the reporting odds ratio (ROR).

RESULTS—Ninety-three reports of bladder cancer were retrieved, corresponding to 138 drug-reaction pairs (pioglitazone, 31; insulin, 29; metformin, 25; glimepiride, 13; exenatide, 8; others, 22). ROR was indicative of a definite risk for pioglitazone (4.30 [95% CI 2.82–6.52]), and a much weaker risk for gliclazide and acarbose, with very few cases being treated with these two drugs (6 and 4, respectively).

CONCLUSIONS—In agreement with preclinical and clinical studies, AERS analysis is consistent with an association between pioglitazone and bladder cancer. This issue needs constant epidemiologic surveillance and urgent definition by more specific studies.

Diabetes Care 34:1–3, 2011

Ader cancer first appeared in preclinical studies and was first reported on the U.S. pioglitazone label in 1999, but experimental studies recently suggested that it might be a rat-specific phenomenon (1). In the large PROactive (PROspective pioglitazone Clinical Trial In macroVascular Events) study, 14 bladder cancers occurred in the pioglitazone arm (0.5%) versus 6 in the placebo arm (0.2%) (2,3), and in September 2010, the U.S. Food and Drug Administration (FDA) announced an ongoing investigation on the possible risk in humans (4). Accordingly, the drug manufacturer is conducting a 10-year observational study to address the long-term risk of bladder cancer associated with pioglitazone (4).

Very recently, the European Medicines Agency (EMA) suspended the marketing

authorization of rosiglitazone (5), and the FDA largely restricted its use because of an increased cardiovascular risk (6). These measures will increase the prescription of pioglitazone; thus, the definition of its benefit/risk profile becomes all the more pressing.

Our aim was to contribute to defining the safety profile of pioglitazone, focusing on cases of bladder cancer recorded in the FDA Adverse Event Reporting System (AERS) database associated with antidiabetic drug treatment.

RESEARCH DESIGN AND METHODS

The reports recorded in FDA AERS from January 2004 to December 2009 were downloaded from the FDA website. The system contains all reports of adverse drug events spontaneously reported by health care professionals, manufacturers,

and consumers from the U.S. and serious and unlabeled spontaneous reports from non-U.S. countries. The adverse events are codified by Medical Dictionary for Regulatory Activities (MedDRA) terminology. Reports concerning antidiabetic drugs were selected, provided that age, sex, and event date were available. Duplicates and multiple records, a well-known drawback of FDA AERS (7), were excluded by a semiautomated multistep process (8).

The association between antidiabetic drugs and bladder cancer was analyzed by the case/noncase methodology (9). Cases were the reports retrieved under the MedDRA high-level term “bladder neoplasms” for any given drug; noncases were all of the other reports related to the same drug. The association between the drug and bladder cancer was calculated by the adverse drug reaction reporting odds ratio (ROR) as a measure of disproportionality. The ratio cases/noncases for each drug were compared with the ratio of cases/noncases for all other antidiabetic drugs. Stratified analyses weighed the influence of male sex and old age. The possible effect of notoriety bias was tested by a year-by-year analysis. Epi Info 3.4.3-2007 software (Centers for Disease Control and Prevention, Atlanta, GA) was used for statistical analyses.

RESULTS—From 2004 to 2009, 86,987 reports involving antidiabetic drugs were recorded in FDA AERS, corresponding to 599,085 drug-reaction pairs (obtained by splitting comedications and multiple reactions reported for each case), with 37,841 reports concerning pioglitazone. Overall, 93 reports of bladder cancer were retrieved, corresponding to 138 drug-reaction pairs, with 31 concerning pioglitazone; 29 insulin; 25 metformin; 13 glimepiride; 8 exenatide; 6 gliclazide; 5 glipizide; 4 sitagliptin, acarbose and rosiglitazone; 3 glibenclamide; 2

From the ¹Department of Pharmacology, University of Bologna, Bologna, Italy; and ²Metabolic Diseases & Clinical Dietetics, Department of Clinical Medicine, University of Bologna, Bologna, Italy.
 Corresponding author: Elisabetta Poluzzi, e-mail: elisabetta.poluzzi@unibo.it.

Received 22 December 2010 and accepted 11 March 2011.

DOI: 10.2337/dc10-2412

This article contains Supplementary Data online at <http://care.diabetesjournals.org/lookup/suppl/doi:10.2337/dc10-2412/-/DC1>.

© 2011 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See <http://creativecommons.org/licenses/by-nc-nd/3.0/> for details.

care.diabetesjournals.org

nateglinide and repaglinide; and 1 phenformin and voglibose.

The ROR of bladder cancer was significantly .1 for pioglitazone (ROR 4.30 [95% CI 2.82–6.52]; *P* , 0.001) as well as

DIABETES CARE, VOLUME 34, JUNE 2011

1

Pioglitazone and bladder cancer risk

for gliclazide and acarbose (Table 1). Among the 31 cases of bladder cancer reported in pioglitazone users (mean age, 70 years; range 53–84), 23 occurred in men (3.86 [2.37–6.26]; Supplementary Table A1) and 8 were in women (5.19 [2.15–12.11]). When stratified by age (cutoff, 65), ROR for pioglitazone was only significant in older patients (5.10 [3.14–8.23]). Four cases of bladder cancer were reported in 2004, three in 2005, nine in 2006, five in 2007, six in 2008, and four in 2009 (ROR not statistically significant in 2005 and 2009; Supplementary Table A2).

Ten cases occurred during clinical studies. The length of drug use, which was recorded in 15 cases, was .6 months in 6 patients, 6–24 months in 5, and .24 months in 4. Antiplatelet agents (e.g., aspirin and clopidogrel), antihypertensive drugs (e.g., ACE inhibitors and diuretics), lipid-lowering agents (e.g., statins), other antidiabetic drugs (e.g., glimepiride, metformin, and acarbose), and glucocorticoid (fluticasone and mometasone) were the cotreatments most frequently recorded (24 patients). One patient was being treated with cytotoxic therapy (infliximab and methotrexate for psoriatic arthropathy), and one was treated with interferon-β-1α for multiple sclerosis.

CONCLUSIONS—Bladder cancer is the fourth most common cancer and the ninth leading cause of cancer death among U.S. men (10). Cigarette smoking, urinary tract infections, occupational

exposure to aromatic amines and polycyclic aromatic hydrocarbons, and drugs (e.g., cyclophosphamide) are risk factors for the disease, as might be the systematic use of glucocorticoids (11).

We found a definite signal for bladder cancer associated with pioglitazone use. The demographic characteristics of the selected cases were consistent with bladder cancer epidemiology (male sex, old age) (10). A weaker signal was also associated with gliclazide, and a much weaker signal was associated with acarbose. Of note, the occurrence of fewer than five events, although resulting in a statistically significant ROR, may be considered clinically meaningless because it is too susceptible to reporting biases (12).

Although notoriety bias may have contributed to part of the association between pioglitazone use and bladder cancer (13), we also observed a significant relationship in 2004, which preceded publication of the PROactive study (2) and label revision. Therefore, we do not believe that our findings can be explained by notoriety bias alone. A greater use of pioglitazone could also have influenced this result (14).

Preliminary data found an increasing risk of bladder cancer with pioglitazone exposure, with statistical significance after 24 months (4). This issue could not be confirmed by our analysis, with only four cases of bladder cancer occurring in patients exposed to pioglitazone for more than 2 years and several missing data. In general, the association with bladder

cancer does not seem to derive from concomitant drug use or comorbidity, with only two patients receiving treatments potentially favoring carcinogenesis and five patients receiving glucocorticoids.

The ROR analysis has several limitations: generic under-reporting, over-reporting generated by notoriety bias, dependence on the drug-marketing period (Weber effect), missing or misspelled data (7,13,15), and lack of information on patients' habits (smoking) or occupational risks. Despite limitations, the higher-than-expected reporting of bladder cancer for pioglitazone users compared with users of other antidiabetic drugs should stimulate specific case-control studies aimed at verifying the magnitude of the hazard; until the final data of the FDA investigation are available, physicians should pay careful attention to this possible risk.

Acknowledgments—No potential conflicts of interest relevant to this article were reported.

C.P. researched data and wrote the manuscript. D.M. reviewed and edited the manuscript. G.M. contributed to discussion and reviewed and edited manuscript. E.P. researched data and wrote the manuscript.

Parts of this work was submitted in abstract form to the 10th Congress of the European Association for Clinical Pharmacology and Therapeutics, Budapest, Hungary, 26–29 June 2011.

References

1. Suzuki S, Arnold LL, Pennington KL, et al. Effects of pioglitazone, a peroxisome proliferator-activated receptor gamma agonist, on the urine and urothelium of the rat. *Toxicol Sci* 2010;113:349–357
2. Dormandy JA, Charbonnel B, Eckland DJ, et al.; PROactive investigators. Secondary prevention of macrovascular events in patients with type 2 diabetes in the PROactive Study (PROspective pioglitAZone Clinical Trial In macroVascular Events): a randomised controlled trial. *Lancet* 2005;366:1279–1289
3. Dormandy J, Bhattacharya M, van Troostenburg de Bruyn AR; PROactive investigators. Safety and tolerability of pioglitazone in high-risk patients with type 2 diabetes: an overview of data from PROactive. *Drug Saf* 2009;32:187–202
4. Food and Drug Administration. *FDA Drug Safety Communication: ongoing safety re-*

Table 1—ROR of bladder cancer for antidiabetic drugs

Active substance	Cases*	All ADR	ROR	95% CI	<i>P</i> †
Pioglitazone	31	37,841	4.30	2.82–6.52	,0.001
Insulin	29	124,873	1.01	0.06–1.55	0.961
Metformin	25	138,900	0.73	0.46–1.15	0.158
Glimepiride	13	35,388	1.66	0.89–3.01	0.080
Exenatide	8	100,946	0.30	0.14–0.64	0.001
Gliclazide	6	7,560	3.56	1.42–8.39	0.001
Glipizide	5	34,816	0.61	0.22–1.54	0.272
Sitagliptin	4	11,638	1.51	0.48–4.22	0.416
Acarbose	4	3,479	5.12	1.61–14.33	,0.001
Rosiglitazone	4	44,006	0.38	0.12–1.05	0.045
Glibenclamide	3	38,214	0.33	0.08–1.06	0.043
Nateglinide	2	4,994	1.75	N.A.	N.A.
Repaglinide	2	6,060	1.44	N.A.	N.A.

Phenformin	1	65	68.30	N.A.	N.A.
Voglibose	1	2,938	1.48	N.A.	N.A.
Other antidiabetic drugs	0	7,367	N.A.	N.A.	N.A.
Total	138	599,085			

ADR, adverse drug reaction; N.A., not available. *Cases of bladder cancer. †Mantel-Haenszel corrected.

view of actos (pioglitazone) and potential increased risk of bladder cancer after two years exposure. [Internet] 2010. Silver Spring, MD, Food and Drug Administration. Available from <http://www.fda.gov/>

- Drugs/DrugSafety/ucm226214.htm. Accessed 21 October 2010
- European Medicines Agency. European Medicines Agency update on ongoing benefit-risk review of Avandia, Avandia and Avaglim. [Internet] 2010. London, U.K., European Medicines Agency. Available from http://www.ema.europa.eu/docs/en_GB/document_library/Press_release/2010/07/WC500094981.pdf. Accessed 25 October 2010
 - Food and Drug Administration. FDA significantly restricts access to the diabetes drug Avandia. [Internet] 2010. Silver Spring, MD, Food and Drug Administration. Available from <http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/ucm226956.htm>. Accessed 25 October 2010
 - Hauben M, Reich L, DeMicco J, Kim K. 'Extreme duplication' in the US FDA Adverse Events Reporting System database. *Drug Saf* 2007;30:551–554
 - Poluzzi E, Raschi E, Motola D, Moretti U, De Ponti F. Antimicrobials and the risk of torsades de pointes: the contribution from data mining of the US FDA Adverse Event Reporting System. *Drug Saf* 2010;33:303–314
 - Moore N, Kreft-Jais C, Haramburu F, et al. Reports of hypoglycaemia associated with the use of ACE inhibitors and other drugs: a case/non-case study in the French pharmacovigilance system database. *Br J Clin Pharmacol* 1997;44:513–518
 - Jemal A, Siegel R, Xu J, Ward E. Cancer statistics, 2010. *CA Cancer J Clin* 2010;60:277–300
 - Dietrich K, Schned A, Fortuny J, et al. Glucocorticoid therapy and risk of bladder cancer. *Br J Cancer* 2009;101:1316–1320
 - van Puijenbroek EP, Bate A, Leufkens HG, Lindquist M, Orre R, Egberts AC. A comparison of measures of disproportionality for signal detection in spontaneous reporting systems for adverse drug reactions. *Pharmacoepidemiol Drug Saf* 2002;11:3–10
 - Pariante A, Gregoire F, Fourier-Reglat A, Haramburu F, Moore N. Impact of safety alerts on measures of disproportionality in spontaneous reporting databases: the notoriety bias. *Drug Saf* 2007;30:891–898
 - Cohen A, Rabbani A, Shah N, Alexander GC. Changes in glitazone use among office-based physicians in the U.S., 2003–2009. *Diabetes Care* 2010;33:823–825
 - Almenoff J, Tonning JM, Gould AL, et al. Perspectives on the use of data mining in pharmaco-vigilance. *Drug Saf* 2005;28:981–1007

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES

- Q1: According to the FDA Web site, "System" is part of the acronym; should be AERS rather than AER System.
- Q2: The title of ref 5 has been adjusted to match the title of the PDF that is cited in the reference. Please check.
- Q3: Trademarked items generally require the supplier and location. Please check for EpiInfo. Note name changed per CDC Web site.
- Q4: Check that the conflict of interest information for each author is presented in full in the Acknowledgments section.