

Lactose intolerance: diagnosis, genetic, and clinical factors

Rejane Mattar
Daniel Ferraz de Campos
Mazo
Flair José Carrilho

Department of Gastroenterology,
University of São Paulo School
of Medicine, São Paulo, Brazil

Abstract: Most people are born with the ability to digest lactose, the major carbohydrate in milk and the main source of nutrition until weaning. Approximately 75% of the world's population loses this ability at some point, while others can digest lactose into adulthood. This review discusses the lactase-persistence alleles that have arisen in different populations around the world, diagnosis of lactose intolerance, and its symptomatology and management.

Keywords: hypolactasia, lactase persistence, lactase non-persistence, lactose, *LCT* gene, *MCM6* gene

Introduction

Lactose is a disaccharide that is abundant in mammalian milk and essential for the nourishment of newborn infants. It is hydrolyzed by the intestinal brush-border enzyme, lactase, into absorbable sugars, namely glucose and galactose. In most infants, intestinal lactase activity is maximal during the perinatal period; however, after 2–12 years of age, two distinct groups emerge, ie, a “lactase non-persistence” group with low lactase activity (hypolactasia) and a “lactase-persistence” group of individuals who retain their neonatal level of lactase activity into adulthood.^{1–3}

Reduction in lactase activity causes primary maldigestion of lactose, a condition that is occasionally asymptomatic. When symptoms are present, lactose intolerance is diagnosed. It is important to distinguish between primary hypolactasia and secondary causes of maldigestion of lactose, including celiac disease, infectious enteritis, or Crohn's disease, which have distinct pathogenic and therapeutic implications. Moreover, primary hypolactasia should be distinguished from congenital lactase deficiency, a rare autosomal recessive disease with unique molecular mechanisms that affects infants from birth.⁴

Lactase-persistence alleles and polymorphisms for lactose tolerance

The *LCT* gene is 49.3 kb in length and located on the long (q) arm of chromosome 2 at position 21. It contains 17 exons and is translated into a 6 kb transcript (NCBI Reference Sequence NG_008104.1). Individuals with hypolactasia and lactase persistence have identical coding sequences, except for some silent mutations; thus, both lactases are identical.⁵

Enattah et al⁶ devised a brilliant strategy using polymorphic microsatellite markers flanking *LCT*, encompassing a region of 47 kb, in a haplotype linkage analysis of

Correspondence: Rejane Mattar
Hospital das Clínicas da FMUSP, Av Dr
Enéas de Carvalho Aguiar 255, 9ºAndar,
Sala 9159, São Paulo, SP, Brazil, 05403-000
Tel +55 11 2661 6150
Fax +55 11 2661 7830
Email r.mattar@hc.fm.usp.br

nine Finnish families with hypolactasia. Two variants were associated with lactase persistence. A polymorph variant, *LCT*-13910C>T, in intron 13 of the *MCM6* gene that is 13,910 bp from the initiation codon of *LCT*, demonstrated a complete association, while the *LCT*-22018G>A variant in intron 9 of *MCM6* gene upstream of the *LCT* locus 22,018 bp was strongly, but not completely, associated.^{1,2,6} The functional role of *MCM6* in vertebrates is unknown, but it has been implicated in “licensing” DNA replication during the cell cycle.¹ This association was confirmed in a study of DNA collected from subjects of Finnish, South Korean, Italian, German, French, or white or African North American descent.^{1,6}

In subjects of European descent, the *LCT*-13910C>T variant completely associated with the lactase-persistence phenotype and presented different allelic frequencies in countries within Europe, Oceania, Asia, and the Americas, as shown in Table 1.

Both genotypes of *LCT*-13910CT and *LCT*-13910TT were associated with the lactase-persistence phenotype, indicating that the presence of one single lactase-persistence allele in the heterozygous state has a dominant effect, rendering the person a lactose digester, whereas the genotype *LCT*-13910CC, when the lactase-persistence allele *LCT*-13910T is absent, is consistent with lactose maldigestion.^{2,3}

Despite the association of *LCT*-13910C>T with lactose digestion in Europeans, analysis of this variant in Africa demonstrated its restriction to populations with a high prevalence of the lactase-persistence phenotype (Table 2). This finding suggests the presence of other lactase-persistence alleles (Table 3). Thus, as shown in Figure 1, different alleles have originated in various locations around the world over the course of human history after the emergence of modern man from Africa.¹⁷

Genotyping of *LCT*-13910C>T versus *LCT*-22018G>A has shown almost full agreement. Patients with *LCT*-13910CC were also *LCT*-22018GG, while individuals with *LCT*-13910CT had the *LCT*-22018GA genotype. *LCT*-13910TT was associated with *LCT*-22018AA, except for a few cases in Finland⁶ and China,³² and in Japanese Brazilians.³³

Functional in vitro studies of these polymorphic alleles have shown that *LCT*-13910T,^{1,34,35} *LCT*-13907G, *LCT*-13915G, and *LCT*-14010C act as enhancers of the *LCT* promoter²⁹ unlike in ancestral constructs (*LCT*-13910C, *LCT*-13907C, *LCT*-13915T, and *LCT*-14010G). These effects are most likely mediated by the Oct-1 transcriptional factor binding site in the variant enhancer and by HNF1 α binding

Table 1 Frequencies of the European variant *LCT*-13910C>T in countries within the Americas, Asia, Europe, and Oceania

Country or population	Allele frequency (%)	Reference
US (Utah)	74.5	7
Sweden	73.7	8
New Zealand (Christchurch)	72	9
The Netherlands (Rotterdam Study)	69	10
Basques	65.9	7
Finland	58.1	11
Austria	53	12
Estonia (Väike-Maarja)	51.4	13
Poland	43.9	11
Russia (northern)	38.9	15
Portugal (northern)	37	17
Canary Islands	36.5	18
Hungary	35.9	16
Kola Sami (Murmansk)	30.5	14
Brazil (Caucasian)	24.7	21
Italy (North-east)	23.7	20
Chile (Hispanics)	22	22
India (Northern)	19.5	19
Brazil (African origin)	18.3	21
Uzbekistan (Kazakh, nomadic)	15.7	23
Italy (North-central)	13.3	20
Italy (Central)	13; 11.2	17,20
Uzbekistan (Tajiko-Uzbek)	10	23
Greece	9	20
US (African origin)	9	7
Italy (Southern)	5.5; 8	7,20
Sardinia	7.2	20
India (South)	6.6	19
Chile (Amerindians)	5.8	22
China	0	7
Japanese Brazilian	0	21

Note: In some publications, the percentage of *LCT*-13910-C>T allele frequencies were calculated based on the number of individuals with the *LCT*-13910-CT and *LCT*-13910-TT genotypes in relation to the total.

in the *LCT* promoter. However, further evaluation is required to determine whether these actions correspond to the situation in vivo.^{34–36}

LCT gene regulation of lactase-persistence alleles occurs at the transcriptional level. *LCT* mRNA levels, which are distinguished by polymorphic markers in the coding region of *LCT*, were several times higher in individuals with *LCT*-13910T/-22018A alleles than in individuals with *LCT*-13910C/-22018G alleles.¹ After 5 years of age, an imbalance appears in the mRNA levels of *LCT*-13910C and *LCT*-13910T, with the *LCT*-13910T allele representing approximately 92% of *LCT* mRNA in children heterozygous for *LCT*-13910CT.³

Several transcription factors (Cdx2, GATA-4, GATA-5, GATA-6, and HNF1 α) activate the *LCT* promoter in intestinal cell culture at the –100 to –20 bp binding site regions of

Table 2 Frequencies of the lactase persistence allele (*LCT*-13910C>T) reported in African countries

Country and/or population	Allele frequency (%)	Reference
Cameroon (Fulbe)	11.2, 21, 39	24,17,25
Mali (Fulbe)	37	26
South Africa (Xhosa mixed)	21.8	27
Morocco	17.3	7
Cameroon (Hausa)	13.9	24
Cameroon (agricultural)	4.3	24
São Tomé	4	17
Somalia	3.2	7
Senegal	2.6	24
Mozambique	1	17
Ethiopia	1.9	28
(Somali camel herders)		
Nigeria	0	24
Malawi	0	24
Sudan (north and south)	0	24
Ethiopia	0	24
Uganda	0	24

LCT which are repressed by PDX-1.¹ Mutation of the PDX-1 binding site does not prevent *LCT* promoter repression, which suggests that PDX-1 might function by binding to another DNA binding site or by inhibiting other transcriptional factors. PDX-1 overexpression resulted in strong repression

of *Cdx2* and *HNF1 α* activation of the *LCT* promoter.¹ However, the exact mechanism for downregulation of *LCT* after weaning is unknown.

Haplotype conservation around lactase-persistence alleles indicates that these alleles emerged recently in different parts of the world and have been subject to strong positive selection in communities of high and perhaps intermittently exclusive consumers of fresh milk.²⁸ Nevertheless, the selective advantage provided by drinking fresh milk is not yet clear among populations reliant on agriculture with dairy farming as their main source of income, but has been discussed in detail elsewhere.³⁷ Gene-culture coevolution is a likely hypothesis in Africa, because high lactase-persistence allele frequencies are preferentially found in pastoral communities. In populations more likely to consume agricultural products, cheese and fermented milk, which have lower concentrations of lactose, the frequencies of lactase-persistence variants are possibly due to genetic drift.³⁸

It is estimated that the *LCT*-13910T allele initially originated on the background of a more common haplotype approximately 5000–12,000 years ago and re-emerged recently (1400–3000 years ago) on another haplotype background in restricted populations west of the Urals and north of the Caucasus.⁷ The *LCT*-13907G and *LCT*-13910T alleles share the

Table 3 Frequencies of other lactase persistence alleles in the *MCM6* gene

Country or population	Alleles	Frequency (%)	Reference
Saudi Arabia	<i>LCT</i> -13915T>G	48.9; 59.4	25,30
Jordan		39.1	25
Sudan (Beni Amir)		24.4	25
Ethiopia (Afar)		15	25
Sudan (Jaali)		14.2	25
Ethiopia (Amharic)		13.2	25
Ethiopia (Somali camel herders)		5.1	28
Tanzania	<i>LCT</i> -14010G>C	31.9	29
Kenya		27.6	29
Xhosa (South Africa)		12.8	27
Xhosa (mixed ancestry)		8.1	27
Angola		<7	17
Mozambique		No LP allele	17
Ethiopia (Somali camel herders)		0.5	28
Sudan (Afro-Asiatic Beja)	<i>LCT</i> -13907C>G	20.6	29
Ethiopia (Afar)		20	25
Ethiopia (Somali camel herders)		5.6	28
Northern Russia	<i>LCT</i> -13914G>A	Rare variant	31
Austria		Two individuals	12
China (Kazak)	<i>LCT</i> -22018G>A/	18	32
China (Northern)	<i>LCT</i> -13910CC	6.8	32
Japanese Brazilians		5.3	33
Tanzania (Akie)		One individual	29
Sudan (Jaali)	<i>LCT</i> -14009T>G	6.6	28
Ethiopia (Somali camel herders)		1.4	28

Abbreviation: LP, lactase persistence.




Figure 1 Tendency of lactase-persistence polymorphic variants in the world, based on the reports presented in Tables 1, 2, and 3.

Notes: In places where there was more than one variant, the most frequent variant was considered. ● *LCT-13910C>T*; ● *LCT-22018G>A/-13910CC*; ● *LCT-13915T>G*; ● *LCT-14010G>C*; ● *LCT-13907C>G*.

same ancestral lactase non-persistence haplotype, although they are on backgrounds of different lactase-persistence haplotypes.^{25,28,35} *LCT-13915G* and *LCT-14010C* originated on different haplotype backgrounds,^{25,28,29,35} but age estimates are similar for both, at approximately 4095 ± 2045 years.³⁵

Diagnosis

Initially, the most accurate method available for the diagnosis of lactose maldigestion was direct biochemical assay of lactase activity from a jejunal sample. This assay is performed with a glucose oxidase reagent, which detects glucose liberated from lactose, with a cutoff value of 10 U/g protein.^{1,2} Due to the invasiveness of a jejunal biopsy, this method has been replaced by endoscopic duodenal biopsy.^{39,40} Mean lactase activity was about 40% lower in the duodenum compared with the jejunum,³⁹ but the Quick lactase test performed in samples taken from the postbulbar duodenum effectively identified patients with severe duodenal hypolactasia, with a sensitivity and specificity of 95% and 100%, respectively.⁴⁰

Lactose tolerance tests have been developed to confirm the ability of intestinal lactase to hydrolyze and absorb lactose, and to avoid intestinal biopsies. Blood glucose levels were measured before and after an oral load of lactose at prespecified time intervals, with a maximum rise of 20 mg/dL, indicating lactose tolerance.⁴¹ Oral ethanol administration before lactose load is used to inhibit galactose metabolism for the determination of the blood maximum rise of glucose (at least 20 mg/dl) and galactose (at least 10 mg/dl), indicating

lactose tolerance. Thus, galactose concentration in combination with glucose concentration improves the correlation with jejunal lactase activity than using only glucose maximum rise after lactose load.⁴² Nonetheless, of all the indirect lactose tolerance tests currently available, breath hydrogen after ingestion of 50 g of lactose was considered the most suitable test for population screening for lactase deficiency.⁴³ Use of the 50 g lactose dose has been criticized, because it is equivalent to 4–5 cups of milk, an amount that is far more than an individual usually ingests at one time,⁴⁴ so an oral load of 25 g, ie, the mean quantity contained in 500 mL of semiskimmed milk, may be considered a more appropriate amount, with high sensitivity and specificity.^{41,44,45}

The lactose breath test is based on fermentation of undigested lactose by intestinal flora, producing hydrogen, carbon dioxide, and methane that are absorbed and eliminated via the lungs, but these gases also cause bloating, flatulence, abdominal pain, and diarrhea. Despite being widely used, the reliability of this test depends on the activity of bacterial flora. A false-negative result can occur if antibiotics have been taken within one month of being tested, if colonic pH is acidic enough to inhibit bacterial activity, or if there has been adaptation in the bacterial flora as a result of continuous lactose exposure.^{41,44,45}

The discovery of lactase-persistence alleles prompted use of genetic tests for diagnosis of lactase non-persistence by polymerase chain reaction restriction fragment length polymorphism,^{45–47} real-time polymerase chain reaction,^{48–50} and Pyrosequencing[®] technology.⁵¹ Compared with the lactose

hydrogen breath test, the genetic test is a simple, noninvasive, and more comfortable examination that does not provoke symptoms of lactose intolerance and is less cumbersome,^{46,51} with easy transfer of a venous blood sample to the laboratory.^{45,47} However, other polymorphic variants in Europeans (*LCT*-13914G>A)⁵⁰ and in African and Arab populations (*LCT*-13907C>G, *LCT*-13913T>C, and *LCT*-13915T>G, close to *LCT*-13910C>T, depicted in Table 3) affect the diagnostic accuracy of *LCT*-13910C>T typing by altering the melting profiles of the real-time polymerase chain reaction kit.⁵⁰ The reverse-hybridization strip assay based on multiplex DNA amplification and ready-to-use membrane test strips that detect *LCT* polymorphic variants (-13907C>G, -13910C>T, -13913T>C, -13914G>A, -13915T>G, and -22018G>A) represents a reliable tool for genetic diagnosis of lactase non-persistence, overcoming the interference of different melting profiles of the real-time polymerase chain reaction kit by the other polymorphic variants.¹²

The genetic test provides a more direct result, ie, a hypolactasia or lactase persistence genotype, whereas interpretation of the lactose breath test depends on the cutoff level, dose of lactose given, and duration of the test and age of the individual, among the other factors already discussed,^{45–47,49,51} and is costly.^{47,51} The discovery of other single nucleotide polymorphisms associated with lactase persistence (see Table 3) implies that DNA genotyping should provide information on the DNA sequence around the polymorphic site of the *MCM6* gene.⁵¹ In addition to the reverse-hybridization strip assay,¹² Pyrosequencing technology may be a cost-effective option (€10 per test for polymerase chain reaction and Pyrosequencing reagents) for direct DNA sequencing, allowing genotyping of other single nucleotide polymorphisms.⁵¹ The genetic test does not provide information on symptoms of lactose tolerance; however, measurement of lactase activity in intestinal biopsy does not provide it either.⁴⁷

Contribution of lactose ingestion to symptomatology

The age of onset of primary hypolactasia varies between different ethnic groups. Hypolactasia does not cause any disturbance or discomfort unless lactose-containing food is consumed. Colonic microflora ferment undigested lactose in the intestinal lumen, which leads to production of short-chain fatty acids, hydrogen, carbon dioxide, and methane. These byproducts cause bloating, flatulence, and abdominal pain. Undigested lactose acidifies the colon and increases the osmotic load, resulting in loose stools and diarrhea.⁵² Stools are usually voluminous, foamy, and aqueous. Although hypolactasia-related diarrhea can become

chronic, affected individuals typically do not lose weight. However, some patients can experience constipation due to decreased intestinal motility, possibly caused by production of methane.⁵²

Some authors have reported that the clinical presentation of lactose intolerance is not restricted to gut symptoms. Systemic complaints, such as headache, vertigo, memory impairment, lethargy, muscle and joint pains, allergy, cardiac arrhythmia, mouth ulcers, and sore throat, have been reported in less than 20% but up to 86% of these patients.^{53,54} Putative toxic metabolites, such as acetaldehyde, acetoin, ethanol, peptide, and protein toxins, can alter cell signaling mechanisms and are possibly responsible for these systemic symptoms. They are generated by lactose fermentation in colonic bacteria.^{53,55} When systemic complaints are present, it is important to assess whether they result from lactose intolerance, are coincidental, or emanate from an allergy to cow's milk protein, which is present in up to 20% of patients with symptoms of lactose intolerance.⁵² Minenna et al reported a possible association between gastroesophageal reflux disease and lactose malabsorption in 30 subjects; however, further studies are required to ascertain a causal relationship, given that both lactose intolerance and reflux are very common conditions.⁵⁶

There is considerable intraindividual and interindividual variability in the severity of symptoms, according to the amount of lactose ingested and the patient's ability to digest it. Factors contributing to this variability include osmolality and the fat content of lactose-containing food, gastric emptying rate, ability of colonic microflora to ferment lactose, intestinal transit time, colonic water absorption capacity, and individual perception of abdominal pain and discomfort.^{52,57} Valid evidence is missing for a relationship between symptoms and amount of lactose ingested.⁵⁷ Most studies have included a small number of participants and/or subjects, with lactose maldigestion diagnosed by the breath hydrogen test but not always concomitant with lactose intolerance. In this regard, the available data⁵⁸ demonstrate that a single dose of lactose (up to 12 g, equivalent to that contained in approximately one glass of milk) administered alone produces no or minor symptoms in persons with lactose intolerance or maldigestion. Lactose doses of 15–18 g are well tolerated when offered together with other nutrients. With doses larger than 18 g, intolerance becomes progressively more frequent, and quantities over 50 g elicit symptoms in most individuals.⁵⁸

Various reports indicate that symptoms typically considered secondary to lactose ingestion are not truly

related to maldigestion.^{59,60} On self-report questionnaires, individuals commonly associate ingestion of lactose-containing products with onset of abdominal symptoms, even in the absence of objective evidence for lactose maldigestion, such as an altered lactose breath test.^{61,62} Symptoms frequently attributed to lactose maldigestion can be secondary to irritable bowel syndrome,⁶² which shares a similar clinical presentation, or food allergy.⁶³ Even a “nocebo effect”, ie, occurrence of symptoms after ingestion of an inert substance when negative expectations about its content exist,⁶⁴ has been considered to be contributory to this exaggerated perception of lactose intolerance.⁶⁵ However, this concept requires more consistent evidence.⁶⁶ The misleading diagnosis of lactose intolerance and subsequent implementation of a dairy-restricted diet is not without consequences. The negative clinical impact of imposed restrictions, which mainly involve bone metabolism, is a topic that will be discussed in a following section.

Along with irritable bowel syndrome and cow’s milk protein allergy, the differential diagnosis of lactose intolerance includes bacterial overgrowth, celiac disease, and inflammatory bowel disease.^{57,67} When bloating and flatulence are the predominant symptoms, it is also advisable to rule out the possible contribution of other dietary sources of intestinal gas, such as beans, which contain two indigestible sugars, stachyose and raffinose.^{57,68}

Management

The goal of treatment is to improve symptoms while maintaining an adequate intake of calcium, thus preventing secondary bone disease caused by a milk-restricted diet. Considerable efforts have been made to confirm whether decreased lactase enzyme activity can impair calcium absorption and prevent attainment of optimal peak bone mass. When evaluating peak bone mass and bone turnover rate in a young population with molecularly defined lactose maldigestion, Enattah et al showed that hypolactasia and lactose maldigestion do not alter calcium absorption or bone turnover rate, nor do they impair acquisition of peak bone mass. Moreover, the *LCT*-13910CC genotype does not appear to be a risk factor for stress fractures in this population.⁶⁹ Although decreased calcium absorption, evaluated by the strontium absorption test in patients with the *LCT*-13910CC genotype, was reported by Obermayer-Pietsch et al,⁷⁰ the predominant idea in the literature is that low calcium intake, rather than deficient calcium absorption, is the major factor contributing to loss of bone mass.^{57,69} Several studies in patients with presumed or confirmed lactose intolerance have also reported lower calcium intake in this population.^{57,71,72}

Several reports have been published that address the relationship among the *LCT*-13910C>T genotype, lactose intolerance, bone mineral density, and fracture risk. Studies in postmenopausal women⁷³ and elderly people⁷⁴ with the *LCT*-13910CC genotype have identified lower bone mineral density and a higher incidence of bone fractures in comparison with individuals with other lactase genotypes. However, these results have not been confirmed by other studies^{75,76} or in younger subjects.^{69,77}

Recently, Tolonen et al showed that young men with the *LCT*-13910TT genotype had the highest bone trabecular density at the distal radius and tibia, but other bone traits or low-energy fractures were not associated with the *LCT*-13910C>T genotype.⁷⁸ In addition to height and bone parameters, Koek et al assessed the correlation between vitamin D receptor polymorphisms and *LCT*-13910C>T genotypes in the elderly. This study found that the *LCT*-13910CC genotype was associated with lower dietary calcium intake and lower serum calcium levels, but not with bone mineral density and fracture risk. No interaction was detected between *LCT*-13910C>T genotypes and vitamin D receptor polymorphisms.¹⁰

The available data suggest that deficient calcium intake plays a major role in lactose intolerance that may be related to bone disease. Therefore, an objective diagnosis through either the hydrogen breath test or molecular detection of hypolactasia is key to the appropriate clinical management of patients with symptoms suggestive of lactose intolerance. This approach avoids inappropriate calcium-restricted diets and adverse consequences for bone health.

The initial recommendation for management of lactose intolerance is to aim for remission of symptoms by temporarily avoiding milk and dairy products. As mentioned earlier, most individuals with lactose malabsorption can tolerate up to 12 g of lactose without significant symptoms. After the initially restricted diet, lactose should be gradually reintroduced until the patient’s threshold for symptoms is reached.⁷⁹ At this point, several behavioral measures can be adopted to overcome possible symptoms, including having fermented and matured milk products in the diet, consuming lactose together with other foods, and distributing lactose intake over the day. Although lactose tablets have been cited as a potential trigger of symptoms of lactose intolerance, such a small amount of lactose cannot be blamed for provoking symptoms, even when differences in individual symptom thresholds are considered.⁸⁰

If the measures suggested here do not suffice in reducing symptoms, pharmacological strategies can be implemented.

The main pharmacological measures in use include lactase supplements, lactose-hydrolyzed or lactose-reduced milk, probiotics, colonic adaptation, and rifaximin. Ingestion of probiotics containing lactase may have the potential to aid lactose digestion in intolerant patients, but studies that have investigated this have published conflicting results. Therefore, the role of probiotics in lactose intolerance management is currently uncertain.⁷⁹ Yoghurt containing live cultures providing endogenous beta galactosidase are an alternative source of calories and calcium, and are well tolerated by many lactose-intolerant patients. However, yoghurt containing milk or its derivatives added after fermentation can cause symptoms.⁷⁹ Overall, the available evidence-based data are insufficient to ascertain the efficacy of these interventions, as discussed at a recent National Institutes of Health conference.⁵⁸

Attention must be paid to daily ingestion of calcium and vitamin D, with supplementation as required. For adolescents and young adults, the dietary calcium recommendation is generally 1200–1500 mg. In adults, the amount varies according to gender and menopausal status. Calcium should be supplemented if there is not enough in the diet, and vitamin D should also be monitored and supplemented if necessary.⁷⁹ Well designed, randomized, placebo-controlled trials are still required before strong clinical recommendations can be made for the management of patients who are intolerant of lactose-hydrolyzed milk and yoghurt.

Conclusion

Random mutations have occurred in regions upstream of the *LCT* gene that have an enhancer effect on the *LCT* promoter, which enables carriers with the lactase-persistence phenotype to exist in populations all over the world. No “gold standard” test is available for the diagnosis of lactose intolerance. The lactose breath test, although considered the best method, may be influenced by confounding factors. Genetic testing has been a new tool for the diagnosis of hypolactasia/lactase persistence, but may not detect all the single nucleotide polymorphisms associated with this disorder. Symptoms of lactose intolerance might have been exaggerated, such that up to 12 g of lactose is possibly well tolerated by lactase non-persistence individuals, which negates the need for restrictions on lactose-hydrolyzed milk, as well as fermented and matured milk products, preventing any subsequent effects on bone mass density.

Acknowledgment

The authors thank Marcos Antonio Retzer for his contribution to Figure 1.

Disclosure

The authors have no conflict of interests to declare in this work.

References

1. Troelsen JT. Adult-type hypolactasia and regulation of lactase expression. *Biochim Biophys Acta*. 2005;1723:19–32.
2. Rasinperä H, Savilahti E, Enattah NS, et al. A genetic test which can be used to diagnose adult-type hypolactasia in children. *Gut*. 2004;53:1571–1576.
3. Rasinperä H, Kuokkanen M, Kolho KL, Lindahl H, Enattah NS, Savilahti E. Transcriptional down regulation of the lactase (*LCT*) gene during childhood. *Gut*. 2005;54:1660–1661.
4. Robayo-Torres CC, Nichols BL. Molecular differentiation of congenital lactase deficiency from adult-type hypolactasia. *Nutr Rev*. 2007;65:95–98.
5. Boll W, Wagner P, Mantei N. Structure of the chromosomal gene and cDNAs coding for lactase-phlorizin hydrolase in humans with adult-type hypolactasia or persistence of lactase. *Am J Hum Genet*. 1991;48:889–890.
6. Enattah NS, Sahi T, Savilahti E, Terwilliger JD, Peltonen L, Järvelä I. Identification of a variant associated with adult-type hypolactasia. *Nat Genet*. 2002;30:233–237.
7. Enattah NS, Trudeau A, Pimenoff V, et al. Evidence of still-ongoing convergence evolution of the lactase persistence T-13910 alleles in humans. *Am J Hum Genet*. 2007;81:615–625.
8. Almon R, Engfeldt P, Tysk C, Sjöström M, Nilsson TK. Prevalence and trends in adult-type hypolactasia in different age cohorts in Central Sweden diagnosed by genotyping for the adult-type hypolactasia-linked *LCT*-13910C>T mutation. *Scand J Gastroenterol*. 2007;42:165–170.
9. Upton J, George P. The prevalence of lactose intolerance (adult hypolactasia) in a randomly selected New Zealand population. *N Z Med J*. 2010;123:117–118.
10. Koek WN, van Meurs JB, van der Eerden BC, et al. The T-13910C polymorphism in the lactase phlorizin hydrolase gene is associated with differences in serum calcium levels and calcium intake. *J Bone Miner Res*. 2010;25:1980–1987.
11. Kuokkanen M, Butzow R, Rasinperä H, et al. Lactase persistence and ovarian carcinoma risk in Finland, Poland and Sweden. *Int J Cancer*. 2005;117:90–94.
12. Tag CG, Oberkanins C, Kriegshäuser G, et al. Evaluation of a novel reverse-hybridization StripAssay for typing DNA variants useful in diagnosis of adult-type hypolactasia. *Clin Chim Acta*. 2008;392:58–62.
13. Lember M, Torniainen S, Kull M, et al. Lactase non-persistence and milk consumption in Estonia. *World J Gastroenterol*. 2006;12:7329–7331.
14. Kozlov A, Borinskaya S, Vershubsky G, et al. Genes related to the metabolism of nutrients in the Kola Sami population. *Int J Circumpolar Health*. 2008;67:56–66.
15. Khabarova Y, Torniainen S, Nurmi H, Järvelä I, Isokoski M, Mattila K. Prevalence of lactase persistent/non-persistent genotypes and milk consumption in a young population in north-west Russia. *World J Gastroenterol*. 2009;15:1849–1853.
16. Nagy D, Tömöry G, Csányi B, et al. Comparison of lactase persistence polymorphism in ancient and present-day Hungarian populations. *Am J Phys Anthropol*. 2011;145:262–269.
17. Coelho M, Luiselli D, Bertorelle G, et al. Microsatellite variation and evolution of human lactase persistence. *Hum Genet*. 2005;117:329–339.
18. Almon R, Álvarez-Leon EE, Engfeldt P, Serra-Majem L, Magnuson A, Nilsson TK. Associations between lactase persistence and the metabolic syndrome in a cross-sectional study in the Canary Islands. *Eur J Nutr*. 2010;49:141–146.

19. Babu J, Kumar S, Babu P, Prasad JH, Ghoshal UC. Frequency of lactose malabsorption among healthy southern and northern Indian populations by genetic analysis and lactose hydrogen breath and tolerance tests. *Am J Clin Nutr*. 2010;91:140–146.
20. Anagnostou P, Battaggia C, Coia V, et al. Tracing the distribution and evolution of lactase persistence in southern Europe through the study of the T-13910 variant. *Am J Hum Biol*. 2009;21:217–219.
21. Mattar R, Monteiro MS, Villares CA, Santos AF, Silva JMK, Carrilho FJ. Frequency of *LCT*-13910C>T single nucleotide polymorphism associated with adult-type hypolactasia/lactase persistence among Brazilians of different ethnic groups. *Nutr J*. 2009;8:46.
22. Morales E, Azocar L, Maul X, Perez C, Chianale J, Mique JF. The European lactase persistence genotype determines the lactase persistence state and correlates with gastrointestinal symptoms in the Hispanic and Amerindian Chilean population: a case-control and population-based study. *BMJ Open*. 2011;1:e000125.
23. Heyer E, Brazier L, Ségurel L, et al. Lactase persistence in Central Asia: phenotype, genotype, and evolution. *Hum Biol*. 2011;83:379–392.
24. Mulcare CA, Weale ME, Jones AL, et al. The allele of a single-nucleotide polymorphism 13.9 kb upstream of the lactase gene (*LCT*) (C13.9kbT) does not predict or cause the lactase-persistence phenotype in Africans. *Am J Hum Genet*. 2004;74:1102–1110.
25. Ingram CJE, Elamin FF, Mulcare CA, et al. A novel polymorphism associated with lactose tolerance in Africa: multiple causes for lactase persistence? *Hum Genet*. 2007;120:779–788.
26. Lokki AI, Järvelä I, Israelsson E, et al. Lactase persistence genotypes and malaria susceptibility in Fulani of Mali. *Malar J*. 2011;10:9.
27. Torniaainen S, Parker MI, Holmberg V, Lahtela E, Dandara C, Järvelä I. Screening of variants for lactase persistence/non-persistence in populations from South Africa and Ghana. *BMC Genet*. 2009;10:31.
28. Ingram CJE, Raga TO, Tarekegn A, et al. Multiple rare variants as a cause of a common phenotype: several different lactase persistence associated alleles in a single ethnic group. *J Mol Evol*. 2009;69:579–588.
29. Tishkoff SA, Reed FA, Ranciaro A, et al. Convergent adaptation of human lactase persistence in Africa and Europe. *Nat Genet*. 2007;39:31–40.
30. Imtiaz F, Savilahti E, Sarnesto A, et al. The T/G_13915 variant upstream of the lactase gene (*LCT*) is the founder allele of lactase persistence in an urban Saudi population. *J Med Genet*. 2007;44:1–4.
31. Khabarova Y, Torniaainen S, Savilahti E, Isokoski M, Mattila K, Järvelä I. The -13914G>A variant upstream of the lactase gene (*LCT*) is associated with lactase persistence/non-persistence. *Scand J Clin Lab Invest*. 2010;70:354–357.
32. Xu L, Sun H, Zhang X, et al. The -22018A allele matches the lactase persistence phenotype in northern Chinese populations. *Scand J Gastroenterol*. 2010;45:168–174.
33. Mattar R, Monteiro MS, Silva JMK, Carrilho FJ. *LCT*-22018G>A single nucleotide polymorphism is a better predictor of adult-type hypolactasia/lactase persistence in Japanese-Brazilians than *LCT*-13910C>T. *Clinics (São Paulo)*. 2010;65:1399–1400.
34. Lewinsky RH, Jensen TGK, Møller J, Stensballe A, Olsen J, Troelsen JT. T-13910 DNA variant associated with lactase persistence interacts with Oct-1 and stimulates lactase promoter activity in vitro. *Hum Mol Genet*. 2005;14:3945–3953.
35. Enattah NS, Jensen TGK, Nielsen M, et al. Independent introduction of two lactase-persistence alleles into human populations reflects different history of adaptation to milk culture. *Am J Hum Genet*. 2008;82:57–72.
36. Olds LC, Ahn JK, Sibley E. -13915*G DNA polymorphism associated with lactase persistence in Africa interacts with Oct-1. *Hum Genet*. 2011;129:111–113.
37. Ingram CJE, Mulcare CA, Itan Y, Thomas MG, Swallow DM. Lactose digestion and the evolutionary genetics of lactase persistence. *Hum Genet*. 2009;124:579–591.
38. Gerbault P, Moret C, Currat M, Sanchez-Mazas A. Impact of selection and demography on the diffusion of lactase persistence. *PLoS One*. 2009;4:e6369.
39. Lagman JM, Rowland R. Activity of duodenal disaccharidases in relation to normal and abnormal mucosal morphology. *J Clin Pathol*. 1990;43:537–540.
40. Kuokkanen M, Myllyniemi M, Vauhkonen M, et al. A biopsy-based quick test in the diagnosis of duodenal hypolactasia in upper gastrointestinal endoscopy. *Endoscopy*. 2006;38:708–712.
41. Law D, Conklin J, Pimentel M. Lactose intolerance and the role of the lactose breath test. *Am J Gastroenterol*. 2010;105:1726–1728.
42. Jussila J. Diagnosis of lactose malabsorption by the lactose tolerance test with peroral ethanol administration. *Scand J Gastroenterol*. 1969;4:361–368.
43. Newcomer AD, McGill DB, Thomas PJ, Hofmann AF. Prospective comparison of indirect methods for detecting lactase deficiency. *N Engl J Med*. 1975;293:1232–1236.
44. Romagnuolo J, Schiller D, Bailey RJ. Using breath tests wisely in a gastroenterology practice: an evidence-based review of indications and pitfalls in interpretation. *Am J Gastroenterol*. 2002;97:1113–1126.
45. Mattar R, Monteiro MS, Villares CA, Santos AF, Carrilho FJ. Single nucleotide polymorphism C/T-13910, located upstream of the lactase gene, associated with adult-type hypolactasia: validation for clinical practice. *Clin Biochem*. 2008;41:628–630.
46. Büning C, Genschel J, Jurga J, et al. Introducing genetic testing for adult-type hypolactasia. *Digestion*. 2005;71:245–250.
47. Hogenauer C, Hammer HF, Mellitzer K, Renner W, Krejs GJ, Toplak H. Evaluation of a new DNA test compared with the lactose hydrogen breath test for the diagnosis of lactase non-persistence. *Eur J Gastroenterol Hepatol*. 2005;17:371–376.
48. Bodlaj G, Stöcher M, Hufnagl P, et al. Genotyping of the lactase-phlorizin hydrolase -13910 polymorphism by lightCycler PCR and implications for the diagnosis of lactose intolerance. *Clin Chem*. 2006;52:148–151.
49. Szilagyi A, Malolepszy P, Hamard E, et al. Comparison of a real-time polymerase chain reaction assay for lactase genetic polymorphism with standard indirect tests for lactose maldigestion. *Clin Gastroenterol Hepatol*. 2007;5:192–196.
50. Tag CG, Schiffers MC, Mohnen M, Gressner AM, Weiskirchen R. A novel proximal -13914G>A base replacement in the vicinity of the common-13910T/C lactase gene variation results in an atypical light cycler melting curve in testing with the MutaREAL lactase test. *Clin Chem*. 2007;53:146–148.
51. Torbjörn K, Olsson LA. Simultaneous genotyping of the three lactose tolerance linked polymorphisms *LCT*-13907C>G, *LCT*-13910C>T and *LCT*-13915T>G with Pyrosequencing technology. *Clin Chem Lab Med*. 2008;46:80–84.
52. Lomer MC, Parkes GC, Sanderson JD. Review article: lactose intolerance in clinical practice – myths and realities. *Aliment Pharmacol Ther*. 2008;27:93–103.
53. Matthews SB, Waud JP, Roberts AG, Campbell AK. Systemic lactose intolerance: a new perspective on an old problem. *Postgrad Med J*. 2005;81:167–173.
54. Harrington LK, Mayberry JF. A re-appraisal of lactose intolerance. *Int J Clin Pract*. 2008;62:1541–1546.
55. Campbell AK, Matthews SB, Vassel N, et al. Bacterial metabolic ‘toxins’: a new mechanism for lactose and food intolerance, and irritable bowel syndrome. *Toxicology*. 2010;278:268–276.
56. Minenna MF, Palieri A, Panella C, Ierardi E. Gastro-oesophageal reflux disease and lactose malabsorption: casual comorbidity or neglected association? *Dig Liver Dis*. 2006;38:437–438.
57. Suchy FJ, Brannon PM, Carpenter TO, et al. National Institutes of Health Consensus Development Conference: lactose intolerance and health. *Ann Intern Med*. 2010;152:792–796.
58. Shaukat A, Levitt MD, Taylor BC, et al. Systematic review: effective management strategies for lactose intolerance. *Ann Intern Med*. 2010;152:797–803.
59. Jellema P, Schellevis FG, van der Windt DA, Kneepkens CM, van der Horst HE. Lactose malabsorption and intolerance: a systematic review on the diagnostic value of gastrointestinal symptoms and self-reported milk intolerance. *Q J Med*. 2010;103:555–572.

60. Savaiano DA, Boushey CJ, McCabe GP. Lactose intolerance symptoms assessed by meta-analysis: a grain of truth that leads to exaggeration. *J Nutr*. 2006;136:1107–1113.
61. Casellas F, Varelam E, Aparici A, Casaus M, Rodríguez P. Development, validation, and applicability of a symptoms questionnaire for lactose malabsorption screening. *Dig Dis Sci*. 2009;54:1059–1065.
62. Casellas F, Aparici A, Casaus M, Rodríguez P, Malagelada JR. Subjective perception of lactose intolerance does not always indicate lactose malabsorption. *Clin Gastroenterol Hepatol*. 2010;8:581–586.
63. Erminia R, Ilaria B, Tiziana M, et al. HRQoL questionnaire evaluation in lactose intolerant patients with adverse reactions to foods. *Intern Emerg Med*. May 26, 2011. [Epub ahead of print.]
64. Colloca L, Miller FG. The nocebo effect and its relevance for clinical practice. *Psychosom Med*. 2011;73:598–603.
65. Vernia P, Di Camillo M, Foglietta T, Avallone VE, De Carolis A. Diagnosis of lactose intolerance and the “nocebo” effect: the role of negative expectations. *Dig Liver Dis*. 2010;42:616–619.
66. Di Stefano M, Corazza GR. The patient’s expectation during H₂ breath testing: don’t underestimate the reader’s expectation. *Dig Liver Dis*. 2011;43:86.
67. Novillo A, Peralta D, Dima G, Besasso H, Soifer L. Frequency of bacterial overgrowth in patients with clinical lactose intolerance. *Acta Gastroenterol Latinoam*. 2010;40:221–224.
68. Suarez FL, Levitt MD. An understanding of excessive intestinal gas. *Curr Gastroenterol Rep*. 2000;2:413–419.
69. Enattah N, Välimäki VV, Välimäki MJ, Löyttyniemi E, Sahi T, Järvelä I. Molecularly defined lactose malabsorption, peak bone mass and bone turnover rate in young Finnish men. *Calcif Tissue Int*. 2004;75:488–493.
70. Obermayer-Pietsch BM, Gugatschka M, Reitter S, et al. Adult-type hypolactasia and calcium availability: decreased calcium intake or impaired calcium absorption? *Osteoporos Int*. 2007;18:445–451.
71. Lovelace HY, Barr SI. Diagnosis, symptoms, and calcium intakes of individuals with self-reported lactose intolerance. *J Am Coll Nutr*. 2005;24:51–57.
72. Nicklas TA, Qu H, Hughes SO, et al. Self-perceived lactose intolerance results in lower intakes of calcium and dairy foods and is associated with hypertension and diabetes in adults. *Am J Clin Nutr*. 2011;94:191–198.
73. Bácsi K, Kósa JP, Lazáry A, et al. LCT 13910 C/T polymorphism, serum calcium, and bone mineral density in postmenopausal women. *Osteoporos Int*. 2009;20:639–645.
74. Enattah NS, Sulkava R, Halonen P, Kontula K, Järvelä I. Genetic variant of lactase-persistent C/T-13910 is associated with bone fractures in very old age. *J Am Geriatr Soc*. 2005;53:79–82.
75. Enattah N, Pekkarinen T, Välimäki MJ, Löyttyniemi E, Järvelä I. Genetically defined adult-type hypolactasia and self reported lactose intolerance as risk factors of osteoporosis in Finnish postmenopausal women. *Eur J Clin Nutr*. 2005;59:1105–1111.
76. Gugatschka M, Hoeller A, Fahrleitner-Pammer A, et al. Calcium supply, bone mineral density and genetically defined lactose maldigestion in a cohort of elderly men. *J Endocrinol Invest*. 2007;30:46–51.
77. Laaksonen MM, Impivaara O, Sievänen H, et al. Associations of genetic lactase non-persistence and sex with bone loss in young adulthood. *Bone*. 2009;44:1003–1009.
78. Tolonen S, Laaksonen M, Mikkilä V, et al. Lactase gene c/t(-13910) polymorphism, calcium intake, and pQCT bone traits in Finnish adults. *Calcif Tissue Int*. 2011;88:153–161.
79. Montalto M, Curigliano V, Santoro L, et al. Management and treatment of lactose malabsorption. *World J Gastroenterol*. 2006;12:187–191.
80. Montalto M, Gallo A, Santoro L, et al. Low-dose lactose in drugs neither increases breath hydrogen excretion nor causes gastrointestinal symptoms. *Aliment Pharmacol Ther*. 2008;28:1003–1012.

Clinical and Experimental Gastroenterology

Publish your work in this journal

Clinical and Experimental Gastroenterology is an international, peer-reviewed, open access journal, publishing all aspects of gastroenterology in the clinic and laboratory, including: Pathology, pathophysiology of gastrointestinal disease; Investigation and treatment of gastrointestinal disease; Pharmacology of drugs used in the alimentary tract;

Submit your manuscript here: <http://www.dovepress.com/clinical-and-experimental-gastroenterology-journal>

Dovepress

Immunology/genetics/genomics related to gastrointestinal disease. This journal is indexed on CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.