

## Research Paper

# Manganese provides antioxidant protection for sperm cryopreservation that may offer new consideration for clinical fertility

Ranjna S. Cheema,\* Amrit K. Bansal and Gurmail Singh Bilaspuri

Department of Animal Reproduction and Gynecology; Guru Angad Dev Veterinary and Animal Sciences University; Ludhiana, India

**Abbreviations:** EYC, egg yolk citrate; EYC-G, egg yolk citrate-glycerol; HOS, hypo-osmotic swelling; LPO, lipid peroxidation; MDA, malondialdehyde; ROS, reactive oxygen species; SOD, superoxide dismutase; TBA, thiobarbituric acid

**Key words:** cattle, lipid peroxidation, manganese, semen, oxidative stress

Reactive oxygen species (ROS) are generated by sperm metabolism. While, ROS are required for maturation, capacitation and acrosome reaction, they also modify many peroxidable cellular compounds. There is production of ROS during cryopreservation and frozen spermatozoa are highly sensitive to lipid peroxidation (LPO). Antioxidants exert a protective effect on the plasma membrane of frozen bovine sperm preserving both metabolic activity and cellular viability. Manganese ( $Mn^{++}$ ) is proved to be a chain breaking antioxidant in biological system. Therefore, we examined the role of ( $Mn^{++}$ ) during cryopreservation of cattle bull semen. Semen was divided into four parts and cryopreserved in egg-yolk-citrate extender + glycerol (EYC-G), EYC-G + 100  $\mu M$  of  $Mn^{++}$ , EYC-G + 150  $\mu M$  of  $Mn^{++}$  and EYC-G + 200  $\mu M$  of  $Mn^{++}$ . After four hours of cooling and 24 hrs of freezing, the spermatozoa were examined for percentage motility, Hypo-osmotic swelling (HOS), LPO and protein leakage. Addition of manganese to the semen during cryopreservation showed a protective effect and accounted for an increase in semen quality parameters [percentage motility, HOS percent and decrease in malondialdehyde (MDA) production and protein leakage]. The effect of manganese on motility and HOS was non-significant ( $p < 0.05$ ) in cooled spermatozoa but significant with 150  $\mu M$  of  $Mn^{++}$  in frozen-thawed spermatozoa. MDA production and protein leakage decreased to a significant and maximum level ( $p < 0.05$ ) on addition of 200  $\mu M$  of manganese. The addition of manganese to EYC-G dilutor will improve the quality/fertility of semen, which will result in improvement of in vitro fertilization and artificial insemination success rate.

## Introduction

It is generally accepted that the consequences of sperm cryoinjury caused by the cryopreservation are impaired transport and poor survival in the female reproductive tract.<sup>1</sup> The sperm plasma membrane is the primary site of damage induced by cryopreservation.<sup>2-4</sup> Both freezing and thawing cause tremendous alterations in cell water volume, which confer considerable mechanical stress on the cell membrane.<sup>2</sup> Although a significant physiological role of ROS during normal sperm function has been reported,<sup>5,6</sup> such as they facilitate fusogenicity of the spermatozoa, which enables them to bind to the zona pellucida, undergo the acrosome reaction through membrane lipid peroxidation and phospholipase-A activity,<sup>7</sup> traverse through the zona pellucida and fuse with oocyte membrane,<sup>8</sup> but when the balance between ROS production and detoxification by antioxidants is disrupted, an excess of ROS create oxidative stress. Unsaturated fatty acids, predominant in sperm membrane are susceptible to peroxidation<sup>9</sup> and leads to membrane damage, inhibition of respiration and leakage of intracellular enzymes.<sup>10</sup> LPO increases after cryopreservation in bull semen.<sup>11</sup>

Improvement in sperm quality upon addition of antioxidants to semen indicates indirect evidences for the damaging effects of ROS in sperm function. The scavenging potential of the ejaculate is normally maintained by adequate levels of antioxidants, present in seminal plasma. Sperm oxidative damage is the result of an improper balance between ROS generation and scavenging activities. In domestic animals, defective and dead spermatozoa have been identified as a major source of ROS generation during cryopreservation. Radical scavenging activity of manganese related to the rapid quenching of peroxy radicals has been demonstrated in the biological system viz, a typical antioxidative property of  $Mn^{++}$  may protect substantial nigra compact neurons from iron induced oxidative stress;<sup>12</sup> It could reduce the ferrous-ascorbate-mediated lipid peroxidation in placental membranes;<sup>13</sup> It induces an increase in the iron level, which provides direct evidence for Fe-mediated lipid peroxidation in the rat's brain that play an important role in the mechanism of Mn induced neurotoxicology;<sup>14</sup>

\*Correspondence to: Ranjna S. Cheema; Department of Animal Reproduction; Gynecology and Obstetrics; Guru Angad Dev Veterinary and Animal Sciences University; Ludhiana-141004; Punjab, India; Tel.: +91.9464416085; Email: ranjna\_cheema@hotmail.com

Submitted: 03/20/09; Revised: 04/21/09; Accepted: 04/21/09

Previously published online as an *Oxidative Medicine and Cellular Longevity* E-publication:  
<http://www.landesbioscience.com/journals/oximed/article/8804>

Mn-SOD transgene product protects frozen bone marrow cells/irradiated at 24°C;<sup>15</sup> Mn mew-tetrakas porphyrin also results in significantly blunt nitration of renal proteins during renal ischemia/reperfusion by blocking oxidant production and inducing ATP-synthase-beta subunit and prevention of ATP depletion.<sup>16</sup> Therefore, the present study evaluated the potential effect of Mn<sup>++</sup> in a dose dependent manner against cryopreservation induced damage to cattle bull spermatozoa. Spermatozoa were monitored for post thaw survival, HOS test, LPO and protein leakage.

## Results

**Effect of Mn<sup>++</sup> on motility and HOS.** Manganese added showed a protective effect and accounted for significantly higher values of motility and HOS-test response in cooled and frozen-thawed semen samples. A sperm motility declined to 63.3 ± 6.8% and 45 ± 2.4% in the cooled and frozen-thawed semen samples, diluted in EYC-G without Mn<sup>++</sup> respectively. Whereas the percentage of motile spermatozoa declined to 67.5 ± 2.0, 45.0 ± 2.4, 66.6 ± 5.9, 66.6 ± 3.0 and 48.3 ± 4.9, 58.3 ± 3.6, 51.6 ± 3.0 in the cooled and frozen thawed semen sample, supplemented with 100 µM, 150 µM, 200 µM of Mn<sup>++</sup> respectively (Table 1, Fig. 1).

There was 16.7% and 35% loss of sperm motility in the semen samples cryopreserved in EYC-G without Mn<sup>++</sup> after cooling and freezing-thawing respectively. But the loss of motility decreased to only 12.5%, 13.4%, 13.4% and 31.7%, 21.7%, 28.4% on addition of 100 µM, 150 µM, 200 µM of Mn<sup>++</sup> in the cooled and frozen-thawed semen samples respectively (Table 1 and Fig. 1). It indicates non-significant ( $p < 0.05$ ) effect of Mn<sup>++</sup> on percentage motility of cooled spermatozoa and a significant ( $p < 0.05$ ) effect on frozen-thawed spermatozoa, but 150 µM concentration of Mn<sup>++</sup> showed a maximum protective effect.

The percentage of HOS +ve spermatozoa was 55.7 ± 2.8% in the freshly diluted semen, which declined to 46.7 ± 2.8% and 19.17 ± 4.4% in the cooled and frozen-thawed semen samples, diluted in EYC-G without Mn<sup>++</sup> respectively. The percentage of HOS +ve spermatozoa further declined to 45.1 ± 5.1%, 43.0 ± 6.4%, 44.2 ± 3.1% in the cooled semen samples and 23.19 ± 3.9%, 30.1 ± 6.9%, 24.25 ± 5.2% in the frozen-thawed semen samples, supplemented with 100 µM, 150 µM, 200 µM of Mn<sup>++</sup> respectively (Table 2 and Fig. 2).

The percentage of HOS +ve spermatozoa declined to 8.97% in cooled semen samples, diluted in EYC-G without manganese. But this decline was increased to 10.60%, 12.70%, 11.50% in cooled semen samples supplemented with 100, 150 and 200 µM of Mn<sup>++</sup>. A percentage of 27.56% in HOS was observed in semen samples, cryopreserved in EYC-G without Mn<sup>++</sup>, whereas a decrease 21.91%, 12.84% & 19.95% in HOS +ve spermatozoa was observed in frozen thawed semen samples, supplemented with 100, 150 & 250 µM of Mn<sup>++</sup> (Table 2 and Fig. 2).

The results indicate non-significant ( $p < 0.05$ ) effect of Mn<sup>++</sup> on sperm membrane integrity during the cooling process and significant ( $p < 0.05$ ) effect during the freezing process, but to a maximum level on addition of 150 µM of Mn<sup>++</sup>.

**Effect of Mn<sup>++</sup> on lipid peroxidation and protein-leakage.** Only 45.0 ± 3.3 µmoles/10<sup>9</sup> cells of MDA were produced in the

**Table 1 Manganese increases sperm motility during cryopreservation of cattle bull semen**

Concentration of Mn <sup>++</sup> (µM)	Motility (Mean ± S.E.)			
	Pre-Freezing	Decrease in motility (%)	Post-Thaw	Decrease in motility (%)
Control	80.0 ± 0.00 <sup>a</sup>	-	-	-
0	63.3 ± 6.8 <sup>bc</sup>	16.7	45.0 ± 2.4 <sup>d</sup>	35
100	67.5 ± 2.0 <sup>c</sup>	12.5	48.3 ± 4.9 <sup>d</sup>	31.7
150	66.6 ± 5.9 <sup>c</sup>	13.4	58.3 ± 3.6 <sup>b</sup>	21.7
200	66.6 ± 3.0 <sup>c</sup>	13.4	51.6 ± 3.0 <sup>d</sup>	28.4

Superscripts (a, b, c and d) indicate the difference at 5% level of significance within the columns. The table depicts non-significant protective effect of Mn<sup>++</sup> on percentage motility of cooled spermatozoa and significant and significant ( $p \leq 0.05$ ) effect on that of frozen thawed spermatozoa. 150 µM concentration of Mn showed a maximum protective effect.

freshly diluted semen, which increased to 73.3 ± 1.6 and 122.5 ± 6.9 µmoles/10<sup>9</sup> cells in the cooled semen samples, diluted in EYC-G without Mn<sup>++</sup>. MDA production was reduced to 66.96 ± 2.0, 63.43 ± 4.2, 56.26 ± 1.6 µmoles/10<sup>9</sup> cells in the cooled semen samples and to 101.8 ± 1.4, 96.8 ± 7.7, 75.3 ± 2.2 in the frozen-thawed semen samples, supplemented with 100, 150 and 200 µM of Mn<sup>++</sup> respectively (Table 3 and Fig. 3).

It shows a significant increase in MDA production in the cooled as well as frozen-thawed spermatozoa irrespective of the addition of Mn<sup>++</sup> to EYC-G (Table 3). The results indicate an increase of 38.82% and 40.16% in the cooled and frozen thawed spermatozoa, diluted in EYC-G without Mn<sup>++</sup>. Comparatively the percentage of MDA production decreased to 30.82%, 29.08% and 19.96% in the cooled spermatozoa and to 36.05%, 34.47% and 25.18% in frozen-thawed spermatozoa, supplemented with 100 µM, 150 µM and 200 µM of Mn<sup>++</sup> respectively. However, progressive decline in MDA production with an increase in concentration of Mn<sup>++</sup> was observed. Anti-oxidant effect of Mn<sup>++</sup> was significant ( $p < 0.05$ ) in cooled as well as frozen-thawed semen samples but to a maximum level on addition of 200 µM of Mn<sup>++</sup>.

The total protein content of freshly diluted washed spermatozoa was 18.72 mg/10<sup>9</sup> spermatozoa. The protein content of cooled spermatozoa declined to 15.53 ± 1.8, 16.35 ± 1.4, 16.73 ± 8.4 and 16.84 ± 8.0 mg/10<sup>9</sup> cells, diluted in EYC-G, EYC-G + 100 µM, EYC-G + 150 µM and EYC-G + 200 µM of Mn<sup>++</sup> respectively. The protein content of frozen-thawed spermatozoa further declined to 8.65 ± 1.0, 10.71 ± 8.5, 9.68 ± 5.3 and 12.50 ± 6.2 mg/10<sup>9</sup> cells, cryopreserved in EYC-G, EYC-G + 100 µM, EYC-G + 150 µM and EYC-G + 200 µM of Mn<sup>++</sup> respectively (Table 4, Fig. 4).

There was 17.04% and 44.30% leakage of total sperm proteins in the cooled and frozen-thawed semen samples, diluted in EYC-G without Mn<sup>++</sup> (Fig. 5, Table 4). The leakage of total sperm proteins was reduced to 12.66%, 10.63% and 10.04% and 34.49%, 42.14% and 25.77% in cooled and frozen thawed semen, supplemented with 100 µM, 150 µM and 200 µM of Mn<sup>++</sup> respectively. The results indicate that the




Figure 1. Manganese increases sperm motility (% Mean  $\pm$  SE) during cryopreservation of cattle bull semen. The figure depicts non-significant protective effect of  $Mn^{++}$  on percentage motility of cooled spermatozoa and significant and significant ( $p \leq 0.05$ ) effect on that of frozen-thawed spermatozoa. 150  $\mu M$  concentration of  $Mn^{++}$  showed a maximum protective effect.

Table 2 Manganese increases HOS percentage (mean  $\pm$  S.E) during cryopreservation of cattle bull spermatozoa

Conc of $Mn^{++}$ ( $\mu M$ )	Spermatozoa cooled at 4°C		Frozen-thawed spermatozoa	
	HOS (%)	Difference in HOS (%) between control & cooled spermatozoa	HOS (%)	Difference in HOS (%) between cooled and frozen-thawed spermatozoa
Control	55.7 $\pm$ 2.8 <sup>a</sup>	-	-	-
0	46.7 $\pm$ 2.8 <sup>b</sup>	8.97	19.17 $\pm$ 4.4 <sup>c</sup>	27.56
100	45.1 $\pm$ 5.1 <sup>b</sup>	10.60	23.19 $\pm$ 3.9 <sup>c</sup>	21.91
150	43.0 $\pm$ 6.4 <sup>b</sup>	12.70	30.1 $\pm$ 6.9 <sup>d</sup>	12.84
200	44.2 $\pm$ 3.1 <sup>b</sup>	11.50	24.25 $\pm$ 5.2 <sup>c</sup>	19.95

Superscripts (a, b, c and d) indicate the difference at 5% level of significance within the columns. This table depicts non-significant ( $p \geq 0.05$ ) effect of  $Mn^{++}$  on sperm membrane integrity in cooled semen samples and significant ( $p \leq 0.05$ ) effect in frozen thawed semen samples, but to a maximum level on addition of 150  $\mu M$   $Mn^{++}$ .

addition of  $Mn^{++}$  to EYC-G could control the leakage of sperm membrane proteins to a significant level ( $p < 0.05$ ) both in cooled and frozen-thawed semen, but to a maximum level on addition of 200  $\mu M$  of  $Mn^{++}$ .

## Discussion

Much of cryopreservation sperm damage depends on the structural stability of the plasma membrane.<sup>17</sup> The structural re-organization of sperm head plasma membrane after cryopreservation appears to disrupt the ability of the sperm to interact normally with cells of the female genital tract. Cryopreservation in the presence of cryoprotectants (typically glycerol) is considered to be a more moderate treatment than cold shock. Since impaired sperm membrane function due to cryopreservation inevitably affects capacitation, acrosome reaction, penetration into zona, fusion and penetration into the oolema and so diminishes successful fertilization in vivo. That is why when equal number of motile spermatozoa are inseminated, the fertility of fresh semen is superior to that of frozen semen. Manganese is an element of great importance in the life cycle of plants/animals and it plays an essential role as an activator of various enzymatic systems. Antioxidant effect of  $Mn^{++}$  or its protective effect against LPO has been studied in various biological systems.<sup>12-14,18</sup> In various organisms, high intracellular manganese provide protection against oxidative damage through unknown pathways and recently it has been found that manganese for oxidative protection is provided by the Nramp transporters.<sup>19</sup> The potential role of manganese in evaluation of infertile males has also been reported.<sup>20</sup> Elbetieha et al.<sup>21</sup> also postulated that ingestion of high dose of manganese chloride by




Figure 2. Cattle bull spermatozoa showing progressive and non-progressive motility.

male and female mice causes some adverse effects on fertility and reproduction.  $Mn^{++}$  has a peroxy radical scavenging activity, which seems to be related to the rapid quenching of peroxy radicals according to the reaction  $R-OO + Mn^{++} + H(+) \rightarrow ROOH + Mn$ .<sup>22</sup>

**Effect of  $Mn^{++}$  on motility and HOS.** A subpopulation of sperm survives and maintains fertilizing capacity not withstanding an overall reduction in the percentage of motile and viable cells. However, computer assisted motility of bovine sperm, immediately after thawing is not correlated with fertility.<sup>23</sup> Similarly Lapointe et al.<sup>24</sup> suggested the beneficial effect of 0.1 mM  $MnCl_2$  for the maintainance of sperm motility without detrimental effects on mucus penetration and fertilizing ability and hypothesized that  $Mn^{++}$  would have an effect on sperm cyclase activity that lead to increased  $Ca^{++}$  concentration and motility. However, despite preservation of adequate motility, cryopreserved human sperm exhibit significant membrane damage as indicated by subnormal hypo-osmotic swelling tests.<sup>25</sup> During the present studies, although the post thaw motility ranged from 45.0% to 51.6% in all the experiments but the percentage of HOS +ve spermatozoa ranged only from 19.95% to 27.56%. It shows that even motile spermatozoa have damaged membrane. About 36–47% reduction in normal morphology i.e., motility, viability, mitochondrial function has been observed in frozen-thawed human spermatozoa.<sup>26</sup> It has also been reported that the release of oxidase from dead sperms in EYC reduces the motility and viability of remaining living bull spermatozoa.<sup>27</sup> Therefore, the addition of  $Mn^{++}$  to EYC-G resulted in significant improvement in the post-thaw motility and HOS of frozen cattle bull spermatozoa in a dose dependent manner, which was maximum on addition of 150  $\mu M$  of  $Mn^{++}$ . Similarly the addition of  $Mn^{++}$  to human washed spermatozoa resulted in stimulation of progressive motility in a time and dose dependent manner.<sup>28</sup> The positive effects of antioxidants vitamin E and  $Mn^{++}$  have been studied on buffalo and cattle bull spermatozoa incubated with lipid peroxidation catalysts.<sup>29,30</sup>

**Effect of  $Mn^{++}$  on lipid peroxidation and protein-leakage.** Although a significant negative correlation between the ROS and IVF fertilization rate has been found,<sup>31</sup> controlled quantities of

Table 3 **Manganese decreases the malondialdehyde production (mean  $\pm$  SE) during cryopreservation of cattle bull spermatozoa**

Conc of $Mn^{++}$ ( $\mu M$ )	Spermatozoa cooled at 4°C		Frozen-thawed spermatozoa	
	MDA produced $\mu M/10^9$ cells	Difference in MDA production (freshly diluted & cooled sperms)	MDA produced $\mu M/10^9$ cells	Difference in MDA production (cooled & frozen-thawed sperms)
Control	45.03 $\pm$ 3.3 <sup>a</sup>	-	-	-
0	73.33 $\pm$ 1.6 <sup>b</sup>	38.55	122.5 $\pm$ 6.9 <sup>e</sup>	40.16
100	66.96 $\pm$ 2.0 <sup>c</sup>	30.82	101.8 $\pm$ 1.4 <sup>f</sup>	36.05
150	63.43 $\pm$ 4.2 <sup>c</sup>	29.08	96.8 $\pm$ 7.7 <sup>g</sup>	34.47
200	56.26 $\pm$ 1.6 <sup>d</sup>	19.96	75.2 $\pm$ 2.2 <sup>h</sup>	25.18

Superscripts (a, b, c, d, e, f, g and h) indicate the difference at 5% level of significance within the columns. This table depicts significant antioxidant effect of  $Mn^{++}$  in cooled as well as frozen thawed semen samples but to a maximum level on addition of 200  $\mu M$  of  $Mn^{++}$ .

ROS have shown to be essential for the development of capacitation and hyper activation,<sup>32</sup> two physiological processes of the sperm that are necessary to ensure fertilization. The maintenance of a suitable ROS level is therefore essential for adequate sperm functionality.

When the balance between ROS production and detoxification by antioxidants is disrupted, an excess of ROS such as  $H_2O_2$ , SO etc., creates oxidative stress and results in membrane damage. The activation of an aromatic amino acid oxidase following the death of ram and bull spermatozoa has been identified as major source of ROS production in the semen of these animals.<sup>33</sup> It has been reported that LPO in bull spermatozoa increases after cryopreservation.<sup>11</sup> The cryopreservation of cattle bull spermatozoa also resulted in an increase in LPO in terms of MDA production irrespective of the addition of  $Mn^{++}$ . Frozen-thawed bull spermatozoa are more easily per oxidized than freshly ejaculated spermatozoa.<sup>34</sup> The supplementation of  $Mn^{++}$  could reduce the level of MDA




Figure 3. Manganese increases HOS percentage (Mean ± SE) during cryopreservation of cattle bull spermatozoa. The figure depicts non-significant ( $p \geq 0.05$ ) effect of  $Mn^{++}$  on sperm membrane integrity in cooled semen samples and significant ( $p \leq 0.05$ ) effect in frozen-thawed semen samples, but to a maximum level on addition of 150 µM of  $Mn^{++}$ .

Table 4 **Manganese decreases the leakage of total protein (mean ± SE) during cryopreservation of cattle bull spermatozoa**

Conc of $Mn^{++}$ (µM)	Spermatozoa cooled at 4°C		Frozen-thawed spermatozoa	
	Protein leakage µM/10 <sup>9</sup> cells	Increase in protein leakage as compared to control	Protein leakage µM/10 <sup>9</sup> cells	Increase in protein leakage as compared to cooling
Control	18.72 ± 3.6 <sup>a</sup>	-	-	-
0	15.53 ± 1.8 <sup>b</sup>	17.04	8.65 ± 1.0 <sup>c</sup>	44.30
100	16.35 ± 1.4 <sup>b</sup>	12.66	10.71 ± 5.5 <sup>cd</sup>	34.49
150	16.73 ± 5.4 <sup>b</sup>	10.63	9.68 ± 2.3 <sup>c</sup>	42.14
200	16.84 ± 5.0 <sup>b</sup>	10.04	12.50 ± 4.2 <sup>d</sup>	25.17

Superscripts (a, b, c, d, e, f, g and h) indicate the difference at 5% level of significance within the columns. This table depicts that the addition of  $Mn^{++}$  to EYC-G could reduce the leakage of total sperm proteins to a significant level both in cooled and frozen-thawed semen samples, which was maximum on addition of 200 µM of  $Mn^{++}$ .

production significantly in spermatozoa cooled at 4°C and frozen-thawed spermatozoa, but to a maximum level on addition of 200 µM of  $Mn^{++}$ . Inclusion of natural oxidants ( $\alpha$ -tocopherol and ascorbate) had a protective effect on metabolic activity and cellular viability of

cryopreserved bovine sperm.<sup>35,36</sup> Similarly  $Mn^{++}$  as an antioxidant showed a protective effect on cattle bull spermatozoa during cryopreservation.

The presence of antioxidant enzymes, SOD, glutathione peroxidase (GPx) and catalase in human;<sup>37</sup> bull<sup>38</sup> and ram<sup>39,40</sup> semen and the effect of semen dilution results in reducing their protective capacity. Marti et al.<sup>41</sup> indicated a 65% decrease in SOD activity of ram spermatozoa after freezing-thawing and addition of seminal plasma proteins with oleic-linoleic acid & vitamin E accounted for an increase in enzyme activity levels. In our study, sperm membrane damage due to oxidative stress during cryopreservation of cattle bull spermatozoa also resulted in protein leakage irrespective of the addition of  $Mn^{++}$ . However, the addition of  $Mn^{++}$  could prevent the leakage of proteins to a significant level in a dose dependent manner and the effect was maximum with 200 µM of  $Mn^{++}$ . Bilodeau et al.<sup>40</sup> also observed a reduction in glutathione levels of bull spermatozoa during cryopreservation. Literature indicate that the enzyme system comprising SOD, glutathione peroxidase/reductase and catalase function as a defense against lipid peroxidation in mammalian sperm and defect of these enzyme activities could produce a loss to cell function. It can be further postulated from our results that supplementation of  $Mn^{++}$  as an antioxidant to EYC-G during cryopreservation may maintain the enzyme system by its scavenging activity for free radicals or by reducing the oxidative stress and could reduce the effect of cryoinjury to cattle bull spermatozoa.




Figure 4. Manganese decreases the malondialdehyde (MDA) production (Mean  $\pm$  SE) during cryopreservation of cattle bull spermatozoa. The figure depicts significant antioxidant effect of  $Mn^{++}$  in cooled as well as frozen thawed semen samples but to a maximum level on addition of 200  $\mu M$  of  $Mn^{++}$ .

It can be concluded that 200  $\mu M$  concentration of  $Mn^{++}$  is an optimum dose, which can be added to EYC-G during cryopreservation of cattle bull spermatozoa to reduce the oxidative stress/ improve the quality of semen.

## Materials and Methods

**Procurement and evaluation of semen.** Freshly ejaculated semen was collected with the help of artificial vagina, immediately transported to the lab and evaluated for its mass activity. A drop of semen was placed on a clean glass slide and observed under microscope (10 x 10X) for waves, swirls and eddies. Grading was done on the basis of observations; Immotile/dead sperm (0), no waves but sperm movement (+), slow wave formation (++), relatively more wave formation with swirls (+++), waves with swirls and eddies (++++). The semen with only ++++ mass activity was used for freezing.

**Experiment design.** Three separate experiments were performed to evaluate the effect of adding different concentrations of  $Mn^{++}$  to EYC-G on the post thaw sperm survival performance (pre-freezing motility, post thaw motility, HOS-test, LPO, protein leakage) as follows:-

Experiment 1. Addition of 100  $\mu M$  of  $MnCl_2$ .

Experiment 2. Addition of 150  $\mu M$  of  $MnCl_2$ .

Experiment 3. Addition of 200  $\mu M$  of  $MnCl_2$ .

In all experiments treatments were compared to a sample without  $Mn^{++}$  and to a control (freshly diluted semen).

**Cryo-preservation of semen.** Immediately EYC-G in the ratio of 1:1 was added to the semen and kept at 37°C for 5–10 minutes. Semen with 80% initial motility was further diluted to 1:8 at a sperm concentration of about  $100 \times 10^6$  cells/ml. Diluted semen samples of experiments 1–3 and without  $Mn^{++}$  were kept at 4°C in a cold handling cabinet for four hours. After checking the pre-freezing motility, half of the semen was taken out to study various sperm functions. Second part of each experiment was cryopreserved using the straw freezing procedure. After 24 hours, semen was processed for post thaw motility and various sperm functions.

**Sperm concentration.** Sperm concentration was estimated spectrophotometrically by taking absorbance at 545 nm, a standard curve for which was prepared (Sperm concentration, calculated with haemocytometer versus absorbance at 545 nm.)

**Percentage motility.** Motility of pre-cooled and frozen-thawed semen was observed microscopically using CCTV. A total of 200 motile and non-motile sperms were observed on the monitor and percent of motile spermatozoa was calculated.

**HOS-test.**<sup>41</sup> HOS-test indicates the membrane integrity of the spermatozoa and damage is caused to the sperm membrane during cryopreservation. Therefore, HOS-test was performed to find out the protective effect of  $Mn^{++}$  on sperm damage. Briefly, 0.2 ml of semen was incubated with 1.0 ml of 100  $\mu M$  of HOS solution and 0.85% saline separately for 30 minutes. After 30 minutes spermatozoa with swollen and coiled tails were observed under microscope




Figure 5. Manganese decreases the leakage of total protein (Mean  $\pm$  SE) during cryopreservation of cattle bull spermatozoa.

at 10 x 40X. A total of 200 coiled/uncoiled sperms were counted and percent coiled spermatozoa were calculated. Percent of HOS positive spermatozoa was calculated by subtracting number of coiled sperms in normal saline from that in HOS solution.

**Membrane lipid peroxidation.** Oxidative stress is induced during cryopreservation of semen and as a result ROS are produced. Therefore, LPO was measured to find out an antioxidant effect of  $Mn^{++}$  on oxidative stress. Membrane LPO was estimated by the end point generation of MDA determined by the TBA test.<sup>42</sup> Briefly, extended spermatozoa [(40–70)  $\times$  10<sup>6</sup> cells] in 0.5 ml of cold 10% (wt/vol) chilled trichloroacetic acid to proliferate proteins. The precipitate was pelleted by centrifugation (6,000 rpm for 10 minutes) and 1 ml of 0.375% (wt/vol) TBA was added to the supernatant and kept in a boiling water bath at 100°C for 10 minutes. After cooling, the absorbance was read at 532 nm (UV-VIS spectrophotometer, systronics, India). MDA production was calculated by the formula:

$$\frac{[O.D. \text{ at } 532 \text{ nm} \times \text{Volume of assay mixture}]}{[\text{Molar extinction coefficient} \times \text{Volume of sample}]}$$

$$(1.56 \times 10^5 \text{ M}^{-1}\text{C}^{-1})$$

**Total proteins.** There is leakage of sperm membrane enzymes and total proteins during cryopreservation. Therefore, total sperm protein was measured to find out the protective effect of  $Mn^{++}$  on protein leakage. It was estimated by the standard method<sup>43</sup> using 4–7  $\times$  10<sup>6</sup> sperms/0.05 ml. Bovine serum albumin was run as a standard.

Increase in MDA production and protein leakage due to cooling and freezing-thawing of spermatozoa was calculated by subtracting the values of control from that of cooled and of cooled samples from that of frozen-thawed ones respectively.

**Statistical analysis.** The data were analyzed by using computerized soft ware programmed for analysis of RBD (Randomized block design).

#### Acknowledgements

The authors are thankful to the semen freezing lab, Department of Animal Breeding and Genetics and GADVASU for providing the semen samples.

#### References

- Salmon S, Maxwell WMC. Frozen storage of ram semen II. Cause of low fertility after cervical insemination and methods of improvement. *Anim Reprod Sci* 1995; 38:1-36.
- Hammerstedt RH, Graham JK, Nolan JP. Cryopreservation of mammalian sperm: what we ask them to survive. *J Androl* 1990; 11:73-88.
- Parks JE, Graham JK. Effects of cryopreservation procedures on sperm membrane. *Theriogenol* 1999; 51:461-72.
- Watson PE. Recent developments and concepts in cryopreservation of spermatozoa and assessment of their post-thawing functions. *Reprod Fertil Dev* 1995; 7:871-91.
- Kodama H, Kuribayashi Y, Gagnon C. Effect of sperm lipid peroxidation on fertilization. *J Androl* 1996; 17:151-7.
- de Lamirande E, Jiang H, Zini A, Kodama H, Gagnon C. Reactive oxygen species and sperm physiology. *Rev Reprod* 1997; 2:48-54.
- Awda BJ, Machenzie-Bell M, Buhr MM. Reactive oxygen species and boar sperm function. *Biol Reprod* 2009; In press.
- Michael AJ, Alexopoulos C, Pontiki EA, Hadjippavlou-Lintina DJ, Saratsis P, Ververdis HN, et al. Effect of antioxidant supplementation in semen extenders on semen quality and reactive oxygen species of chilled canine spermatozoa. *Anim Reprod Sci* 2009; 112:119-35.
- Halliwell B, Gutteridge J. Lipid peroxidation, oxygen radicals, cell damage and antioxidant therapy. *Lancet* 1984; 1:1396-8.
- White IG. Lipids and calcium uptake of sperm in relation to cold shock and preservation: a review. *Reprod Fertil Dev* 1993; 5:639-58.
- Slaweta R, Wasowicz W, Laskowska T. Selenium content, glutathione peroxidase activity and lipid peroxide level in fresh bull semen and its relationship to motility of spermatozoa after freezing and thawing. *Zentralb Veterinarmed A* 1988; 35:455-60.
- Srizaki I, Mohanakumar KP, Rauhala P, Kim HG, Yeh KJ, Chiueh CC. Manganese: a transition metal protects nigrostriatal neurons from oxidative stress in the iron-induced animal model of Parkinsonism. *Neurosci* 1998; 85:1101-11.

13. Anand RK, Kanwar U. Role of some traces metal ions in placental membrane lipid peroxidation. *Biol Trace Elem Res* 2001; 82:61-75.
14. Chen MT, Cheng GW, Lin CC, Chen BH, Huang YL. Effects of acute manganese chloride exposure on lipid peroxidation and alteration of trace metal in rat brain. *Biol Trace Elem Res* 2006; 110:163-78.
15. Epperly MW, Epperly LD, Min Y, Wang H, Zhang X, Franicola D, et al. Overexpression of the MnSOD transgene product protects cryopreserved bone marrow hematopoietic progenitor cells from ionizing radiation. *Radiant Res* 2007; 168:560-6.
16. Saba H, Ines BH, Shambar M, Tanecia M, Cheryl L, Judit M, et al. Manganese porphyrin reduces renal injury and mitochondrial damage during ischemia/reperfusion. *Free Radic Bio Med* 2007; 42:1571-8.
17. de Leeuw FE, de Leeuw AM, DenDass JH, Colenbrander B, Verkleij AJ. Effects of various cryoprotective agents and membranestabilizing compounds on bull sperm membrane integrity after cooling and freezing. *Cryobiol* 1993; 3:32-44.
18. Campanella L, Gatta T, Ravera O. Relationship between anti-oxidant capacity and manganese accumulation in the soft tissues of two fresh water mollusks: *Unio pictorum mancus* (Lamellibranchia, Unionidae) and *viviparus ater* (Gastropoda, Prosobranchia). *J Limnol* 2005; 64:153-8.
19. Reddi AR, Jensen LT, Naranuntarat A, Rosenfeld L, Leung E, Shah R, et al. The overlapping roles of manganese and Cu/Zn SOD in oxidative stress protection. *Free Radic Biol Med* 2009; 46:154-62.
20. Adejuwon CA, Ilesanmi AO, Ode EO, Akinlade KS. Biophysical and biochemical analysis of semen in infertile Nigerian males. *Afr J Med Sci* 1996; 25:217-9.
21. Elbetieha A, Bataineh H, Darmani H, Al-Hamood MH. Effects of long term exposure to manganese chloride on fertility of male and female mice. *Toxicol letters* 2001; 119:193-201.
22. Coassin M, Ursini F, Bindoli A. Antioxidant effect of manganese. *Arch Biochem Biophys* 1992; 299:330-3.
23. Check ML, Check JH. Poor hypoosmotic swelling test results from cryopreserved sperm despite preservation of sperm motility. *Arch Androl* 1991; 26:37-41.
24. Lapointe S, Ahmad I, Bhur MM, Sirard MA. Modulation of post-thaw motility, survival, calcium uptake and fertility of bovine sperm by magnesium and manganese. *J Dairy Sci* 1996; 79:2163-9.
25. Bailey JL, Robertson L, Buhr MM. Calcium regulation, computerized motility parameters and the fertility of bovine spermatozoa. *Can. J Anim Sci* 1994; 74:53-8.
26. Cornell MO, McClure N, Lewis SEM. The effect of cryopreservation on sperm morphology, motility and mitochondrial function. *Hum Reprod* 2002; 17:704-9.
27. Shanon P, Curron B. Toxic effect and action of dead sperm on dilute bovine semen. *J Dairy Sci* 1972; 55:614-20.
28. Magnus O, Brekke I, Abyholm T, Purvis K. Effect of manganese and other divalent cations on progressive motility of human sperm. *Arch Androl* 1990; 24:159-66.
29. Singh P, Chand D, Georgie GC. Effect of vitamin E on lipid peroxidation in buffalo *Bubalus bubalis* L. *Ind J Exp Biol* 1989; 27:14-6.
30. Bilaspuri GS, Bansal AK. Mn<sup>2+</sup>: A potent antioxidant and stimulator of sperm capacitation and acrosome reaction in crossbred cattle bulls. *Arch Anim Breed* 2008; 51:149-58.
31. Aggarwal A, Allamaneni S, Nallella K, George A, Mascha E. Correlation of reactive oxygen species levels with the fertilization rate after in vivo fertilization rate after in vitro fertilization: a qualified meta-analysis. *Fertil Steril* 2005; 84:228-31.
32. de Lamirande E, Gagnon C. A positive role for the superoxide anion in triggering hyperactivation and capacitation of human spermatozoa. *Int J Androl* 1993; 16:21-5.
33. Upreti GC, Jensen K, Munday R, Dugan Zich DM, Vishwanath R, Smith JF. Studies on aromatic amino acid oxidase activity in ram spermatozoa: Role of pyruvate as an antioxidant. *Anim Reprod Sci* 1998; 51:275-87.
34. Trincherro GD, Affranchino MA, Schang LM, Beconi MT. Antioxidant effect of bovine spermatozoa on lipid peroxidation. *Comp Biol* 1990; 8:339-50.
35. Beconi MT, Affranchino MA, Schang LM, Beorlegui NB. Influence of antioxidants on SOD activity in bovine sperm. *Biochem Int* 1991; 23:545-53.
36. Beconi MT, Francia CR, Mora NG, Affranchino MA. Effect of natural antioxidants on frozen bovine semen preservation. *Theriogenol* 1993; 40:841-51.
37. Mann T, Mann CL. Male reproductive function and semen. Springer-Verlag, New York 1981; 195-268.
38. Marti E, Marti JI, Muino Balanco T, Cebrian-Perez JA. Effect of the cryopreservation process on the activity and immunolocalization of antioxidant enzymes in ram spermatozoa m. *J Androl* 2008; 1:25.
39. Marti E, Mara L, Marti J, Muino Blanco T, Cebrian-Perez J. Seasonal variations in antioxidant enzyme activity in ram seminal plasma. *Theriogenol* 2007; 67:1446-54.
40. Bilodeau JF, Chatterjee S, Sirard MA, Gagnon C. Level of antioxidant defenses are decreased in bovine spermatozoa after a cycle of freezing and thawing. *Mol Reprod Dev* 2000; 55:282-8.
41. Jeyendran R, Vandervan, Znebel L. The hypo-osmotic swelling test an update. *Arch Androl* 1992; 29:115-6.
42. Buege JA, Steven AD. In: Colowick SP, Kaplan NO, (Eds.) *Methods in Enzymology* 52 (Biomembranes: Part C: Biological oxidants; Fleischers, Pecker L, Eds.,) Academic Press, New York 1978; 302-10.
43. Lowry RR, Tansly TJ. Rapid colorimetric determination of free fatty acids. *J Am Oil Chem Soc* 1976; 53:470-2.





# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

