

Natural hydroxyanthraquinoid pigments as potent food grade colorants: an overview

Yanis CARO,^{a,b,*} Linda ANAMALE,^{a,b} Mireille FOUILLAUD,^{a,b} Philippe LAURENT,^{b,c} Thomas PETIT,^{b,c} and Laurent DUFOSSE^{a,b}

^aDépartement Agroalimentaire, ESIROI, Université de La Réunion, Sainte-Clotilde, Ile de la Réunion, France

^bLCSNSA, Faculté des Sciences et des Technologies, Université de La Réunion, Sainte-Clotilde, Ile de la Réunion, France

^cDépartement Génie Biologique, IUT, Université de La Réunion, Saint-Pierre, Ile de la Réunion, France

Received 24 October 2012; Accepted 12 November 2012

© The Author(s) 2012. This article is published with open access at Springerlink.com

Abstract: Natural pigments and colorants are widely used in the world in many industries such as textile dyeing, food processing or cosmetic manufacturing. Among the natural products of interest are various compounds belonging to carotenoids, anthocyanins, chlorophylls, melanins, betalains... The review emphasizes pigments with anthraquinoid skeleton and gives an overview on hydroxyanthraquinoids described in Nature, the first one ever published. Trends in consumption, production and regulation of natural food grade colorants are given, in the current global market. The second part focuses on the description of the chemical structures of the main anthraquinoid colouring compounds, their properties and their biosynthetic pathways. Main natural sources of such pigments are summarized, followed by discussion about toxicity and carcinogenicity observed in some cases. As a conclusion, current industrial applications of natural hydroxyanthraquinoids are described with two examples, carminic acid from an insect and Arpink red™ from a filamentous fungus.

Keywords: anthraquinone, hydroxyanthraquinone, natural colorant, food colorant, microbial pigment, biotechnology, mycotoxin contamination

Introduction

Food grade colorants can loosely be categorized as ‘*natural*’ or ‘*synthetic*’. The term ‘natural colorants’ indicates that the source of the colorant is natural even if varying definitions and regulations exist according to the country in question. For example, under the United States (US) Food and Drug Administration (FDA) regulations, a colorant added to a food product cannot be considered as ‘natural’, no matter what the source is; unless the colorant is natural to the food product itself. The FDA regulates the natural and synthetic colorants of food applications in two classes. In general, the synthetic colorants (that do not exist in nature) are subjected to a certification requirement to assure that each batch of material manufactured meets the standard specifications, while natural colorants are “exempt from certification” and may be manufactured and marketed without certification of FDA (no US Food, Drug and Cosmetic Act (FD&C)-number). In contrast, E-numbers are used for all colorants for food applications in the European Union (EU). Colorants for food applications listed by both the FDA and the EU are tested for biosafety before their promotion and commercialization, and are further controlled by national legislation specifying those

colorants that may be used, the type of food that may be coloured, the quantity that may be added and the limit of maximum daily intake.

For a very long time, the use of food colorants focused on synthetic ones. However, over the last few decades, synthetic colorants tend to be perceived as undesirable by consumers, due to the harmful effects of some synthetic pigments on human health, including allergic reactions, mutagenicity and potential carcinogenicity (e.g. skin cancer)¹. Many manufacturers have considered replacing synthetic colorants in their food products with natural colouring alternatives in response to pressure from both customers and regulators². Whereas 43 colorants were authorized in the EU as food additives in 1994, actually almost a hundred of food grade colorants are authorized in the EU and have been assigned by an ‘E-number’; almost 40% of these were of natural origin^{1,2}. These natural colorants are usually applied in several industrial food processes for the same reasons as the synthetic counterparts: (i) to enhance the product’s natural colour whose ingredients are unable to provide a sufficient colour; (ii) to standardize the colour and appearance of product like confectionery; (iii) to restore what has been lost during processing or (iv) to add a novel sensory aspect that attracts customers.

Among the natural pigments of interest are various compounds belonging to carotenoids, anthocyanins,

*To whom correspondence should be addressed. E-mail: yanis.caro@univ-reunion.fr

chlorophylls, melanins, betalains, quinones... This review emphasizes pigments with anthraquinoid skeleton and gives an overview on hydroxyanthraquinoid pigments which are widely present in Nature and are gaining increasing interest by academics and people from the industry.

1 Trends in consumption, production and regulation of natural food grade colorants in the current global market

Currently, the natural food colouring industry market is growing 10%–15% annually. Natural varieties share of the global food colorant market increased from about 31% in 2005 to 36% in 2009. The current consumer preference for natural food grade colorants is associated with their image of being healthy and of good quality. According to a report from Leatherhead Food Research, Shaun Weston mentions that the global market for food grade colorants is expected to reach \$1.6 billion USD by 2015, up to 10% from its present levels and fuelled mainly by the growth in natural colorants and colouring foodstuffs (data from Leatherhead Food International LFI) (www.leatherheadfood.com). The main industrial technology used for the production of natural colorants for food applications depends on the extraction of coloured pigments from edible plants, fruits or vegetables. Table 1 shows the main natural food grade colorants authorized and currently available in the current global market. Common natural colorants include turmeric, curcumin, annatto, paprika, caramel and cochineal extract. Natural colorants are often commercially available in powder, oil-soluble emulsion, or water-soluble emulsion forms.

Table 1. Main natural food grade colorants authorized and currently available in the current global market

Color/shade	E-number*	Natural colorant	Chemical category
<i>From plants, fruit or vegetables:</i>			
Yellow	E100, E100 (i)	Curcumin	Curcuminoid
Yellow	E100 (ii)	Turmeric	Curcuminoid
Green	E140	Chlorophylls	Tetrapyrrole
Green	E141	Chlorophyllins	Tetrapyrrole
Brown	E150a–d	Caramel	Melanoidin
Orange-yellow	E160a (i)	Mixed Carotenes	Carotenoid
Orange-yellow	E160a (ii)	β-carotene	Carotenoid
Yellow to orange	E160b	Annatto	Carotenoid
Yellow to orange	E160b (i)	Annato (Bixin)	Carotenoid
Yellow to orange	E160b (ii)	Annato (Norbixin)	Carotenoid
Red	E160c	Paprika (Capsanthin)	Carotenoid
Yellow to red	E160d	Lycopene	Carotenoid
Yellow to red	E160e	Apocarotenal	Carotenoid
Orange-yellow	E161a	Flavoxanthin	Carotenoid
Orange-yellow	E161b	Lutein	Carotenoid
Orange-yellow	E161d	Rubixanthin	Carotenoid
Orange-yellow	E161e	Violaxanthin	Carotenoid
Orange-yellow	E161f	Rhodoxanthin	Carotenoid
Orange, Red	E161h	Zeaxanthin	Carotenoid
Red	E162	Red Beet Juice	Betalain
Red, Blue or Violet	E163a	Cyanidin	Anthocyanin
Red, Blue or Violet	E163e	Peonidin	Anthocyanin

From animal:

Yellow	E101, E101a	Riboflavin	Flavin
Magenta-red	E120 (ii)	Carmine acid (Cochineal extract)	Anthraquinone
Orange-yellow	E161c	Cryptoxanthin	Carotenoid
Orange, Red	E161g	Canthaxanthin	Carotenoid

From microorganisms:

Yellow	E101 (iii)	Riboflavin (from <i>Bacillus subtilis</i>) Other sources: <i>Ashbya gossypii</i> , <i>Candida guilliermondii</i> , <i>Clostridium acetobutylicum</i> and <i>Debaryomyces subglabrous</i>	Flavin
Orange-yellow	E160a (ii)	β-carotene (from <i>Blakeslea trispora</i>)	Carotenoid
Orange-yellow	E160a (iv)	β-carotene (from <i>Dunaliella salina</i>) Other sources: <i>Dunaliella bardawil</i>	Carotenoid
Yellow to red	E160d (iii)	Lycopene (from <i>Blakeslea trispora</i>)	Carotenoid
Yellow to red	E-161j	Astaxanthin (from <i>Haematococcus pluvialis</i>) Other sources: <i>Haematococcus lacustris</i> , <i>Xanthophyllomyces dendrorhous</i>	Carotenoid
Orange, Red	E161g	Canthaxanthin (from <i>Haematococcus lacustris</i>) Other sources: <i>Bradyrhizobium sp.</i>	Carotenoid

*E-number of the corresponding authorized food colorant in the European Union

Many scientific papers describe the extraction, characterization and properties of natural pigments from fruits, vegetables, lichens and marine life^{1,3}. However, the potential of these renewable resources as sources for new commercial natural food grade colorants would still be limited both by the manufacturing costs and the availability of the raw material, which would need to be cultivated in sufficient quantities for industrial extraction. The microbial pigment production by biotechnology would have the advantage of producing higher yields. This kind of pigment production is not at all dependent on the availability and external supply of particular raw materials. In addition, microbial pigments are often more stable and water-soluble than those of plant sources^{4,5}. The really first European success story in pigment production using a microorganism is β -carotene (additive E-160a(ii); orange-yellow pigment) from the fungus *Blakeslea trispora* by DSMTM. Among microalgae, some successful stories yield to efficient production of carotenoids using *Dunaliella salina* (e.g., β -carotene, additive E-160a(iv)) or *Haematococcus pluvialis* (e.g., astaxanthin, additive E-161j; yellow to red pigment)⁶ (Table 1). Nowadays, fermentative productions of natural food grade colorants are available in the global market⁷. This approval of microbial carotenoids as food colorants has strengthened the prospects for new natural colorants².

However, some new microbial pigments might not be accepted if they were to be introduced into industrial food manufacturing today⁸. The commercially available *Monascus* pigments are a perfect example. These fungal pigments are natural azaphilone pigment mixtures. The red colorant obtained is produced commercially using strains of *Monascus* fungi in the Orient for centuries used as a food colorant for making red rice wine, red soybean cheese, meat and marine

products. However, *Monascus* pigments are still not allowed as a food additive in either the US or the EU; there have been controversial views presented over their safe use⁸. This could be essentially because of the presence of the mycotoxin citrinin (yellow compound) and some other potential toxic metabolites which may occur in some batches with *Monascus* fungi^{9–11}. The production of citrinin limits the commercial use of *Monascus* fungi as producers of natural food grade colorants¹². Research continues on new azaphilone pigments produced from non-mycotoxigenic fungal strains, such as *Epicoccum nigrum*, *Penicillium aculeatum* or *P. pinophilum*—that are incapable of co-producing citrinin—in the prospects for new natural food grade colorants^{8,13}. The case of the fungal Arpink red™ colorant, *i.e.* a natural food colorant manufactured by the Czech company, Ascolor Biotech s.r.o. is also atypical. This company has produced a chromophore of the anthraquinoid type as a natural food colorant, by fermentation and bioprocess engineering using the strain *Penicillium oxalicum* var. *Armeniaca* CCM 8242 obtained from soil (the variety was never formally described). The Arpink red™ colorant has received a two-year temporary approval by the EU for distribution as a food additive, exclusively in the Czech Republic from 2004 to 2006. The extraction, isolation and characterization of natural anthraquinoid pigments have been also reported from other filamentous fungi with different shades such as red, reddish brown, bronze and maroon.

2 Natural hydroxyanthraquinoid pigments: chemical structures of the main colouring components, their properties and their biosynthetic pathways

Anthraquinones are a class of compounds of the quinone family that consists of several hundreds of compounds that differ in the nature and positions of substituent groups. Anthraquinone derivatives are derivatives of the basic structure 9,10-anthracenedione or also called 9,10-dioxoanthracene, *i.e.* a tricyclic aromatic organic compound with formula C₁₄H₈O₂ and whose ketone groups are on the central ring in position C-9 and C-10. Figure 1 shows the skeleton structure of anthraquinoid derivatives. In general, for each anthraquinoid derivative there are eight possible hydrogens that can be substituted. The term ‘hydroxyanthraquinoid (HAQN)’ derivatives usually refers to derivatives of 9,10-hydroxyanthraquinone, *i.e.* derivatives of 9,10-anthraquinone where any number *n* of hydrogen atoms have been replaced by *n* hydroxyl (-OH) groups. In this case the number *n* of hydroxyl group is indicated by a multiplier prefix (mono-, di-, tri-, up to octa-). The HAQN derivatives absorb visible light and are coloured, whereas strictly 9,10-anthraquinone derivatives are colorless like tectoquinone¹⁴.

Most HAQN colour compounds of natural origin have




Figure 1. The skeleton structure of anthraquinoid derivatives

complex structures with several functional groups, which modify their absorption spectra. The chemical structures of natural HAQN pigments and their main physical and functional properties^{15–52} are shown on Table 2. In the UV region, substituted 9,10-anthraquinone derivatives show intense benzenoid absorption bands fairly regularly within the ranges 240–260 and 320–330 nm. The quinonoid bands appear in a range from 260 to 290 nm and 9,10-hydroxyanthraquinone derivatives show an absorption band between 220 and 240 nm. The HAQN derivatives have attracted the attention of many researchers due to their large list of possible applications related to their interesting photoactivity and more particularly based on their chromatic properties. They possess good light-fastness properties, which makes metallization unnecessary. HAQN derivatives can form coordination complexes with several cations. They are relatively stable and the advantage of pigments of HAQN-type compared to azo pigments is their superior brightness. Moreover, ionization of a hydroxylic group results in a bathochromic shift. It appears that the colour of the HAQN pigments depends on the position and number of the hydroxyl substituents in the different rings⁵³.

Natural HAQN pigments are produced by the secondary metabolism of organisms. One of the remarkable features of natural HAQN biosynthesis is that they are derived from a variety of different precursors and pathways. There are at least two biosynthetic pathways leading to HAQN pigments. On one hand, the most important is the polyketide pathway (acetate-malonate pathway) that includes suitable folding and condensation of an octaketide chain derived from acetate (acetyl-CoA) and malonate (malonyl-CoA) units^{3,32,54}. The resultant polycarbonyl compounds serve as substrates for various cyclases that produce aromatic compounds that represent typical fungal metabolites. Natural HAQN pigments that are synthesized following this acetate-malonate pathway (see Figure 2) always show a characteristic substitution pattern, *i.e.* they show substitution on both aromatic rings and more particularly at least one hydroxyl group in position R1 and one hydroxyl or methoxyl (-OCH₃) group in position R8: examples being emodin, physcion, endocrocin, dermolutein, dermorubin, dermocybin and dermocybin. According to this polyketide pathway, the biosynthetic relationships show that the yellow compounds (e.g., emodin, physcion, endocrocin and dermolutein) exist in the beginning of the synthesis pathway whereas the red compounds like dermorubin and dermocybin are more complicated in structure and occur in the latter part of the biosynthesis pathway³². More recently, Bringmann et al.⁵⁵ revealed that the pigment chrysophanol is shown to be formed, in an organism-specific way, by a third folding mode involving a remarkable cyclization of a bicyclic diketo precursor, thus establishing the first example of multiple convergence in polyketide biosynthesis.

On the other hand, HAQN pigments are formed *via* the shikimate or chorismate/*o*-succinylbenzoic acid pathway (Fig. 2). HAQN pigments that are synthesized *via* this pathway only have one of the rings unsubstituted and at least one hydroxyl group in position R1 on the ring C. The rings A and B are derived from chorismate and α -ketoglutarate *via* *o*-succinylbenzoic acid (the HAQN biosynthesis branch at 1,4-dihydroxy-2-naphthoic acid), whereas ring C is formed from isopentenyl diphosphate either formed *via* the mevalonic acid pathway or the 2-C-methyl-D-erythritol 4-phosphate pathway^{54,56,57}. The relevant colouring compounds are alizarin

Table 2. Chemical structures, physical and functional properties of natural hydroxyanthraquinoid (HAQN) pigments

HAQN dye (and it's abbreviation)	Chemical structure	Physical and functional properties	References
Monohydroxyanthraquinones (OH)₁:			
Damnacanthal 3-hydroxy-1-methoxy-anthraquinone-2-carboxaldehyde		C ₁₆ H ₉ O ₄ (OH) ₁ – Mw: 282 Mp: 210–211 °C Shade: pale yellow	(15, 16)
Lucidin primeveroside 3-O-primeverose-1-hydroxy-2-hydroxymethyl-anthraquinone (LuP)		C ₂₆ H ₂₇ O ₁₃ (OH) ₁ – Mw: 564 Mp: 210–212 °C Shade: red	(17)
Pachybasin 2-methyl-1-hydroxy-anthraquinone		C ₁₅ H ₉ O ₂ (OH) ₁ – Mw: 238 Mp: 176 °C; shade: yellow UV (EtOH) λ _{max} 403, 281, 252, 224 nm; IR (KBr) ν _{max} 3083, 2967, 2933, 2862, 1677, 1642, 1593, 1578, 1403, 1380, 1338, 1311, 1290, 1235, 1148, 1068, 1039, 1002, 929, 870, 805, 793, 769, 712 cm ⁻¹ ; ¹H NMR (CDCl ₃ , 500 MHz) δ 12.67, 8.20, 7.74, 7.55, 7.17, 2.45.	(18)
Ruberythric acid (= alizarin primeveroside) 2-O-primeverose-1-hydroxy-anthraquinone (Rba)		C ₂₅ H ₂₅ O ₁₂ (OH) ₁ – Mw: 534 Mp: 259–261 °C Shade: yellow Soluble in hot water Slightly soluble in alcohols and ether Insoluble in benzene	(17, 19)
- Dihydroxyanthraquinones (OH)₂:			
2-(1-hydroxyethyl)-3,8-dihydroxy-6-methoxy-anthraquinone		C ₁₇ H ₁₂ O ₄ (OH) ₂ – Mw: 314 Mp: 208–212 °C; shade: orange UV (MeOH) λ _{max} 466, 341, 305, 285, 219 nm; IR (KBr) ν _{max} 3420, 1667, 1630, 1580, 1480, 1440, 1390, 1360, 1305, 1270, 1240, 1215, 1160, 1080, 1030, 1005, 970, 930, 900, 875, 835, 800 cm ⁻¹ ; ¹H NMR (CDCl ₃ , 500 MHz) δ 12.96, 7.94, 7.76, 7.35, 6.70, 5.20, 3.93, 1.68.	(20, 21)
2-acetyl-3,8-dihydroxy-6-methoxy-anthraquinone		C ₁₇ H ₁₀ O ₄ (OH) ₂ – Mw: 312 Mp: 256–270 °C; shade: yellow UV (MeOH) λ _{max} 420, 348, 307, 280, 233 nm; IR (KBr) ν _{max} 1672, 1655, 1630, 1575, 1480, 1450, 1410, 1380, 1300, 1250, 1205, 1170, 1125, 1025, 970, 935, 905, 890, 865, 830, 805, 755, 745, 715, 695 cm ⁻¹ ; ¹H NMR (CDCl ₃ , 500 MHz) δ 12.91, 12.72, 8.76, 7.81, 7.39, 6.74, 2.81.	(20, 21)
Alizarin 1,2-dihydroxy-anthraquinone (Ali)		C ₁₄ H ₆ O ₂ (OH) ₂ – Mw: 240 Mp: 278–280 °C Shade : Yellow to red (acid); red to violet (alkaline) Soluble in water, in NaOH 1M and in alcohols UV (MeOH) λ _{max} 609, 567, 429, 422 nm.	(22, 23)
Aloe-emodin 3-hydroxymethyl-1,8-dihydroxy-anthraquinone (Ale)		C ₁₅ H ₈ O ₃ (OH) ₂ – Mw: 270 Mp: 223–224 °C Shade: orange-yellow	(24, 25, 26)





Anthraflavic acid (= anthraflavin) 2,6-dihydroxy-anthraquinone (Afv)		$C_{14}H_6O_3(OH)_2$ – Mw: 240 Mp ≥ 320 °C Shade: yellow	(19)
Arpink red™		$C_{22}H_{13}O_4(OH)_1$ Shade: red	(2, 5, 15)
Austrocortinin 2-methoxy-7-methyl-1,4-dihydroxy-anthraquinone		$C_{16}H_{10}O_3(OH)_2$ – Mw: 284 Shade: red	(27)
Chrysophanol (=chrysophanic acid) 3-methyl-1,8-dihydroxy-anthraquinone (Chr)		$C_{15}H_8O_2(OH)_2$ – Mw: 254 Mp: 186 °C Shade: orange-yellow	(18, 28-31)
Danthron (= dantron or chrysazin) 1,8-dihydroxy-anthraquinone (Dan)		$C_{14}H_6O_2(OH)_2$ – Mw: 240 Mp: 190-195 °C Shade: reddish to orange	(27)
Dermolutein 8-methoxy-3-methyl-1,6-dihydroxy-anthraquinone-2-carboxylic acid		$C_{17}H_{10}O_5(OH)_2$ – Mw: 326 Shade: yellow	(32)
Fallacinal 3-formyl-6-methoxy-1,8-dihydroxy-anthraquinone		$C_{16}H_8O_4(OH)_2$ – Mw: 298 Mp: 227–228 °C Shade: yellow	(33)
Frangulin A (=franguloside) 3-O-primeverose-6-methyl-1,8-dihydroxy-anthraquinone (Fran)		$C_{21}H_{18}O_7(OH)_2$ – Mw: 416 Shade: orange	(24)

Lucidin (=henine) 2-hydroxymethyl-1,3-dihydroxy-anthraquinone		$C_{15}H_8O_3(OH)_2$ – Mw: 270 Mp: 300 °C Shade: red	(19, 23, 34)
Munjistin 1,3-dihydroxy-anthraquinone-2-carboxylic acid (Mun)		$C_{15}H_6O_4(OH)_2$ – Mw: 284 Shade: orange-red	(17)
Nordamnacanthal 2-formyl-1,3-dihydroxy-anthraquinone (Nor)		$C_{15}H_6O_3(OH)_2$ – Mw: 268 Mp: 214–218 °C Shade: orange-yellow	(16, 17)
Phomarin 3-methyl-1,6-dihydroxy-anthraquinone		$C_{15}H_8O_2(OH)_2$ – Mw: 254 Shade: yellow UV (MeOH) λ_{max} 410, 293, 268, 219 nm; IR (KBr) ν_{max} 3414, 3310, 2925, 2856, 1677, 1640, 1603, 1579, 1488, 1389, 1305, 1268, 1191, 1153, 1020, 945, 894, 854, 817, 795, 795 cm^{-1} ; ¹H NMR (acetone, 500 MHz) δ 8.14, 7.65, 7.56, 7.28, 7.10.	(18)
Physcion (=parietin) 6-methoxy-3-methyl-1,8-dihydroxy-anthraquinone (Phy)		$C_{16}H_{10}O_3(OH)_2$ – Mw: 284 Mp: 207 °C; shade: yellow UV (EtOH) λ_{max} 435, 283, 265, 253, 226 nm; IR (CHCl ₃) ν_{max} 2929, 2854, 1626, 1568, 1483, 1140 cm^{-1} ; ¹H NMR (CDCl ₃ , 500 MHz) δ 7.08, 7.62, 7.36, 6.69, 3.94, 2.45.	(35)
Questin 8-methoxy-3-methyl-1,6-dihydroxy-anthraquinone		$C_{16}H_{10}O_3(OH)_2$ – Mw: 284 Mp: 301–303 °C Shade: yellow to orange-brown Soluble in aqueous sodium carbonate	(36)
Quinizarin 1,4-dihydroxy-anthraquinone (Qza)		$C_{14}H_6O_2(OH)_2$ – Mw: 240 Mp: 198–199 °C Shade: Orange to red-brown Slightly soluble (but soluble in hot water) UV (MeOH) λ_{max} 480 nm	(19, 34, 37, 38)
Rhein (=cassic acid) 1,8-dihydroxy-anthraquinone-3-carboxylic acid (Rhe)		$C_{15}H_6O_4(OH)_2$ – Mw: 284 Mp: 320–321 °C; shade: orange Insoluble in water IR (KBr) ν_{max} 3441, 3407, 1693, 1673, 1624, 1560, 1449 cm^{-1} ; ¹H NMR (CDCl ₃ , 500 MHz) δ 11.87, 8.07, 7.81, 7.71, 7.69, 7.38.	(33)
Rubiadin 2-methyl-1,3-dihydroxy-anthraquinone		$C_{15}H_8O_2(OH)_2$ – Mw: 254 Mp: 290 °C (in alcohols) and 302 °C (in glacial acetic acid) Shade: yellow UV (EtOH) λ_{max} 415, 280, 246 nm.	(19, 39, 40)

Soranjidiol 2-methyl-1,6-dihydroxy-anthraquinone		$C_{15}H_8O_2(OH)_2$ – Mw: 254	(39, 40)
Teloschistin (= fallacinol or phallacinol) 3-hydroxymethyl-1,8-dihydroxy-6-methoxy-anthraquinone		$C_{16}H_{10}O_4(OH)_2$ – Mw: 300 Mp: 236–237 °C; Shade: yellow	(33)
Xanthopurpurin 1,3-dihydroxy-anthraquinone (Xpu)		$C_{14}H_6O_2(OH)_2$ – Mw: 240 Shade: red	(19, 23, 37)
- Trihydroxyanthraquinones (OH)₃:			
Anthragallol (= alizarin brown) 1,2,3-trihydroxy-anthraquinone (Agl)		$C_{14}H_5O_2(OH)_3$ – Mw: 256 Mp: 312–313 °C Shade: orange	(19)
Citreosein 6-hydroxymethyl-1,3,8-trihydroxy-anthraquinone		$C_{15}H_7O_3(OH)_3$ – Mw: 286 Shade: yellow	(41, 42)
Dermoglaucin 3-methyl-1,7,8-trihydroxy-6-methoxy-anthraquinone		$C_{16}H_9O_3(OH)_3$ – Mw: 300 Shade: red	(32)
Dermorubin 3-methyl-1,4,6-trihydroxy-8-methoxy-anthraquinone-2-carboxylic acid		$C_{17}H_9O_5(OH)_3$ – Mw: 344 Shade: red	(32)
Emodin (= frangula emodin) 3-methyl-1,6,8-trihydroxy-anthraquinone (Emo)		$C_{15}H_7O_2(OH)_3$ – Mw: 270 Mp: 254–256 °C; Shade: orange	(33, 42, 43)
Endrocroc 3-methyl-1,6,8-trihydroxy-anthraquinone-2-carboxylic acid		$C_{16}H_7O_4(OH)_3$ – Mw: 314 Shade: yellow	(32)

Erythroglauin 3-methyl-1,4,8-trihydroxy- 6-methoxy-anthraquinone			C ₁₆ H ₉ O ₅ (OH) ₃ – Mw: 300 Shade: red	(4)
Flavokermesic acid (= <i>Laccaic acid D</i>) 1-methyl-3,6,8-trihydroxy- anthraquinone-2-carboxylic (Flk) acid			C ₁₆ H ₇ O ₆ (OH) ₃ – Mw: 314 Shade: yellow	(15, 44, 45)
Flavopurpurin (= <i>alizarin Y</i>) 1,2,6-trihydroxy-anthraquinone			C ₁₅ H ₅ O ₅ (OH) ₃ – Mw: 256 Shade: yellow	(19)
Helminthosporin 3-methyl-1,5,8-trihydroxy- anthraquinone			C ₁₅ H ₇ O ₅ (OH) ₃ – Mw: 270 Shade: brown	(4)
Islandicin 2-methyl-1,4,5-trihydroxy- anthraquinone			C ₁₅ H ₇ O ₅ (OH) ₃ – Mw: 270 Mp: 217–219 °C Shade: red	(46)
Morindone 6-methyl-1,2,5-trihydroxy- anthraquinone			C ₁₅ H ₇ O ₅ (OH) ₃ – Mw: 270 Shade: yellowish-red	(47)
Pseudopurpurin 1,3,4-trihydroxy-anthraquinone- 2-carboxylic acid (Psp)			C ₁₅ H ₅ O ₆ (OH) ₃ – Mw: 300 Shade: red	(15)
Purpurin 1,2,4-trihydroxy-anthraquinone (Pur)			C ₁₄ H ₅ O ₅ (OH) ₃ – Mw: 256 Mp: 265–270 °C; shade : yellow to red (acid); red to violet (alkaline)	(15, 23, 37, 38)
Rubrocristin 2-methyl-1,4,7-trihydroxy- 5-methoxy-anthraquinone			C ₁₆ H ₉ O ₅ (OH) ₃ – Mw: 300 Shade: red	(36)

Skyrin		$C_{30}H_{15}O_7(OH)_3$ – Mw: 358 Shade : yellow to red (48)
Tetrahydroxyanthraquinones (OH)₄:		
Averythrin 2-(1-Hexenyl)-1,3,6,8-tetrahydroxy-anthraquinone (Avt)		$C_{20}H_{14}O_2(OH)_4$ – Mw: 354 Shade : orange (49)
Carminic acid 2- α -D-glucopyranosyl-8-methyl-1,3,4,6-tetrahydroxy-anthraquinone-7-carboxylic acid (Car)		$C_{22}H_{16}O_9(OH)_4$ – Mw: 492 Mp: 120 °C; shade: red to violet (alkaline); or orange (acid) (15, 22, 37, 50) Good solubility in water UV (EtOH) λ_{max} 495, 491, 311, 278 nm.
Catenarin 3-methyl-1,4,5,7-tetrahydroxy-anthraquinone		$C_{15}H_6O_2(OH)_4$ – Mw: 286 Mp: 240 °C; shade: red (51) UV (MeOH) λ_{max} 525, 512, 490, 480, 463, 306, 278, 257, 230 nm; ¹ H NMR (CDCl ₃ , 500 MHz) δ 13.34, 12.42, 12.35, 7.32, 7.13, 6.66, 2.35.
Cynodontin 2-methyl-1,4,5,8-tetrahydroxy-anthraquinone (Cyn)		$C_{15}H_6O_2(OH)_4$ – Mw: 286 Shade: bronze (4)
Dermocybin 3-methyl-1,5,7,8-tetrahydroxy-6-methoxy-anthraquinone		$C_{16}H_8O_3(OH)_4$ – Mw: 316 Mp: 228–229 °C Shade: red (52) UV (EtOH) λ_{max} 521, 486, 459, 279, 262, 219 nm.
Kermesic acid: 8-methyl-1,3,4,6-tetrahydroxy-anthraquinone-7-carboxylic acid (Ker)		$C_{16}H_6O_4(OH)_4$ – Mw: 330 Mp > 320 °C Shade : Dark red in acidic pH ; Violet in aqueous NaOH (37, 44, 45) UV (MeOH) λ_{max} 545, 496 nm.
Laccaic acid A (LaA)		$C_{26}H_{15}NO_8(OH)_4$ – Mw: 537 Shade: red (37, 44, 45) UV (H ₂ SO ₄) λ_{max} 558, 518, 361, 302 nm.

Laccaic acid B (LaB)		$C_{24}H_{12}O_8(OH)_4$ – Mw: 496 Shade: red	(37, 44, 45)
Laccaic acid C (LaC)		$C_{25}H_{13}NO_9(OH)_4$ – Mw: 539 Shade: dark red	(37, 44, 45)
Laccaic acid E (LaE)		$C_{24}H_{13}NO_7(OH)_4$ – Mw: 495 Shade: red UV (H_2SO_4) λ_{max} 496 nm.	(37, 44, 45)
Tritisporin 3-hydroxymethyl-1,4,6,8-tetrahydroxy-anthraquinone		$C_{15}H_7O_3(OH)_3$ – Mw: 286 Shade: brownish-red	(4)

*Mw: molecular weight; Mp: Melting point; Pr: primeverose; Rh: rhamnose

(yellow(acid) to red(alkali)), pseudopurpurin (orange), purpurin (dark red) and lucidin (red). A practical HAQN classification, according to the respective biosynthetic pathway of the compound and the position of the functional groups added on the 9,10-anthraquinone skeleton, is shown on Table 3. This classification is partially based on that proposed by Rafaëly et al. in 2008⁴⁵. It appears that the natural HAQN pigments formed *via* the chorismate/*o*-succinylbenzoic acid pathway are all classified in the entire ‘group E’ of HAQN dyes because they have substitution only on one aromatic ring, like alizarin and purpurin. In contrast, the HAQN pigments that are synthesized *via* the polyketide pathway are classified in the ‘group A₁’ (compounds show substitution on both aromatic rings and at least two hydroxyl groups in both R1 and R8 positions, like emodin, chrysophanol or physcion) or into the ‘group A₂’ of HAQN dyes (compounds show substitution on both aromatic rings and at least two hydroxyl groups in R1 and R6 positions and one methoxyl group in R8 position, like dermolutein and dermorubin).

3 The main natural sources of hydroxyanthraquinoid pigments

The natural HAQN pigments are mainly found in plants like *Rubiaceae*, *Polygonaceae*, *Rhamnaceae*, *Fabaceae*, *Liliaceae*, *Bignoniaceae* and *Pedaliaceae*, in lichens and in the animal kingdom (insects).

3.1 Hydroxyanthraquinoid pigments from plants

In plants, the dyestuff is often extracted from dried roots. HAQN pigments are mostly present as sugar derivatives—the glycosides—but the free form—the aglycones—are widely distributed as well. For example, the European madder roots contain 2%–3.5% of the dry weight of di- and tri-

hydroxyanthraquinone-glycosides and, in general for higher plants, the HAQN-based colorant content from the dry mass is often under 5%¹⁵. The anthraquinone glycosides are formed when one or more sugar molecules, mostly glucose or rhamnose, are bound to the aglycone by a β -glycoside linkage to hydroxyl group at position C-8 (in the case of glucose) or the one at C-6 (in the case of rhamnose)⁵⁸. During storage, hydrolysis of the glycosides occurs, which is completed under acidic conditions. In the literature, a total of more than 35 anthraquinoid compounds have been reported to be extracted from roots of European madder (*Rubia tinctorum* Linn., *i.e.* the most important species of the plant family *Rubiaceae*), even if a part of the compounds is believed to be artefacts formed during extraction or drying¹⁵. The main HAQN colouring compounds of plants of the *Rubiaceae* family (e.g. *Rubia* spp., *Galium* spp., *Morinda* spp., *Hypericum* spp., *Polygonum* spp. and *Cinchona* spp.)^{14,15,23,34,38–40,45,59–73} are alizarin (yellow to red, group E₃), pseudopurpurin (orange, group E₂), purpurin (dark red, group E₂), lucidin-3-*O*-primeveroside (red, group E₃), ruberythric acid (golden-yellow, group E₃), nordamnacanthal (orange, group E₃) and munjistin (orange-red, group E₃) (see Table 4). These colouring compounds are all classified in the entire ‘group E’ of HAQN dyes and they are formed through the chorismate/*o*-succinylbenzoic acid pathway as mentioned above.

In contrast, in other higher plants such as the *Polygonaceae* (*Rheum* spp., *Rumex* spp.)^{24,29,31,43,74–80}, *Rhamnaceae* (*Rhamnus* spp.)^{25,81}, *Fabaceae* (*Cassia* spp.)^{30,33,82}, *Liliaceae* (*Aloe* spp.)^{83,84} and *Pedaliaceae* (*Ceratotheca* spp.)⁸⁵ families, the most common naturally occurring HAQN pigments are synthesized *via* the polyketide pathway (Fig. 2). Relevant pigments are emodin (yellow), aloe-emodin (yellow), physcion (yellow), rhein (orange) and chrysophanol (orange-red) (see

Table 4). These HAQN pigments are all classified in the 'group A₁' of HAQN because they show substitution on both

aromatic rings and at least two hydroxyl groups in both R1 and R8 positions.




Figure 2. The two main biosynthetic pathways of hydroxyanthraquinoid (HAQN) pigments in organisms

Table 3. Position of functional groups added on 9,10-anthraquinone skeleton in natural hydroxyanthraquinoid (HAQN) pigments and their classification into several groups

HAQN compound	R1	R2	R3	R4	R5	R6	R7	R8	shade
Group A1: at least two hydroxyl groups in both R1 and R8 positions:									
Aloe-emodin	OH	H	CH ₂ OH	H	H	H	H	OH	orange
Averythrin	OH	C ₆ H ₁₁ H	OH	H	H	OH	H	OH	orange
Chrysophanol	OH	H	CH ₃	H	H	H	H	OH	orange-red
Citreorosein	OH	H	OH	H	H	CH ₂ OH	H	OH	yellow
Cynodontin	OH	H	CH ₃	OH	OH	H	H	OH	bronze
Danthron (= chrysin)	OH	H	H	H	H	H	H	OH	reddish to orange
Dermoglucanin	OH	H	CH ₃	H	H	OCH ₃	OH	OH	red
Dermocycin	OH	H	CH ₃	H	OH	OCH ₃	OH	OH	red
Emodin	OH	H	CH ₃	H	H	OH	H	OH	orange
Endocrocin	OH	COOH	CH ₃	H	H	OH	H	OH	yellow
Erythroglucanin	OH	H	CH ₃	OH	H	OCH ₃	H	OH	red
Fallacinal	OH	H	CHO	H	H	OCH ₃	H	OH	yellow
Frangulin A	OH	H	O-Rh	H	H	CH ₃	H	OH	orange
Helminthosporin	OH	H	CH ₃	H	OH	H	H	OH	maroon
Physcion (= parietin)	OH	H	CH ₃	H	H	OCH ₃	H	OH	yellow
Rhein	OH	H	COOH	H	H	H	H	OH	orange
Teloschistin (= fallacinal)	OH	H	CH ₂ OH	H	H	OCH ₃	H	OH	yellow
Tritisporin	OH	H	CH ₂ OH	OH	H	OH	H	OH	reddish brown
Group A2: at least two hydroxyl groups in R1 and R6 positions, and one methoxyl group in R8 position									
Questin	OH	H	CH ₃	H	H	OH	H	OCH ₃	yellow to orange
Dermolutein	OH	COOH	CH ₃	H	H	OH	H	OCH ₃	yellow
Dermorubin	OH	COOH	CH ₃	OH	H	OH	H	OCH ₃	red
Group B: Four hydroxyl groups in R1, R3, R4 and R6 positions, and one carboxyl group in R7 position:									
Kermesic acid	OH	H	OH	OH	H	OH	COOH	CH ₃	red
Carminic acid	OH	Glc	OH	OH	H	OH	COOH	CH ₃	red
Laccaic acid A	OH	C ₁₀ H ₁₂ O ₂ N	OH	OH	H	OH	COOH	COOH	red
Laccaic acid B	OH	C ₈ H ₉ O ₂	OH	OH	H	OH	COOH	COOH	red
Laccaic acid C	OH	C ₉ H ₁₀ O ₃ N	OH	OH	H	OH	COOH	COOH	red
Laccaic acid E	OH	C ₈ H ₁₀ ON	OH	OH	H	OH	COOH	COOH	red
Group C1: at least two hydroxyl groups in R1 and R4 positions, and at least one functional group (-OH, -CH₃) in R7 position:									
Austrocortinin	OH	OCH ₃	H	OH	H	H	CH ₃	H	red
Catenarin	OH	H	CH ₃	OH	OH	H	OH	H	red
Rubrocristin	OH	CH ₃	H	OH	OCH ₃	H	OH	H	red
Group C2: at least two hydroxyl groups in R1 and R6 positions, and position R8 unsubstituted:									
Flavopurpurin	OH	OH	H	H	H	OH	H	H	yellow
Phomarin	OH	H	CH ₃	H	H	OH	H	H	yellow
Soranjidiol	OH	CH ₃	H	H	H	OH	H	H	yellowish-red
Group C3: at least two hydroxyl groups in R1 and R5 positions, and position R8 unsubstituted:									
Islandicin	OH	CH ₃	H	OH	OH	H	H	H	red
Morindone	OH	OH	H	H	OH	CH ₃	H	H	yellowish-red
Group D: no hydroxyl group in R1 position, at least one hydroxyl group in R2 position, and position R8 unsubstituted:									
Anthraflavic acid (= anthraflavin)	H	OH	H	H	H	OH	H	H	yellow
Skyrin	C ₁₅ H ₉ O ₅	OH	H	OH	OH	H	CH ₃	H	yellow to red
Compounds with functional groups only on one aromatic ring (group E):									
Group E1: no hydrogen intramolecular bonds with carbonyl function:									
3-MeO-hystazarin	H	OH	OCH ₃	H	H	H	H	H	
Damnacanthal	OCH ₃	CHO	OH	H	H	H	H	H	pale yellow
Group E2: at least 2 hydroxyl groups in R1 and R4 positions:									
Purpurin	OH	OH	H	OH	H	H	H	H	dark red
Pseudopurpurin	OH	COOH	OH	OH	H	H	H	H	orange
Quinizarin	OH	H	H	OH	H	H	H	H	orange-red
Group E3: 2 or 3 hydroxyl groups but always one in position R1 and none in position R4:									
Alizarin	OH	OH	H	H	H	H	H	H	yellow to red
Anthragallol	OH	OH	OH	H	H	H	H	H	orange
Lucidin-3-O-primeveroside	OH	CH ₂ OH	O-Pr	H	H	H	H	H	red
Lucidin (= henine)	OH	CH ₂ OH	OH	H	H	H	H	H	red
Munjistin	OH	COOH	OH	H	H	H	H	H	orange-red
Nordamnacanthal	OH	CHO	OH	H	H	H	H	H	orange-yellow
Pachybasin	OH	CH ₃	H	H	H	H	H	H	yellow
Ruberythric acid	OH	O-Pr	H	H	H	H	H	H	golden-yellow
Rubiadin	OH	CH ₃	OH	H	H	H	H	H	yellow
Xanthopurpurin	OH	H	OH	H	H	H	H	H	red

Pr: primeverose; Glc: glucose; Rh: rhamnose

Table 4. Natural occurrence of hydroxyanthraquinoid (HAQN) pigments in plants

Latin name	Main colouring components	References
Rubiaceae		
<i>Rubia tinctorum</i> L. (= European madder)	alizarin; purpurin; pseudopurpurin; lucidin; rubiadin; xanthopurpurin; munjinstin; anthraflavin; quinizarin; danthron; anthragalloyl; nordamnacanthal; ruberythric acid; lucidin primeveroside, alizarin-2-methyl ether; lucidin- ω -ethyl-ether; munjinstin ethyl ether	(23, 34, 38, 59)
<i>Rubia cordifolia</i> L.	rubiadin; alizarin; purpurin; pseudopurpurin; lucidin; munjinstin; xanthopurpurin; tectoquinone	(14, 15, 23, 60)
<i>Rubia akane</i>	purpurin; ruberythric acid	(15, 45)
<i>Rubia peregrina</i> L.	pseudopurpurin	(15)
<i>Galium aparine</i> L.	nordamnacanthal; xanthopurpurin; rubiadin	(61)
<i>Galium sinaicum</i>	7-methyl-anthragalloyl-1,3-dimethyl ether; 7-methyl-anthragalloyl-2-methyl ether; 6-methyl-anthragalloyl-3-methyl ether; 8-hydroxy-anthragalloyl-2,3-dimethyl ether; 7-formyl-anthragalloyl-1,3-dimethylether; 6-hydroxy-xanthopurpurin; 6-methoxy-lucidin- ω -ethyl ether; copareolatin; copareolatin-6,7-dimethyl ether; copareolatin-5,7-dimethyl ether	(62)
<i>Galium verum</i> L. (= Lady's bedstraw)	alizarin; 1,3-dihydroxy-2-methoxymethyl; 1,3-dimethoxy-2-hydroxy, 1,3-dihydroxy-2-acetoxy; 1-hydroxy-2-hydroxymethyl; 1,3-dihydroxy-2-methyl; 1-methoxy-2-hydroxyanthraquinones; 1,3-dihydroxy-2-hydroxymethyl-6-methoxy anthraquinones	(15, 63)
<i>Galium mollugo</i> L.	pseudopurpurin	(15)
<i>Galium spurium</i>	8-hydroxy-3-methoxy-7-methyl-1,2-methylenedioxy-anthraquinone; 2,8-dihydroxy-1,3-dimethoxy-7-methyl-anthraquinone	(64)
<i>Morinda officinalis</i>	alizarin; purpurin; pseudopurpurin; lucidin; rubiadin; 2-hydroxy-1-methoxy-anthraquinone; 1,3,8-trihydroxy-2-methyl-anthraquinone	(40)
<i>Morinda elliptica</i>	alizarin; purpurin; pseudopurpurin; lucidin; rubiadin; moridone; soranjidol; nordamnacanthal; alizarin-1-methylether; lucidin- ω -methylether	(39)
<i>Morinda citrifolia</i>	damnacanthal; morindone; morindin; alizarin; physcion; morenone; morenone; ruberythric acid; rubiadin; lucidin	(65)
<i>Cinchona ledgeriana</i>	purpurin; rubiadin; anthragalloyl-1,2-dimethylether; anthragalloyl-1,3-dimethylether; 1-hydroxy-2-hydroxymethylanthraquinone; 1-hydroxy-2-methylanthraquinone; morindone-5-methylether (or 1,7-dihydroxy-8-methoxy-2-methylanthraquinone)	(66, 67)
<i>Cinchona pubescens</i>	purpurin; alizarin-2-methylether; anthragalloyl-1,2-dimethylether; purpurin-1-methylether; 1-hydroxy-2-hydroxymethyl-anthraquinone; 2-hydroxy-1,3,4-trimethoxy-anthraquinone	(68)
<i>Cinchona succirubra</i>	emodin; anthrapurpurin; quinizarin; 2,6-dihydroxyanthraquinone; 1,8-dihydroxyanthraquinone	(69)
<i>Cinchona robusta</i>	robustaquinones (A–H); 1,3,8-trihydroxy-2-methyl anthraquinone; copareolatin 6-methylether	(70)
<i>Asperula tinctoria</i> L.	alizarin; rubiadin	(15, 45)
<i>Asperula arvensis</i> L.	alizarin	(45)
<i>Oldenlandia umbellata</i> L.	alizarin; 1,2,3-trimethoxyanthraquinone; 3-MeO-hystazarin, ruberythric acid; 1,3-dimethoxy-2-hydroxyanthraquinone; 1,2-dimethoxyanthraquinone; 1-methoxy-2-hydroxyanthraquinone; 1,2-dihydroxyanthraquinone	(45, 71, 72)
<i>Hedyotis auricularia</i> L.	alizarin	(45)
<i>Crucianella maritima</i> L.	alizarin; 3-formyl-1-hydroxy-2-methoxy anthraquinone; alizarin-1-methyl ether; 1,4-dihydroxy-2-methoxy-anthraquinone	(45, 73)
<i>Coprosma lucida</i>	anthragalloyl; lucidin; rubiadin	(45)
<i>Hymenodictyon excelsum</i>	anthragalloyl	(45)
Polygonaceae		
<i>Rheum officinale</i>	emodin; chrysophanol; rhein	(43, 74)
<i>Rheum palmatum</i>	chrysophanol; aloë-emodin; rhein; physcion; citreosein	(24, 75)
<i>Rheum emodi</i>	emodin; chrysophanol; aloë-emodin; rhein; physcion	(29, 76)
<i>Rheum rhabarbarum</i>	emodin; chrysophanol; aloë-emodin; rhein; physcion	(31)
<i>Rumex dentatus</i>	chrysophanol; physcion	(24)
<i>Rumex crispus</i>	chrysophanol; parietin	(77, 78)
<i>Rumex acetosa</i>	chrysophanol; physcion; emodin; emodin-8-O- β -D-glucopyranoside	(79)
<i>Rumex obtusifolius</i>	aloë-emodin; chrysophanol; emodin	(80)
<i>Rumex spp. (19 spp.)</i>	emodin; chrysophanol; physcion; aloë-emodin; rhein	(80)
Rhamnaceae		
<i>Rhamnus saxatilis</i>	emodin; chrysophanol; aloë-emodin; rhein; physcion	(25)
<i>Rhamnus alpinus</i> L.	aloë-emodin; rhein; emodin; chrysophanol; physcion	(81)
Fabaceae		
<i>Cassia occidentalis</i> L.	emodin, chrysophanol, aloë-emodin, rhein, physcion	(30)
<i>Cassia tora</i>	emodin; rhein; physcion	(33)
<i>Senna alata</i>	aloë-emodin; emodin; rhein; chrysophanol	(82)
Liliaceae		
<i>Aloes spp. (32 spp.)</i>	chrysophanol; asphodelin; chrysophanol-8-methyl ether; aloëchryson; helminthosporin; aloësaponols; aloësaponarins	(83, 84)
Bignoniaceae		
<i>Tecoma ipes</i>	tectoquinone	(45)
Pedaliaceae		
<i>Ceratotheca triloba</i> (Bernh.)	1-hydroxy-4-methylanthraquinone	(85)

3.2 Hydroxyanthraquinoid pigments from lichens

HAQN pigments found in some lichens are synthesized *via* the polyketide pathway. For example, the main colouring compounds in the lichens of the family *Teloschistaceae* (*Caloplaca* sp., *Xanthoria* sp. or *Teloschistes* sp.)^{15,48,86–90} and the family *Trypetheliaceae* (*Laurera benguelensis*)^{87,88} are emodin, physcion, teloschistin (yellow, group A₁) and fallacinal (yellow, group A₁) (see Table 5). The lichens *Nephroma laevigatum* and *Heteroderma obscurata* also contain emodin⁹¹ whereas skyrin (*i.e.* a yellow to red pigment classified in the ‘group D’ of HAQN which show substitution on both aromatic rings; see Table 3) is the main component of *Cladonia* species⁴⁸.

3.3 Hydroxyanthraquinoid pigments from insects

In animals, HAQN-type pigments are known to be present only in a few insect species (see Table 5). Concerning the red carminic acid, kermesic acid and laccaic acid obtained from cochineal (*Dactylopius coccus*)^{15,22,37,50} kermes (*Kermes vermilio*)^{15,44} and lac (*Kerria lacca*)^{15,44,45}, respectively, they contain functional groups on both aromatic rings and particularly four hydroxyl groups in R1, R3, R4 and R6 positions, and one carboxyl group in R7 position. So these animal anthraquinoid glycosides are all classified in the ‘group B’ of HAQN (see Table 3). In both cochineal and kermes the pigments were obtained from the body and eggs of the female insect. Although the various species of the genus *Porphyrophora*, e.g. Armenian cochineal (*P. hamelli*) and Polish cochineal (*P. polonica*), also contain carminic acid, dried specimens of *Dactylopius coccus* have a much higher content (15%–20%) of carminic acid, compared with only 0.8% and 0.6% for the Armenian and Polish ones, respectively^{15,44,45}. Lac insects of the *Kerria* family (e.g.

Kerria lacca and *K. chinensis*) contain mainly laccaic acids like laccaic acid A.

3.4 Hydroxyanthraquinoid pigments from fungi

HAQN pigments are widespread in nature and have been also found abundantly in microorganisms, particularly in filamentous fungi belonging to *Penicillium* spp. and *Aspergillus* spp., with different shades (see Table 6). For example, the pigment emodin was isolated from strains of *Penicillium citrinum* and *P. islandicum*^{5,92}. The natural food colorant Arpink red™ manufactured by the Ascolor Biotech Czech company was claimed to be produced by fermentation and bioprocess engineering using the strain *Penicillium oxalicum* var. *Armeniaca* CCM 8242 obtained from soil^{5,92}. On the second day of cultivation of this fungus in liquid broth containing carbohydrates, zinc sulfate and magnesium sulfate, a red colorant is released in the medium, increasing up to 1.5–2.0 g/L of broth after 3–4 days². After biosynthesis of the red colorant, the liquid is separated from the biomass by centrifugation or filtration. The liquid is then acidified to pH 3.0–2.5 to precipitate the colorant. The precipitate is dissolved in ethyl alcohol and filtered. Following removal of alcohol, the colorant is obtained in the crystalline form as a dark red powder. In strains of *Penicillium purpurogenum*⁹³, a red pigment of HAQN-type (none completely characterized) was also observed.

Some strains of *Aspergillus* (*A. glaucus*, *A. cristatus* and *A. repens*)^{4,5,35,36,92–94} were found to produce known yellow and red HAQN compounds such as emodin (yellow, group A₁), physcion (yellow, group A₁), questin (yellow to orange-brown, group A₂), erythroglaucon (red, group A₁), catenarin (red, group C₁; see Table 3) and rubrocristin (red, group C₁; see Table 3). However, by using *Penicillium* or *Aspergillus* strains,

Table 5. Natural occurrence of hydroxyanthraquinoid (HAQN) pigments in lichens and in insects

Dye source	Latin name	Main colouring components	References
Lichens:			
Teloschistaceae	<i>Xanthoria</i> spp.	physcion; emodin; parietin; fallacinal; teloschistin; citreoresein; erythroglaucon; fallacinal	(15, 48, 86, 87)
	<i>Xanthoria parietina</i> L.	physcion; fallacinal; fallacinal; emodin; parietic acid	(88)
	<i>Xanthoria mandshurica</i>	erythroglaucon; parietin	(89)
	<i>Xanthoria fallax</i>	fallacinal; emodin; fallacinal; erythroglaucon; parietin	(89)
	<i>Caloplaca</i> spp.	2-chloroemodin; citreoresein; emodin; fallacinal; parietin; physcion; teloschistin	(48, 87)
	<i>Caloplaca cerina</i>	emodin; fallacinal; physcion; teloschistin	(33)
	<i>Caloplaca erythrantha</i>	emodin; 7-chloroemodin	(86)
	<i>Teloschistes exilis</i>	parietin; teloschistin	(86)
	<i>Teloschistes</i> spp. (29 spp.)	parietin; emodin; teloschistin; fallacinal; parietic acid; erythroglaucon	(90)
	Trypetheliaceae	<i>Laurera benguelensis</i>	parietin; physcion; citreoresein; emodin; fallacinal; teloschistin
Nephromataceae	<i>Nephroma laevigatum</i>	emodin; 7-chloroemodin; 7-chloro-1-O-methylemodin; 5-chloro- ω -hydroxyemodin; 7-chloro-1-O-methyl- ω -hydroxyemodin; 5-chloroemodin; 5-chloro-1-O-methylemodin; 5-chloro-1-O-methyl- ω -hydroxyemodin;	(91)
Physciaceae	<i>Heteroderma obscurata</i>	emodin	(91)
Cladoniaceae	<i>Cladonia</i> spp.	skyrin	(48)
Insects:			
Dactylopius	<i>Dactylopius coccus</i> Costa (cochineal)	carminic acid [food additive E120(ii)]	(15, 22, 37, 50)
Porphyrophora	<i>Porphyrophora hameli</i> B. & <i>Porphyrophora polonica</i> L.	carminic acid; flavokermesic acid (LaE); kermesic acid	(15, 44, 45)
Kermes	<i>Kermes vermilio</i> Planchon (= <i>Kermococcus vermilio</i>)	kermesic acid	(15, 44)
Kerria	<i>Kerria lacca</i> Kerr	laccaic acids (A, B, C, E)	(15, 44, 45)

several known mycotoxins were coproduced in the medium, e.g. secalonin acid D, oxaline, citrinin, tanzawaic acid A, cyclochlorotene, islanditoxin, luteoskyrin, erythrokyrin, rugulosin or aspergillide A (Table 6). Many of these mycotoxins are pigmented, that is, naphthoquinones from *Aspergillus* and *Penicillium*. All these fungal secondary metabolites (on one hand, the yellow and red HAQN pigments that show substitution on both aromatic rings and, on the other hand, the naphthoquinone-type mycotoxins) arise biosynthetically by the same polyketide pathway. The cytotoxic activity of naphthoquinones, and of mycotoxins in general, against mouse leukemia and HeLa cells has been mainly reported in the literature. Moreover, along with the antibiotic and toxic activities, naphthoquinones revealed mutagenic and carcinogenic properties. The results suggested that these fungal strains could not be used to provide safe fungal hydroxyanthraquinoid pigments as potent natural food grade colorants.

Species of *Eurotium* spp. (*E. amstelodami*, *E. chevalieri* and *E. herbariorum*)^{5,92} were found to produce the yellow pigment physcion and the red pigment erythroglaucon (group A₁), however they produce in addition the mycotoxin echinulin and two benzaldehyde colouring compounds: flavoglaucon (yellow) and auroglaucon (red) (see Table 6). In the same way, it has been demonstrated that a coproduction of red hydroxyanthraquinoid pigments (with no hydroxyl substituents at the positions R1 and R4) and mycotoxins such as fusaric acid, nectriafurone, monoliformin and gibepyrone, occurs by using strains of *Fusarium oxysporum* isolated from roots of diseased citrus trees^{95–97} (Table 6).

Apart from those mycotoxigenic fungi, there are other filamentous fungi that have the ability to produce known HAQN pigments that arise biosynthetically by the polyketide pathway more particularly, without coproduction of mycotoxins. A strain of *Dermocybe sanguinea* (= *Cortinarius sanguineus*) has been identified as producing the red HAQN glycoside dermocybin-1-β-D-glycopyranoside giving the typical red colour of the fruiting bodies and the spores, in addition with both emodin and physcion pigments^{3,15,18,52}. In the fresh fungi as much as 90% of the pigments exist as glycosides. The detection of emodin-glycosides and physcion-glycosides was also pointed from a strain of *Dermocytes* spp.⁹⁸

Strains of *Trichoderma aureoviride*⁹⁹ and *T. harzianum*^{29,100} were found to produce yellow pigment pachybasin (group E₃) and also the orange-red pigment chrysophanol (group A₁). Both species of *Trichoderma polysporum* and *T. viride*^{99,101,102} can also produce pachybasin in addition to the emodin and chrysophanol pigments. Several HAQN-type pigments have been isolated from cultures of *Curvularia lunata*^{3–5}. The main pigments characterized were erythroglaucon (red, group A₁), catenarin (red, group C₁), chrysophanol (orange-red, group A₁), helminthosporin (maroon, group A₁) and cynodontin (bronze, group A₁). Cynodontin extracted from the biomass of *C. lunata* has been converted successfully to two anthraquinone biodyes (Disperse blue 7 and Acid Green 28). The properties of these biodyes applied to knitted polyamides were compared with those of conventional dyes and found to be identical to all-important aspects³. Several species of *Drechslera* (e.g. *D. teres*, *D. graminea*, *D. tritici-repentis*, *D. phlei*, *D. dictyoides* and *D. avenae*) give HAQN pigments like catenarin (red, group C₁), helminthosporin (maroon, group A₁), cynodontin

(bronze, group A₁), tritisporin (reddish brown, group A₁) and erythroglaucon (red, group A₁), without coproduction of mycotoxins^{3,103}. Other HAQN pigments like averythrin (orange, group A₁) and averythrin-6-monomethyl ether were isolated and identified from a culture of *Herpotrichia rhodosticta* without coproduction of known mycotoxins¹⁰⁴. More recently, a red pigment produced by a strain of *Isaria farinosa* was recently elucidated as a chromophore of the anthraquinone type¹⁰⁵. Similarly, the red pigment produced by *Paecilomyces sinclairii*, which was beforehand discovered but uncharacterized¹⁰⁶, is certainly of an identical chemical nature, i.e. an amino group linked to an anthraquinone structure¹⁰⁵.

4 Toxicity and carcinogenicity of some natural hydroxyanthraquinoid pigments

Anthraquinoid derivatives, including natural HAQN pigments, possess a broad spectrum of biological activities, including anti-inflammatory, anti-cancer, anti-viral, anti-fungal, anti-bacterial, astringent and purgative. In general, natural HAQN pigments and their intermediates have not been reported as strongly toxic substances, even if it is known that some anthraquinoid dyes are toxic or mutagenic^{107,108}. Due to its use as a food colorant in Japan, the safety of European madder extracts has been studied in the literature. For example, in an extensive study the European madder roots were extracted using different solvents and extracts were fractionated by chromatography. Several colour components extracted from madder roots were positive to mutagenicity tests as the yellow rubiadin pigment (group E₃; see Table 3) and the red lucidin pigment (group E₃) aglycones, which are metabolites of lucidin-3-*O*-primeveroside. From structure mutagenicity studies it was concluded that 1,3-dihydroxyanthraquinones that bear a methyl (-CH₃) or hydroxymethyl (-CH₂OH) group in position R2, e.g. rubiadin or lucidin, respectively, are mutagenic. For direct mutagenicity an oxygenated state of the benzylic carbon-2 is required. Mutagenic studies about lucidin more particularly showed that a reactive compound is formed from the metabolism of the pigment, which then reacts with DNA and possibly other macromolecules to form covalent adducts^{109,110}. Other 1,3-dihydroxyanthraquinones that do not possess a methyl or hydroxymethyl group in position R2, such as the orange pigment nordamnacanthal (group E₃) and the orange-red munjistin pigment (group E₃), are not found to be mutagenic, since the dehydration to the exomethylene compound is not possible under physiological conditions¹⁵. In a 13-week repeated oral dose toxicity study of madder colour, which was performed using F344 rats, the animals were fed a diet containing 0, 0.6, 1.2, 2.5 or 5.0% of colouring compounds extracted from madder roots. The results suggested that madder colour exerts mild toxicity, targeting liver, kidneys and possibly red blood cells and white blood cells, some renal changes being evident from 0.6% madder colour in diet. This is considered to be the lowest-observed adverse effect level (305.8–309.2 mg/kg of body weight per day)¹¹¹. Data are in agreement with another study performed in the same year in a medium-term multiorgan carcinogenesis bioassay in male F344 rats, which reported that madder colour demonstrated significant tumour-promoting effects in the liver and kidneys¹¹². More recently, an additional two-year carcinogenicity study conducted on male and female F344

Table 6. Natural occurrence of hydroxyanthraquinoid (HAQN) pigments in microorganisms

Genus	Species	HAQN colouring compounds (shade)		References	
		Non-toxic HAQN pigments	Potent toxic HAQN pigments (mycotoxins, color or colorless...)		
<i>Penicillium</i>	<i>P. oxalicum</i>	Arpink red TM (red)	-	(5, 92)	
	<i>P. citrinum</i>	-	emodin (yellow)	(5, 92)	
	<i>P. islandicum</i>	skyrin (yellow to red)	emodin (yellow)	(5, 92)	
<i>Aspergillus</i>	<i>P. purpurogenum</i>	red HAQN pigment (none completely characterized)	-	(93)	
	<i>A. glaucus</i>	erythroglauca (red), catenarin (red), cynodontin (bronze), helminthosporin (maroon), tritispurin (reddish brown)	emodin & physcion (yellow)	(4, 5, 92)	
	<i>A. cristatus</i>	catenarin (red), erythroglauca (red), rubrocrisin (red), questin (yellow)	emodin & physcion (yellow)	(36)	
	<i>A. repens</i>	erythroglauca (red)	physcion (yellow)	(35, 94)	
	<i>Eurotium spp.</i>	catenarin (red), erythroglauca (red), cynodontin (bronze), helminthosporin (maroon), tritispurin (reddish brown)	physcion (yellow)	(5, 92)	
	<i>Fusarium</i>	<i>Fusarium spp.</i>	catenarin (red), erythroglauca (red), cynodontin (bronze), helminthosporin (maroon), tritispurin (reddish brown)	chrysophanol (red)	(5)
		<i>F. oxysporum</i>	2-(1-hydroxyethyl)-3,8-dihydroxy-6-methoxy-anthraquinone 2-acetyl-3,8-dihydroxy-6-methoxy-anthraquinone (red)	-	(92, 95-97)
		<i>D. sp. WAT2963</i>	-	emodin & physcion-glycosides	(98)
		<i>Dermocybe D. sanguinea</i>	dermocybin-1-β-D-glycopyranoside (red), dermorubin (red), dermolutein (yellow), dermoglauca (red), 5-chlorodermorubin	emodin & physcion-glycosides	(3, 15, 18, 52)
	<i>Pachybasium</i>	<i>P. candidum</i>	pachybasin (yellow)	chrysophanol (red)	(18)
<i>Phoma P. exigua</i>		pachybasin (yellow), phomarin (yellow)	emodin (yellow), chrysophanol (red)	(18)	
<i>Trichoderma</i>	<i>T. aureoviride</i>	pachybasin (yellow)	chrysophanol (red)	(99)	
	<i>T. harzianum</i>	pachybasin (yellow)	chrysophanol (red)	(29, 100)	
<i>Curvularia</i>	<i>T. polysporum</i>	pachybasin (yellow)	-	(99, 101)	
	<i>T. viride</i>	pachybasin (yellow), 1,3,6,8-tetraHAQN, 2,4,5,7-tetraHAQN	emodin (yellow), chrysophanol (red)	(99, 102)	
	<i>C. lunata</i>	catenarin (red), erythroglauca (red), cynodontin (bronze), helminthosporin (maroon), tritispurin (reddish brown)	chrysophanol (red)	(3-5)	
	<i>Drechslera D. teres</i>	catenarin (red)	-	(103)	
<i>Herpotrichia</i>	<i>D. graminea</i>	catenarin (red)	-	(103)	
	<i>D. tritici-repentis</i>	catenarin (red)	-	(103)	
	<i>D. phlei</i>	catenarin (red)	-	(103)	
	<i>D. dactyloides</i>	catenarin (red)	-	(103)	
	<i>D. avenae</i>	cynodontin (bronze), helminthosporin (maroon)	-	(103)	
	<i>H. rhodosticta</i>	averythrin (orange), averythrin-6-monomethyl ether	-	(104)	
<i>Isaria I. farinosa</i>	red HAQN pigment (none completely characterized)	-	(105)		
<i>Fungi K_BK5</i>	austrorocortinin (red)	-	(27)		



which were fed a diet containing 0, 2.5 or 5.0% of colouring compounds extracted from madder roots clearly indicate that this dyestuff—rich in alizarin, lucidin-3-*O*-primeveroside and ruberythric acid, all classified in the ‘group E₃’ of HAQN and synthesized *via* the chorismate/*o*-succinylbenzoic acid pathway as mentioned above—exerts a carcinogenic potential in both the kidney and the liver, even with the lower dose of the study¹¹³. These studies support data in other previous studies^{114,115} and provide clear evidence that madder colour exerts unequivocal carcinogenicity against renal tubule cells and hepatocytes in rats. Therefore, the authors¹¹³ concluded that further studies on these individual HAQN components should be performed to clarify which anthraquinone is responsible for carcinogenicity.

Other recent studies indicate that the dark red purpurin pigment extracted from Indian madder (*Rubia cordifolia*)—classified in the ‘group E₃’ of HAQN dyes and synthesized *via* the chorismate/*o*-succinylbenzoic acid pathway—has an antimutagenic effect on the Ames *Salmonella* bacterial mutagenicity assay. The antigenotoxic effect was observed in *Drosophila melanogaster* against a range of environmental carcinogens. Inhibition of the formation of hepatic DNA adducts in male C57bl6 mice after a single dose of the heterocyclic amine dietary carcinogen Trp-P-2 (30 mg/kg) was observed by short-term dietary supplementation with purpurin¹¹⁶. In another study, purpurin was found to show inhibition of mutagenicity of a number of heterocyclic amines in the Ames mutagenicity test. The inhibition effect of purpurin was dependent upon pH, being better in neutral than acidic conditions¹⁵.

Concerning anthraquinoid pigments synthesized *via* the polyketide pathway, the orange pigment aloe-emodin (group A₁) induced micronucleus frequencies in the *in vitro* micronucleus test in mouse lymphoma L5178Y cells¹¹⁷. The emodin pigment (group A₁) has toxic and gene mutagenic properties. The activation mechanism of emodin into a direct mutagen to *Salmonella typhimurium* TA1537 was investigated by using the S9 and microsomes of rat livers. Emodin exhibited mutagenicity in the presence of NADPH or NADH¹¹⁸. Another study mentioned that emodin was clearly genotoxic in mouse lymphoma cells¹¹⁹. Emodin of fungal origin has been classified as diarrheagenic and genotoxic mycotoxin⁵⁸. Similarly, as fungal chrysophanol and physcion are hypothesized to exert genotoxicity, they are also considered as mycotoxins today⁹². Thus the detection of emodin, physcion and/or chrysophanol from some strains of *Aspergillus spp.*, *Penicillium spp.*, *Eurotium spp.*, *Dermocybe sanguinea*, *Dermocytes spp.*, *Trichoderma spp.* and *Curvularia lunata* (see Table 6) suggests that some of these fungi are potent mycotoxigenic.

5 Current industrial applications of natural hydroxyanthraquinoid dyestuffs

Traditionally, relevant hydroxyanthraquinoid dyestuffs such as the famous kermes parasite insect, the cochineal insect and the European madder root are essentially used to dye textiles. They are also used for other non-food applications, e.g. printing, cosmetics (hair colorants...) and pharmaceutical applications across the globe. They provide the most important red pigments used in artistic paintings. Hydroxyanthraquinoid red dyes were among the reds that dominated the dye markets of Europe. Natural hydroxyanthraquinoid pigments were

generally applied as isolated compounds or as glycosides. It was discovered that natural hydroxyanthraquinoid pigments, like alizarin from madder root and carminic acid from cochineal insect, coloured hair directly even at room temperature and that they were resistant to perspiration, washing, light and adverse weather conditions. Furthermore, hydroxyanthraquinoid pigments were very stable in solutions of the cosmetic media¹⁵.

Colour compounds extracted from the roots of European madder have been used as mentioned above in Japan as colorants for food, e.g. confectionery, boiled fish and soft drinks, but they are not allowed as a food additive in either the US or the EU. Only the natural red colorant ‘cochineal extract’ (additive E-120(ii)) which is an extract of the dried bodies of the female cochineal insect, with around 20% carminic acid content, is allowed and widely used as a colouring agent in food processes in the EU (at dosage levels from 50 to 500 mg/kg) and in the US (only up to 5 mg/kg)^{7,15,22,38}. Cochineal is commonly cultivated from the wild prickly pear cactus that grows thickly on the mountainsides in central Peru. Cochineal produces the pigment as a deterrent against other insects. The pigment can be obtained from the body and eggs of the insect. The few countries that produce commercial cochineal extracts are Peru, Mexico, the Canary Islands and, more recently, Chile and Bolivia. Only in Peru, the commercial production of cochineal extract is 200 ton/year, whereas in the Canary Islands production is only about 20 ton/year⁷. France is believed to be the world’s largest importer of cochineal extract, but Italy and Japan come next. The insects are killed by immersion in hot water or hot ethanol or by exposure to sunlight, steam, or by oven heat. Approximately 130,000 insects or 2 kg dry insects are required to produce 1 kg of cochineal extract and approximately 200 kg of dried insects are produced weekly at the largest cochineal farm⁷. Cochineal extract is very soluble in water and exhibits shade changes with changes in pH. At pH 4 and below, it is orange; it turns from violet to red by increasing pH from 5 to 7. Traditionally, cochineal extract is extracted with water or aqueous alcohol at 90 °C to 100 °C by batch or continuous process⁷. It is one of the few natural and water-soluble colorants that resist degradation with time. It has a good stability to heat, chemical oxidation, light and oxygen¹⁵. Often it is more stable than some synthetic food grade colorants but unstable at low pH. A water insoluble form of cochineal extract is commonly used to colour several food products, e.g. sausage products, bakery and dairy products, confectionery, and often competes with red root beet (betanin) and anthocyanins in food colouring. The water-soluble form of cochineal extract is currently used in beverages, soft and alcoholic drinks such as aperitifs (e.g. CampariTM). Cochineal extract is not kosher and is not vegetarian. Its main limitation in food application is its insolubility at low pH as mentioned above¹⁵. Carmine, *i.e.* the additive E-120(i), is a complex of carminic acid with various metals: an aluminium lake of carminic acid is currently being used in the commercial preparation of carmine^{7,15,22}. Variation in the ratio of carminic acid to aluminium produces a range of colours from pale strawberry to near black currant. Carmine is commonly traded as powder with a carminic acid content of 40% to 60% and liquid aqueous alkaline forms of carmine (and spray-dried derivatives) are also available with a carminic acid content of 2% to 7%. Cochineal extract and carmine are neither toxic nor known to be carcinogenic. Carmine is widely consumed in foods and beverages and has been rarely

implicated in adverse reactions. It can induce an anaphylactic-shock reaction in a small number of people, due to impurities in the preparation, not due to the pigment itself. In fact, colouring compounds in natural food grade colorants are small molecular weight, non-protein chemicals that cannot be expected to give true food allergies, either immunoglobulin E (IgE)-mediated or cell-mediated allergy. However, natural colorants, for example carmine, are often extracted from biological materials that may contain many other compounds, including proteins in addition to the colouring compounds. In 1998, it was reported that IgE-mediated allergy might be caused by the consumption of carmine, due to the presence of protein residues¹²⁰. Once IgE sensitization to these carmine proteins occurs, the level of exposure to these residual proteins through carmine-containing foods and beverages may be sufficient to elicit allergic reactions. For example, an anaphylactic reaction has been reported in a 34-year-old female atopic patient after ingestion of an orange beverage containing carmine. Symptoms like urticaria, rhinitis, nausea, vomiting, asthma, chills and diarrhea were observed. Skin prick tests carried out on the orange beverage, carmine and cosmetics containing the pigment were positive. In 1995, a reaction to carmine occurred in a 35-year-old woman after she ingested yoghurt that contained mixed fruits. Approximately 2 h after consumption she experienced symptoms of anaphylaxis including generalized urticaria, angioedema (localized swelling) and asthma¹²¹. In 1997, four adverse reactions following consumption of an alcoholic beverage containing carmine were reported in women ranging from 25 to 43 years old, with urticaria and angioedema. A skin prick test was performed and was found positive for carmine contained in the alcoholic beverage. Four instances of acute allergic reactions in a 28-year-old female after ingestion of orange beverage, strawberry milk and a red coloured cocktail containing carmine were also mentioned. An anaphylactic reaction in a 27-year-old woman has been reported after the consumption of a Popsicle coloured with carmine¹²². Carmine and cochineal extract are different from the azo pigment 'cochineal red A' (additive E-124) which is a synthetic colorant.

Concerning the natural food colorant Arpink red™, many toxicological data are also available: acute oral toxicity in mice of the pigment, 90-day subchronical toxicological study, acute dermal irritation/corrosion, acute eye irritation/corrosion, anti-tumour effectiveness, micronucleus test in mice, AMES test (*Salmonella typhimurium* reverse mutation assay), estimation of antibiotic activity, results of estimation of five mycotoxins. The fungal colorant gives a raspberry-red colour in an aqueous solution, stable at pH over 3.5. Neutral solutions are stable even after 30 min of boiling and colour shade does not change in relation with pH². After evaluating all the materials provided by the Ascolor Biotech s.r.o company, the Codex Alimentarius Commission (Rotterdam meeting, March 11–15, 2002) made the following statement: "there will not be any objections to use the red colouring matter Arpink red™"; The Arpink red™ colorant use was recommended as 100 mg/kg in meat products and in non-alcoholic drinks, 200 mg/kg in alcoholic drinks, 150 mg/kg in milk products including ice creams and 300 mg/kg in confectionery products². After the first approval by the Codex Alimentarius, the Arpink red™ safety assessment was discussed during the 63rd meeting of Joint FAO/WHO Expert Committee on Food Additives (JECFA) in Geneva, June 8–17, 2004. The red colorant received a two-year temporary approval by the EU for

distribution as a food additive, exclusively in the Czech Republic from 2004 to 2006². The file was still under progress at the European Food Safety Authority (EFSA) for some years. The situation now is not clear as Ascolor Biotech s.r.o. did not send data to authorities later on and seems to have closed its activities. Thus, there is no particular information on potential mycotoxin production and pathogenicity towards humans, despite the fact that the production of secalonic acid D, *i.e.* a pale yellow teratogenic mycotoxin, is well known from the fungus *Penicillium oxalicum*⁵. It has been shown that the biosynthesis of secalonic acid D (see Fig. 2 for the chemical structure) was dependent on the biosynthesis of the pigment emodin *via* the acetate-malonate pathway in a study conducted on the lichen *Laurera benguelensis*⁴¹. Diverse biological activities of secalonic acid D have been reported, such as a mycotoxin towards chicken and mice embryo, an inhibitor of various isozymes of protein kinase C and protein kinase A in murine secondary palate development¹²³, as well as mouse and human cleft palatal inducing agent¹²⁴.

In conclusion, this review provides relevant information regarding the properties of hydroxyanthraquinoid pigments, their biosynthetic pathway, their toxicity and carcinogenicity in recent decades. The collective information summarized in the review will act as an important segment for development of 'niche' fungal dyestuffs rich in hydroxyanthraquinoid pigments. These conclusions indicate that, even if the toxicological investigations of a new additive are not financially negligible, non-mycotoxigenic filamentous fungi such as strains of *Drechslera spp.*, *Herpotrichia spp.*, *Paecilomyces spp.* and *Isaria spp.* could be used for the production of dyestuffs rich in hydroxyanthraquinoid pigments as potent natural food grade colorants, with different shades according to the biomass composition: such as red (for main components catenarin & erythroglaucon), reddish brown (for tritispurin), bronze (for cynodontin), maroon (for helminthosporin) and orange-yellow (for pachybasin & averythrin). However, further studies should be performed on these fungal HAQN pigments to evaluate their potent carcinogenicity in humans from the food safety perspective. Current data on this topic are therefore insufficient.

Acknowledgments

The authors are grateful to Valérie QUILICI and Lynda GOYER from Université de La Réunion who helped in the preparation of this review.

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] Downham, A.; Collins, P. *Int. J. Food Sci. Technol.* **2000**, *35*, 5–22.
- [2] Dufossé, L. *Food Technol. Biotech.* **2006**, *44*, 313–321.
- [3] Räisänen, R. *Handbook of Natural Colorants*; John Wiley & Sons: Chichester, 2009; Chapter 11, pp 183–200.
- [4] Duran, N.; Maria, F. S. T.; Roseli, D. C.; Elisa, E. *Crit. Rev. Food. Sci. Nutr.* **2002**, *42*, 53–66.
- [5] Mapari, S. A. S.; Nielsen, K. F.; Larsen, T. O.; Frisvad, J. C.; Meyer, A. S.; Thrane, U. *Curr. Opin. Biotechnol.* **2005**, *16*, 231–

- 238.
- [6] Dufossé, L.; Galaup, P.; Yaron, A.; Arad, S. M.; Blanc, P.; Chidambara-Murthy, K. N.; Ravishankar, G. A. *Trends Food Sci. Tech.* **2005**, *16*, 389–406.
- [7] Chattopadhyay, P.; Chatterjee, S.; Sen, S. K. *Afr. J. Biotechnol.* **2008**, *7*, 2972–2985.
- [8] Mapari, S. A. S.; Meyer, A. S.; Thrane, U.; Frisvad, J. C. *Microbial. Cell. Factories.* **2009**, *8*, 24–28.
- [9] Blanc, P. J.; Loret, M. O.; Santerre, A. L.; Pareilleux, A.; Promé, D.; Promé, J. C. *J. Food Sci.* **1995**, *59*, 862–865.
- [10] Hajjaj, H.; Blanc, P. J.; Groussac, E.; Goma, G.; Uribealrea, J. L.; Loubiere, P. *Enzyme Microb. Technol.* **2000**, *27*, 619–625.
- [11] Carvalho, J. C.; Pandey, A.; Batitha, S.; Soccol, C. R. *Agro Food Ind Hi-tech.* **2003**, *14*, 37–42.
- [12] Liu, B. H.; Wu, T. S.; Su, M. C.; Chung, C. P.; Yu, F. Y. *J. Agric. Food Chem.* **2005**, *53*, 170–175.
- [13] Mapari, S. A. S.; Thrane, U.; Meyer, A. S. *Trends Biotechnol.* **2010**, *28*, 300–307.
- [14] Liu, R.; Lu, Y.; Wu, T.; Pan, Y. *Chromatographia* **2008**, *68*, 95–99.
- [15] Bechtold, T. *Handbook of Natural Colorants*; John Wiley & Sons: Chichester, 2009; Chapter 10, pp 151–182.
- [16] Alitheen, N. B.; Mashitoh, A. R.; Yeap, S. K.; Shuhaimi, M.; Abdul-Manaf, A.; Nordin, L. *Int. Food Res. J.* **2010**, *17*, 711–719.
- [17] Bonose-Crosnier de Bellaistre, M.; Nowik, W.; Tchaplá, A.; Heron, S. *J. Chromatogr.* **2011**, *1218*, 778–786.
- [18] Bick, I. R. C.; Rhee, C. *Biochem. J.* **1966**, *98*, 112–116.
- [19] De Santis, D.; Moresi, M. *Ind. Crop. Prod.* **2007**, *26*, 151–162.
- [20] Baker, R. A.; Tatum, J. H. *J. Ferment. Bioeng.* **1997**, *85*, 359–361.
- [21] Nagia, F. A.; El-Mohamedy, R. S. R. *Dyes Pigments* **2007**, *75*, 550–555.
- [22] Socaciu C. *Food Colorants—Chemical and functional properties*; CRC Press Taylor & Francis: New York, 2008.
- [23] Drivas, I.; Blackburn, R. S.; Rayner, C. M. *Dyes Pigments* **2011**, *88*, 7–17.
- [24] Liu, S. Y.; Sporer, F.; Wink, M.; Jourdane, J.; Henning, R.; Li, Y. L.; Ruppel, A. *Trop. Med. Int. Health* **1997**, *2*, 179–188.
- [25] Locatelli, M.; Tammaro, F.; Menghini, L.; Carlucci, G.; Epifano, F.; Genovese, S. *Phytochem. Lett.* **2009**, *2*, 223–226.
- [26] Kremer, D.; Kosalec, I.; Locatelli, M.; Epifano, F.; Genovese, S.; Carlucci, G.; Zovko Koncic, M. *Food Chem.* **2012**, *131*, 1174–1180.
- [27] Muangsin, N.; Wisetsakdakorn, W.; Chaichit, N.; Sihanonth, P.; Petsom, A.; Sangvanich, P. *Dyes Pigments* **2008**, *77*, 653–656.
- [28] Tanaka, C.; Miyagawa, H.; Kuwahara, Y.; Tsuda, M. *Mycoscience* **2002**, *43*, 317–320.
- [29] Liu, S. Y.; Lo, C. T.; Chen, C.; Liu, M. Y.; Chen, J. H.; Peng, K. C. *J. Biochem. Biophys. Methods* **2007**, *70*, 391–395.
- [30] Yadav, J. P.; Arya, V.; Panghal, M.; Kumar, S.; Dhankhar, S. *Fitoterapia* **2010**, *81*, 223–230.
- [31] Zhang, L. M.; Xie, W. G.; Wen, T. T. *J. Therm. Anal. Calorim.* **2010**, *100*, 215–218.
- [32] Gill, M.; Steglich, W. *Progress in the Chemistry of Organic Natural Products*; Springer: New York, 1987; Chapter 51, pp 125–174.
- [33] Manojlovic, I.; Bogdanovic-Dusanovic, G.; Gritsanapan, W.; Manojlovic, N. *Chem. Pap.* **2006**, *60*, 466–468.
- [34] Orban, N.; Boldizar, I.; Szücs, Z.; Danos, B. *Dyes Pigments* **2008**, *77*, 249–257.
- [35] Smetanina, O. F.; Kalinovskii, A. I.; Khudyakova, Y. V.; Slinkina, N. N.; Pivkin, M. V.; Kuznetsova, T. A. *Chem. Nat. Compd.* **2007**, *43*, 327–329.
- [36] Anke, H.; Kolthoum, I.; Zihner, H.; Laatsch, H. *Arch. Microbiol.* **1980**, *126*, 223–230.
- [37] Ackacha, M. A.; Polec-Pawlak, K.; Jarosz, M. *J. Sep. Sci.* **2003**, *26*, 1028–1034.
- [38] Banyai, P.; Kuzovkina, I. N.; Kursinszki, L.; Szöke, E. *Chromatographia* **2006**, *63*, 111–114.
- [39] Jasril; Lajis, N. H.; Mooi, L. Y.; Abdullah, M. A.; Sukari, M. A.; Ali, A. M. *Asia Pacific J. Mol. Biol. Biotechnol.* **2003**, *11*, 3–7.
- [40] Bao, L.; Liu, L.; Wu, Y.; Han, T.; Xue, L.; Zhang, Q. *Chem-Biol. Interact.* **2011**, *194*, 97–105.
- [41] Manojlovic, N. T.; Vasiljevic, P. J.; Gritsanapan, W.; Supabphol, R.; Manojlovic, I. *Biol. Res.* **2010**, *43*, 169–176.
- [42] Tanaka, C.; Miyagawa, H.; Kuwahara, Y.; Tsuda, M. *Mycoscience* **2002**, *43*, 317–320.
- [43] Tang, T.; Yin, L.; Yang, J.; Shan, G. *Eur. J. Pharmacol.* **2007**, *567*, 177–185.
- [44] Schweppe, H. *Historic Textile and Paper Materials II*; American Chemical Society: Washington, 1989; Chapter 13, pp 188–219.
- [45] Rafaëilly, L.; Héron, S.; Nowik, W.; Tchaplá, A. *Dyes Pigments* **2008**, *77*, 191–203.
- [46] Mishchenko, N. P.; Stepanenko, L. S.; Krivoshchekova, O. E.; Maksimov, O. B. *Chem. Nat. Compd.* **1980**, *16*, 117–121.
- [47] Borroto, J.; Coll, J.; Rivas, M.; Blanco, M.; Concepcion, O.; Tandon, Y. A.; Hernandez, M.; Trujillo, R. *Plant Cell Tiss. Org.* **2008**, *94*, 181–187.
- [48] Yamamoto, Y.; Matsubara, H.; Kinoshita, Y.; Kinoshita, K.; Koyama, K.; Takahashi, K.; Ahmadjam, V.; Kurokawa, T.; Yoshimura, I. *Phytochemistry* **1996**, *43*, 1239–1242.
- [49] Gagunashvili, A. N.; Davidsson, S. P.; Jonsson, Z. O.; Andersson, O. S. *Mycol. Res.* **2009**, *113*, 354–363.
- [50] Canamares, M. V.; Garcia-Ramos, J. V.; Domingo, C.; Sanchez-Cortes, S. *Vib. Spectrosc.* **2006**, *40*, 161–167.
- [51] Bouras, N.; Strelkov, S. E. *Physiol. Mol. Plant P.* **2008**, *72*, 87–95.
- [52] Birkinshaw, J. H.; Gourlay, R. *Biochem. J.* **1961**, *80*, 387–392.
- [53] Miliari, C.; Romani, A.; Favaro, G. *J. Phys. Org. Chem.* **2000**, *13*, 141–150.
- [54] Van den Berg, A. J. J.; Labadie, R. P. *Methods in Plant Biochemistry*; Academic press: London, 1989; Vol. 1, pp 451–491.
- [55] Bringmann, G.; Gulder, T. A. M.; Hamm, A.; Goodfellow, M.; Fiedler, H. P. *Chem. Commun.* **2009**, *44*, 6810–6812.
- [56] Leistner, E. *Primary and secondary metabolism of plant cell cultures*; Springer: Berlin, 1985; pp 215–224.
- [57] Han, Y. S.; Van der Heijden, R.; Lefeber, A. W. M.; Erkelens, C.; Verpoorte, R. *Phytochemistry* **2002**, *59*, 45–55.
- [58] Izhazi, I. *New Phytol.* **2002**, *155*, 205–217.
- [59] Angelini, L. G.; Pistelli, L.; Belloni, P.; Bertoli, A.; Panconesi, S. *Ind. Crop. Prod.* **1997**, *6*, 303–311.
- [60] Kaur, P.; Chandel, M.; Kumar, S.; Kumar, N.; Singh, B.; Kaur, S. *Food Chem. Toxicol.* **2010**, *48*, 320–325.
- [61] Morimoto, M.; Tanimoto, K.; Sakatani, A.; Komai, K. *Phytochemistry* **2002**, *60*, 163–166.
- [62] El-Gamal, A. A.; Takeya, K.; Itokawa, H.; Halim, A. F.; Amer, M. M.; Saad, H. E. A.; Awad, S. A. *Phytochemistry* **1995**, *40*, 245–251.
- [63] Banthorpe, D. V.; White, J. J. *Phytochemistry* **1995**, *38*, 107–111.
- [64] Koyama, J.; Ogura, T.; Tagahara, K. *Phytochemistry* **1993**, *33*, 6, 1540–1542.
- [65] Sang, S.; Cheng, X.; Zhu, N.; Stark, R. E.; Badmaev, V.; Ghai, G.; Rosen, R. T.; Ho, C. T. *J. Agric. Food Chem.* **2001**, *49*, 4478–4481.
- [66] Wijnsma, R.; Verpoorte, R.; Mulder-Krieger, T.; Baerheim Svendsen, A. *Phytochemistry* **1984**, *23*, 2307–2311.
- [67] Robins, R. I.; Payne, J.; Rhodes, M. J. C. *Phytochemistry* **1986**, *25*, 2327–2334.
- [68] Wijnsma, R.; Go, J. T. K. A.; Harkes, P. A. A.; Verpoorte, R.; Baerheim Svendsen, A. *Phytochemistry* **1986**, *25*, 1123–1126.
- [69] Khouri, H. E.; Ibrahim, R. K. *Phytochemistry* **1987**, *26*, 2531–2535.
- [70] Schripsema, J.; Ramos-Valdivia, A.; Verpoorte, R. *Phytochemistry* **1999**, *51*, 55–60.
- [71] Arun, P.; Purushotham, K. G.; Jonhsy Jayarani, J.; Vasantha Kumari. *J. Pharm. Sci. Technol.* **2010**, *2*, 198–201.

- [72] Siva, R.; Mayes, S.; Behera, S. K.; Rajasekaran, C. *Ind. Crop. Prod.* **2012**, *37*, 415–419.
- [73] Badr, J. M. *Nat. Prod. Sci.* **2008**, *14*, 227–232.
- [74] Liu, B.; Ge, X.; He, Y.; Xie, J.; Xu, P.; He, Y.; Zhou, Q.; Pan, L.; Chen, R. *Aquaculture* **2010**, *310*, 13–19.
- [75] Li, L.; Zhang, C.; Xiao Y. Q.; Chen, D. D.; Tian, G. F.; Wang, Y. *Chin. J. Nat. Med.* **2011**, *9*, 410–413.
- [76] Malik, S.; Sharma, N.; Sharma, U. K.; Singh, N. P.; Blushan, S.; Sharma, M.; Sinha, A. K.; Ahuja, P. S. *J. Plant Physiol.* **2010**, *167*, 749–756.
- [77] Baskan, S.; Daut-Özdemir, A.; Günaydin, K.; Erim, F. B. *Talanta* **2007**, *71*, 747–750.
- [78] Choi, G. J.; Lee, S. W.; Jang, K. S.; Kim, J. S.; Cho, K. Y.; Kim, J. C. *Crop Prot.* **2004**, *23*, 1215–1221.
- [79] Lee, N. J.; Choi, J. H.; Koo, B. S.; Ryu, S. Y.; Han, Y. H.; Lee, S. I.; Lee, D. U. *Biol. Pharm. Bull.* **2005**, *28*, 2158–2161.
- [80] Fairbairn, J. W.; El-Muhtadi, F. J. *Phytochemistry* **1972**, *11*, 263–268.
- [81] Genovese, S.; Tammara, F.; Menghini, L.; Carlucci, G.; Epifano, F.; Locatelli, M. *Phytochem. Anal.* **2010**, *21*, 261–267.
- [82] Panichayupakaranant, P.; Sakunpak, A.; Sakunphueak, A. *J. Chromatogr. Sci.* **2009**, *47*, 197–200.
- [83] Dagne, E.; Yenesew, A.; Asmellash, S.; Demissew, S.; Mavi, S. *Phytochemistry* **1994**, *35*, 401–406.
- [84] Van Wyk, B. E.; Yenesew, A.; Dagne, E. *Biochem. Syst. Ecol.* **1995**, *23*, 267–275.
- [85] Mohanlall, V.; Steenkamp, P.; Odhav, B. *J. Med. Plants Res.* **2011**, *5*, 3132–3141.
- [86] Rosso, M. L.; Bertoni, M. D.; Adler, M. T.; Maier, M. S. *Biochem. Syst. Ecol.* **2003**, *31*, 1197–1200.
- [87] Manojlovic, N.; Markovic, Z.; Gritsanapan, W.; Boonpragob, K. *Russ. J. Phys. Chem.* **2009**, *83*, 1554–1557.
- [88] Piatelli, M.; Giudici de Nicola, M. *Phytochemistry* **1988**, *7*, 1183–1187.
- [89] Nakano, H.; Komiya, T.; Shibata, S. *Phytochemistry* **1972**, *11*, 3505–3508.
- [90] Sochting, U.; Fröden, P. *Mycol. Prog.* **2002**, *1*, 257–266.
- [91] Cohen, P. A.; Neil Towers, G. H. *Phytochemistry* **1996**, *42*, 1325–1329.
- [92] Frisvad, J. C. *Arch. Environ. Contam. Toxicol.* **1989**, *18*, 452–467.
- [93] Mendez, A.; Perez, C.; Montanez, J.; Martinez, G.; Aguilar, C. N. *J. Zhejiang Univ-Sc. B* **2011**, *12*, 961–968.
- [94] Podojil, M.; Sedmera, P.; Vokocun, J.; Betina, V.; Barathov, H.; Durackov, K.; Horakova, K.; Nemecek, P. *Folia Microbiol.* **1979**, *23*, 438–443.
- [95] Baker, R. A.; Tatum, J. H. *J. Ferment. Bioeng.* **1998**, *85*, 359–361.
- [96] Medentsev, A. G.; Akimenko, V. K. *Phytochemistry* **1998**, *47*, 935–959.
- [97] Nagia, F. A.; El-Mohamedy, R. S. R. *Dyes Pigments* **2007**, *75*, 550–555.
- [98] Gill, M.; Morgan, P. M. *ARKIVOC* **2001**, 145–156.
- [99] Reino, J. L.; Guerrero, R. F. *Phytochem. Rev.* **2008**, *7*, 89–123.
- [100] Vinale, F.; Marra, R.; Scala, F.; Ghisalberti, E. L.; Lorito, M.; Sivasithamparam, K. *Lett. Appl. Microbiol.* **2006**, *43*, 143–148.
- [101] Donnelly, D. M. X.; Helen, M. S. *Phytochemistry* **1986**, *25*, 2303–2304.
- [102] Slater, G. P.; Haskins, R. H.; Hogge, L. R.; Nesbitt, L. R. *Can. J. Chem.* **1967**, *45*, 92–96.
- [103] Engström, K.; Brishammar, S.; Svensson, C.; Bengtsson, M.; Andersson, R. *Mycol. Res.* **1993**, *97*, 381–384.
- [104] Van Eljk, G. W.; Roelijmans, H. J. *Exp. Mycol.* **1984**, *8*, 266–268.
- [105] Cho, Y. J.; Park, J. P.; Hwang, H. J.; Kim, S. W.; Choi, J. W. *J. Biotechnol.* **2002**, *95*, 13–23.
- [106] Velmurugan, P.; Lee, Y. H.; Nanthakumar, K.; Kamala-Kannan, S.; Dufossé, L.; Mapari, S. A. S.; Oh, B. T. *J. Basic Microbiol.* **2010**, *50*, 1–10.
- [107] Wang, Y.; Chen, J. W.; Ge, L. K.; Wang, D. G.; Cai, X. Y.; Huang, L. P.; Hao, C. *Dyes Pigments* **2009**, *83*, 276–280.
- [108] Barbee, G. C.; Santer, M. M.; McClain, W. R. *Crop Prot.* **2010**, *29*, 506–508.
- [109] Kawasaki, Y.; Goda, Y.; Yoshihira, K. *Chem. Pharm. Bull. (Tokyo)* **1992**, *40*, 1504–1509.
- [110] Marec, F.; Kollarova, I.; Jegorov, A. *Planta Med.* **2001**, *67*, 127–131.
- [111] Inoue, K.; Shibutani, M.; Masutomi, N.; Toyoda, K.; Takagi, H.; Uneyama, C.; Nishikawa, A.; Hirose, M. *Food Chem. Toxicol.* **2008**, *46*, 241–252.
- [112] Yokohira, M.; Yamakawa, K.; Hosokawa, K.; Matsuda, Y.; Kuno, T.; Saoo, K.; Imaida, K. *J. Food Sci.* **2008**, *73*, T26–T32.
- [113] Kaoru, I.; Midori, Y.; Miwa, T.; Makoto, S.; Hironori, T.; Masao, H.; Akiyoshi, N. *Food Chem. Toxicol.* **2009**, *47*, 184–191.
- [114] Westendorf, J.; Marquardt, H.; Poginsky, B.; Dominiak, M.; Schmidt, J.; Marquardt, H. *Mutat Res.* **1990**, *240*, 1–12.
- [115] Westendorf, J.; Pfau, W.; Schulte, A. *Carcinogenesis* **1998**, *19*, 2163–2168.
- [116] Marczylo, T.; Sugiyama, C.; Hayatsu, H. *J. Agric. Food Chem.* **2003**, *51*, 3334–3337.
- [117] Mueller, S. O.; Stopper, H.; Dekant, W. *Drug Metab. Dispos.* **1998**, *26*, 540–546.
- [118] Masuda, T.; Ueno, Y. *Mutat Res.* **1984**, *125*, 135–144.
- [119] Mueller, S. O.; Lutz, W. K.; Stopper, H. *Mutat. Res.* **1998**, *414*, 125–129.
- [120] Acero, S.; Tabar, A. I.; Alvarez, M. J.; Garcia, B. E.; Olaguibel, J. M.; Moneo, I. *Allergy* **1998**, *53*, 897–901.
- [121] Beaudouin, E.; Kanny, G.; Lambert, H.; Fremont, S.; Moneret-Vautrin, D. *Ann. Allergy Asthma Immunol.* **1995**, *74*, 427–430.
- [122] Baldwin, J. L.; Chou, A. H.; Solomon, W. R. *Ann. Allergy Asthma Immunol.* **1997**, *79*, 415–419.
- [123] Hanumegowda, U. M.; Judy, B. M.; Welshons, W. V.; Reddy, C. S. *Toxicol. Sci.* **2002**, *66*, 159–165.
- [124] Dhulipala, V. C.; Maddali, K. K.; Welshons, W. V.; Reddy, C. S. *Birth Defects Res. B Dev. Reprod. Toxicol.* **2005**, *74*, 233–242.