

Nonpeptidic urotensin-II receptor antagonists I: *in vitro* pharmacological characterization of SB-706375

*¹Stephen A. Douglas, ¹David J. Behm, ¹Nambi V. Aiyar, ¹Diane Naselsky, ¹Jyoti Disa, ¹David P. Brooks, ¹Eliot H. Ohlstein, ²John G. Gleason, ³Henry M. Sarau, ³James J. Foley, ³Peter T. Buckley, ³Dulcie B. Schmidt, ³William E. Wixted, ⁴Katherine Widdowson, ⁵Graham Riley, ⁶Jian Jin, ⁷Timothy F. Gallagher, ⁷Stanley J. Schmidt, ⁷Lance Ridgers, ⁷Lisa T. Christmann, ⁷Richard M. Keenan, ⁷Steven D. Knight & ⁷Dashyant Dhanak

¹CVU Department of Biology, Cardiovascular and Urogenital and Respiratory and Inflammation Centers of Excellence for Drug Discovery, GlaxoSmithKline, 709 Swedeland Road, UW2510 King of Prussia, PA 19406-0939, U.S.A.; ²CVU Department of Medicinal Chemistry, Cardiovascular and Urogenital and Respiratory and Inflammation Centers of Excellence for Drug Discovery, GlaxoSmithKline, 709 Swedeland Road, King of Prussia, PA 19406-0939, U.S.A.; ³RIRP Department of Biology, Cardiovascular and Urogenital and Respiratory and Inflammation Centers of Excellence for Drug Discovery, GlaxoSmithKline, 709 Swedeland Road, King of Prussia, PA 19406-0939, U.S.A.; ⁴RIRP Department of Medicinal Chemistry, Cardiovascular and Urogenital and Respiratory and Inflammation Centers of Excellence for Drug Discovery, GlaxoSmithKline, 709 Swedeland Road, King of Prussia, PA 19406-0939, U.S.A.; ⁵Assay Development and Compound Profiling, Discovery Research, GlaxoSmithKline, New Frontiers Park (North), Third Avenue, Harlow, Essex CM19 5AW; ⁶High Throughput Chemistry, Discovery Research, GlaxoSmithKline, Collegeville, PA 19426, U.S.A. and ⁷Medicinal Chemistry, Microbial, Musculoskeletal and Proliferative Diseases Center of Excellence for Drug Discovery, GlaxoSmithKline, Collegeville, PA 19426, U.S.A.

1 SB-706375 potently inhibited [¹²⁵I]hU-II binding to both mammalian recombinant and 'native' UT receptors (K_i 4.7 ± 1.5 to 20.7 ± 3.6 nM at rodent, feline and primate recombinant UT receptors and K_i 5.4 ± 0.4 nM at the endogenous UT receptor in SJRH30 cells).

2 Prior exposure to SB-706375 (1 μM, 30 min) did not alter [¹²⁵I]hU-II binding affinity or density in recombinant cells (K_D 3.1 ± 0.4 vs 5.8 ± 0.9 nM and B_{max} 3.1 ± 1.0 vs 2.8 ± 0.8 pmol mg⁻¹) consistent with a reversible mode of action.

3 The novel, nonpeptidic radioligand [³H]SB-657510, a close analogue of SB-706375, bound to the monkey UT receptor (K_D 2.6 ± 0.4 nM, B_{max} 0.86 ± 0.12 pmol mg⁻¹) in a manner that was inhibited by both U-II isopeptides and SB-706375 (K_i 4.6 ± 1.4 to 17.6 ± 5.4 nM) consistent with the sulphonamides and native U-II ligands sharing a common UT receptor binding domain.

4 SB-706375 was a potent, competitive hU-II antagonist across species with pK_b 7.29–8.00 in HEK293-UT receptor cells (inhibition of [Ca²⁺]_i-mobilization) and pK_b 7.47 in rat isolated aorta (inhibition of contraction). SB-706375 also reversed tone established in the rat aorta by prior exposure to hU-II (K_{app} ~ 20 nM).

5 SB-706375 was a selective U-II antagonist with ≥100-fold selectivity for the human UT receptor compared to 86 distinct receptors, ion channels, enzymes, transporters and nuclear hormones (K_i/IC_{50} > 1 μM). Accordingly, the contractile responses induced in isolated aortae by KCl, phenylephrine, angiotensin II and endothelin-1 were unaltered by SB-706375 (1 μM).

6 In summary, SB-706375 is a high-affinity, surmountable, reversible and selective nonpeptide UT receptor antagonist with cross-species activity that will assist in delineating the pathophysiological actions of U-II in mammals.

British Journal of Pharmacology (2005) **145**, 620–635. doi:10.1038/sj.bjp.0706229

Published online 25 April 2005

Keywords: Urotensin; UT receptor; SB-706375; SB-657510; G-protein-coupled receptor; vasoconstriction; GPR-14; hypertension; heart failure; radioligand binding

Abbreviations: ATCC, American Type Culture Collection; BCA, bicinchoninic acid; BSA, bovine serum albumin; [Ca²⁺]_i, intracellular calcium; CHO cell, Chinese hamster ovary cell; DMSO, dimethylsulphoxide; DPBS⁺, Dulbecco's phosphate-buffered saline; EDTA, ethylenediaminetetraacetic acid; FLIPR, fluorometric imaging plate reader; HEK293 cells, human embryonic kidney cells; SB-706375, 2-bromo-4,5-dimethoxy-*N*-[3-(*R*)-1-methyl-pyrrolidin-3-yloxy]-4-trifluoro-methyl-phenyl]-benzenesulphonamide HCl; SB-657510, 2-bromo-*N*-[4-chloro-3-(*R*)-1-methyl-pyrrolidin-3-yloxy]-phenyl]-4,5-dimethoxybenzenesulphonamide HCl; s.e.m., standard error of the mean; SNP, single nucleotide polymorphism; SPA, scintillation proximity assay; (h)U-II, (human) urotensin-II; (h)UT receptor, (human) urotensin-II receptor; WGA, wheat germ agglutinin-coated

Introduction

Human urotensin-II (hU-II) induces profound cardiohaemodynamic effects upon systemic administration in the cat (Behm

*Author for correspondence: E-mail: steve.a.douglas@gsk.com

et al., 2004a), monkey (Ames *et al.*, 1999) and in man (Böhm & Pernow, 2002; Lim *et al.*, 2004). Not only does hU-II constitute 'the most potent mammalian vasoconstrictor identified to date' (Ames *et al.*, 1999), it also influences cardiorenal function by acting as a potent regulator of cardiac contractility (Russell *et al.*, 2003; Kompa *et al.*, 2004), a natriuretic factor (Song *et al.*, 2003) and as a hypertrophic/proinflammatory factor (Watanabe *et al.*, 2001; Zou *et al.*, 2001; Tzanidis *et al.*, 2003; Johns *et al.*, 2004; Onan *et al.*, 2004). As such, hU-II and its G-protein-coupled receptor, UT, are purported to be involved in the (dys)regulation of cardiorenal function (Douglas, 2003; Douglas *et al.*, 2004a).

Based on a series of clinical observations, such as augmented plasma/tissue 'U-II-like' activity and pharmacogenetic associations (i.e. SNP analysis), hU-II has been implicated recently in the aetiology of numerous cardiorenal and metabolic diseases including hypertension (Matsushita *et al.*, 2001; Cheung *et al.*, 2004), heart failure (Douglas *et al.*, 2002; Ng *et al.*, 2002; Richards *et al.*, 2002; Russell *et al.*, 2003; Lapp *et al.*, 2004), atherosclerosis (Bousette *et al.*, 2004; Maguire *et al.*, 2004), renal failure (Totsune *et al.*, 2001; Shenouda *et al.*, 2002; Langham *et al.*, 2004) and diabetes (Totsune *et al.*, 2003; 2004; Wenyi *et al.*, 2003). Unfortunately, however, the lack of suitable UT receptor antagonists (Dhanak *et al.*, 2003; Douglas *et al.*, 2004a) has, thus far, precluded a detailed investigation of the specific role of the U-II/UT receptor system in the pathogenesis of such disorders, either in preclinical species or in man. Although several novel peptidic and nonpeptidic UT receptor ligands have been described in the medical and patent literature, they are of limited functional utility due to poor potency, limited selectivity and/or retention of intrinsic activity/agonism, etc. To address these issues, the present report describes the identification and characterization of SB-706375 (Figure 1a), an arylsulphonamide developed from high-throughput screening leads originally identified in the legacy SmithKline Beecham compound collection. SB-706375 is a potent, surmountable, reversible and selective mammalian UT receptor antagonist. Since SB-706375 exhibits 'pan-species' antagonism, it is proposed that this small molecule inhibitor will serve as a useful agent in delineating the (patho)physiological actions of U-II.

Methods

Radioligand binding studies

The pharmacological properties of SB-706375 were assessed in a variety of radioligand binding studies (intact cells and

membranes) using either (a) HEK293 cells stably expressing recombinant mammalian (mouse, rat, cat, monkey or human) UT receptors or (b) 'native' human cells expressing endogenous UT receptors (human rhabdomyosarcoma SJRH30 cells; Douglas *et al.*, 2004b).

Recombinant HEK293-UT receptor membrane preparation HEK293 cells expressing the UT receptor were detached from 150 cm² flasks with 1 mM EDTA in Ca²⁺/Mg²⁺-free Dulbecco's phosphate-buffered saline (DPBS), washed by centrifugation at 300 × *g* and stored as frozen pellets. Cell pellets were suspended in ice-cold buffer (10 mM Tris-HCl [pH 7.4], 5 mM Na-EDTA, 0.1 mM phenylmethylsulphonylfluoride [PMSF], 1.0 mg ml⁻¹ bacitracin, 0.1 mg ml⁻¹ aprotinin) and homogenized using a Dounce homogenizer (Bellco Glass, Inc., Vineland, NJ, U.S.A.). Homogenates were centrifuged at 47,000 × *g* for 20 min at 4°C. The pellets were washed twice by centrifugation in buffer (25 mM Tris-HCl [pH 7.4], 5 mM MgCl₂, 2 mM Na-EGTA, 0.1 mg ml⁻¹ bacitracin) and resuspended at 5 mg ml⁻¹ for storage at -70°C. Protein concentration was measured by the Pierce (Rockford, IL, U.S.A.) bicinchoninic acid (BCA) method using bovine serum albumin (BSA) as a standard.

Saturation binding studies in recombinant HEK293-UT cell membranes Saturation binding (*K_D*, *B_{max}* determination) was performed by scintillation proximity assay (SPA). Conditions were optimized for membrane protein concentration and amount of wheat germ agglutinin-coated (WGA) SPA beads (Amersham, Arlington Heights, IL, U.S.A.) and binding was carried out with 20–600 pM [¹²⁵I]hU-II either in the absence (total binding) or presence (nonspecific binding) of 1 μM cold hU-II. The apparent equilibrium dissociation constants (*K_D*) and the maximum binding sites (*B_{max}*) from saturation binding experiments were calculated using the interactive nonlinear curve-fitting program of GraphPad Prism (San Diego, CA, U.S.A.).

Competition binding studies in recombinant HEK293-UT cell membranes Competition binding was performed under similar conditions as those described above. [¹²⁵I]hU-II (300 pM) was incubated with cell membranes in the presence of varying concentrations of U-II isoforms (1.0 pM–1.0 μM in 0.1% BSA) or SB-706375 (0.1 nM–10 μM in DMSO vehicle). WGA-SPA beads were suspended (50 mg ml⁻¹) in binding buffer (25 mM Tris-HCl [pH 7.4], 5 mM MgCl₂, 0.1% BSA) and stored at 4°C. At the time of the assay, WGA-SPA beads (12.5 μg ml⁻¹) and recombinant UT receptor HEK293 membranes (50 μg well⁻¹) were precoupled by gentle shaking (1 h,




Figure 1 Structures of the novel nonpeptide sulphonamide antagonists (a) SB-706375 (2-bromo-4,5-dimethoxy-*N*-[3-(*R*)-1-methylpyrrolidin-3-yloxy]-4-trifluoromethyl-phenyl]-benzenesulphonamide HCl) and (b) SB-657510 (2-bromo-*N*-[4-chloro-3-((*R*)-1-methylpyrrolidin-3-yloxy)-phenyl]-4,5-dimethoxybenzenesulphonamide HCl).

25°C). Following preincubation, 100 µl of the complex, 10 µl of SB-706375 (0.1 nM–10 µM) and 50 µl [¹²⁵I]hU-II (0.3 nM) were added to each well of a Packard OptiPlate™-96 microtitre plate (along with adequate binding buffer to bring the final volume of each well to 200 µl) using a Packard MultiProbe II EX robotic liquid handling system (Perkin-Elmer, Shelton, CT, U.S.A.). The assay plates were then sealed and shaken gently on an orbital shaker (1 h, 25°C). Finally, plates were spun at 1500 × *g* for 10 min and cell-bound radioactivity was determined (Packard Top Count). Specific binding was determined using cold hU-II (1 µM).

Reversibility of binding in HEK293 cells stably expressing the monkey recombinant UT receptor Membranes from HEK293-monkey UT receptor cells were incubated with DMSO vehicle or SB-706375 (1 µM) for 30 min at 25°C following which *K_D* and *B_{max}* values were determined. Incubation mixtures were diluted with cold buffer consisting of 25 mM Tris-HCl, pH 7.4, 5 mM MgCl₂, 2 mM Na-EGTA and 0.1 mg ml⁻¹ bacitracin followed by centrifugation at 47,000 × *g* for 20 min at 4°C. Membrane pellets were then washed one more time and resuspended in 2 ml buffer and utilized for [¹²⁵I]hU-II competition binding as detailed above.

Analysis of SB-706375 binding to native UT receptors in human intact SJRH30 cells Binding of [¹²⁵I]hU-II to native human UT receptors was studied according to Douglas *et al.* (2004b) using a whole-cell binding assay format in the human rhabdomyosarcoma cell line, SJRH30 (American Type Culture Collection number CRL-2061, ATCC, Manassas, VA, U.S.A.). Briefly, SJRH30 cells were washed with DPBS⁺ buffer (with 10 mM MgCl₂, 0.7 mM CaCl₂, 1.4 mM glucose, 0.2% BSA) immediately prior to exposure (37°C for 30 min in 1 ml DPBS⁺) to [¹²⁵I]hU-II. After incubation, cells were washed four times with cold DPBS⁺ (1 ml), solubilized with 1 M NaOH (1 ml) and transferred to 12 × 75 mm glass tubes. Radioactivity was then measured in a Packard gamma counter (>85% efficiency). Saturation binding was carried out by incubating the cells with 5–300 pM [¹²⁵I]hU-II in the absence (total binding) or presence of 1 µM hU-II (nonspecific binding). Competition binding was performed under similar conditions using 200 pM [¹²⁵I]hU-II and different concentrations of competing ligands. All assays were performed in duplicate.

Inhibition of [³H]SB-657510 binding to recombinant HEK293-UT cell membranes [³H]SB-657510 (specific activity 87 Ci mmol⁻¹; Figure 1b) saturation binding to monkey recombinant UT-HEK293 cell membranes (10–15 µg) was initiated by the addition of increasing concentrations of radioligand in the absence (total binding) or presence (nonspecific binding) of hU-II (1 µM). Binding was performed at 25°C for 60 min (200 µl total volume of binding buffer; 20 mM Tris-HCl buffer at pH 7.4 containing 5 mM MgCl₂, 0.2% BSA, 0.1 mg ml⁻¹ bacitracin). Competition binding experiments were performed in duplicate using 10 nM [³H]SB-657510 and increasing concentrations of cold ligands. At the end of incubation, the reaction mixture was rapidly diluted with 2 ml cold wash buffer (0.9% NaCl w v⁻¹) followed by rapid filtration over Skatron filtermates (Skatron Instruments, Norway). Radioactivity was counted in a beta liquid scintillation counter.

In vitro selectivity profile of SB-706375

The selectivity of SB-706375 for the UT receptor was assessed by examining the interaction between this sulphonamide ligand and a total of 86 distinct G-protein-coupled receptors, ion channels, enzymes, transporters and nuclear hormone receptor cross-screening assays using established protocols.

G-protein-coupled receptor targets screened included adenosine (A_{1/2A/3}), adrenergic (α_{1A/1B/2B/2A/2C}, β_{1/2/3}), dopamine (D_{1/2/3/4}), muscarinic (M_{1/2/3/4}), serotonin (5-HT_{1A/B/D/E/F}, HT_{2A/B/C}, HT₃, HT₄, HT_{5A}, HT₆, HT₇), histamine (H_{1/2}), angiotensin II (AT_{1/2}), ANP (GC-A), bradykinin (B_{1/2}), calcitonin gene-related peptide (CGRP), melanocortin (MC₄R), melatonin (ML₁), neuropeptide Y (Y_{1/2}), endothelin (ET_{A/B}), neurotensin (NT₁), cholecystokinin (CCK_A), imidazoline (I₂), neurokinin (NK_{2/3}), benzodiazepine (BZD, central), cannabinoid (CB_{1/2}), nicotinic (αBTX-sensitive, neuronal N-type) leukotriene (BLT1), galanin (GalR2), Il-8 (CXCR2), somatostatin (sst_{3/4}), vasoactive intestinal peptide (PAC₁, VPAC₁), opioid (μ, δ, κ, σ), vasopressin (V_{1a}), P_{2X}, GABA_A, AMPA, kainate and NMDA receptors. SB-706375 was also evaluated in a number of nuclear hormone receptor assays (glucocorticoid, estrogen (α,β), progesterone and testosterone) along with a variety of ion channel (including sodium [sites 1 and 2], calcium [L-type], potassium [SK_{Ca}, K_v] and chloride [picrotoxinin-sensitive] channels), enzyme (including phosphodiesterases [isozymes I–V], phospholipase A₂, eNOS, elastase, protein kinase C, Na⁺/K⁺-ATPase and EGF-tyrosine kinase) and biogenic amine uptake (noradrenaline and dopamine transporters) assays.

Evaluation of SB-706375 as a mammalian UT receptor antagonist

The ability of SB-706375 to function as a competitive U-II antagonist was assessed in (a) recombinant cells (inhibition of U-II-mediated [Ca²⁺]_i-mobilization in HEK293 cells expressing mammalian UT receptors) and (b) native tissue (inhibition of U-II-induced contraction of rat aortae).

SB-706375 as a competitive, mammalian UT receptor antagonist in recombinant HEK293-UT cells UT receptor-mediated Ca²⁺-mobilization studies were carried out in Fluo-3-loaded HEK293 cells stably expressing either the mouse, rat, monkey or human UT receptor using a microtitre plate-based fluorometric imaging plate reader (FLIPR, Molecular Devices, Sunnyvale, CA, U.S.A.) assay (Ames *et al.*, 1999). On the day before the assay, cells expressing recombinant HEK293-UT were plated in 96-well black wall/clear bottom Biocoat plates (Becton Dickinson, San Jose, CA, U.S.A.) at ~50,000 cells well⁻¹ (producing ~80–95% confluence on the day of assay). On the day of the experiment, growth media were aspirated and replaced with 100 µl 'loading media' (Eagle's minimal essential media with Earl's salts, L-glutamine, 0.1% BSA, 2.5 mM probenecid, 4 µM Fluo-3-acetoxymethyl ester fluorescent indicator dye [2 mM stock solution in DMSO with 20% pluronic acid; Fluo-3AM, Molecular Probes, Eugene, OR, U.S.A.]). Cells were incubated for 1 h at 37°C at which point media were aspirated and replaced with identical media lacking Fluo-3AM. Cells were incubated for a further 10 min. Cells were then washed (three times with 'assay buffer'; 120 mM NaCl, 4.6 mM KCl, 1.03 mM

KH₂PO₄, 25 mM NaHCO₃, 1.0 mM CaCl₂, 1.1 mM MgCl₂, 11 mM glucose, 20 mM HEPES [pH 7.4], 0.1% gelatin, 2.5 mM probenecid) at which point they were exposed to SB-706375 (0.1–30 μM) for 10 min prior to hU-II administration (10 pM–3.3 μM for the mouse and monkey UT receptor, 30 pM–10 μM for the rat and human UT receptor). The maximum change in agonist-induced fluorescence was quantified and antagonist pK_b determined by Clark analysis (Lew & Angus, 1997).

Rat isolated aortae studies The antagonistic properties of SB-706375 were assessed in rat isolated aortic contraction assays. All animal procedures were performed in accredited facilities in accordance with institutional guidelines (Animal Care and Use Committee, GlaxoSmithKline) and the Guide for the Care and Use of Laboratory Animals (DHSS #NIH 85-23). Male Sprague–Dawley rats (400 g; Charles River, Raleigh, NC, U.S.A.) were anaesthetized with 5% isoflurane in O₂ and euthanized by exsanguination. Isolated aortae were cleaned of adherent tissue and denuded of endothelium using a pair of fine forceps (functional loss was confirmed using 10 μM carbachol). Vessel rings, approximately 2–3 mm in length, were suspended in 10 ml organ baths containing Krebs of the following composition (mM): NaCl, 112.0; KCl, 4.7; KH₂PO₄, 1.2; MgSO₄, 1.2; CaCl₂, 2.5; NaHCO₃, 25.0; dextrose, 11.0; indomethacin, 0.01 (0.1% ethanol, v v⁻¹). Krebs was maintained at 37 ± 1 °C and aerated with 95% O₂ : 5% CO₂ (pH 7.4). Changes in isometric force were measured under 1.0 g optimal resting tension using MLT0201/D force-displacement transducers (Letica Scientific Instruments, Barcelona, Spain) and recorded digitally using ADInstruments Chart 5.0 software (Colorado Springs, CO, U.S.A.). Following a 60 min equilibration period, the vessels were treated with standard concentrations of KCl (60 mM) and noradrenaline (10 μM). Subsequent agonist-induced responses were normalized to these responses. Each tissue was used to generate only one concentration–response curve.

Characterization of SB-706375-induced inhibition of hU-II contraction in rat isolated aortae (pK_b determination): Rat isolated thoracic aortae were pretreated with either vehicle (0.1% DMSO) or SB-706375 (0.3, 1 or 3 μM) for 30 min following which cumulative concentration–response curves to hU-II (10 pM–10 μM) were constructed (pK_b was determined by Clark analysis; Lew & Angus, 1997).

Selectivity of SB-706375 in rat isolated aortae: Paired thoracic aortae were pretreated with either vehicle (0.1% DMSO) or SB-706375 (1 μM, a concentration > 30-fold greater than its pK_b against hU-II in this tissue) for 30 min following which cumulative concentration–response curves to either KCl (5–90 mM), phenylephrine (0.1 nM–10 μM), angiotensin II (0.1 nM–1 μM) or endothelin-1 (0.1 nM–1 μM) were generated.

Reversibility of SB-706375-induced inhibition in rat isolated aortae: Following a 30-min pretreatment with either vehicle (0.1% DMSO) or 1 μM SB-706375, cumulative concentration–response curves to hU-II (0.1 nM–10 μM) were generated. Additional groups of tissues were washed (Krebs was replaced with fresh buffer, which did not contain any antagonist) repeatedly for either 5, 15, 30 or 60 min following which hU-II concentration–response curves were generated. For comparison, an identical washout study was performed in parallel using phenylephrine (1 nM–100 μM) and the competitive and reversible α₁-adrenoceptor antagonist prazosin (3 nM; as with

SB-706375 above, ~30-fold above pK_b determined in the rat isolated aorta; Aboud *et al.*, 1993).

SB-706375-induced inhibition of hU-II established tone in rat isolated aortae: Paired aortae were contracted with 10 nM hU-II (~EC₈₀ concentration). Once the contractile response had reached a plateau (~25 min), cumulative concentration–response curves to SB-706375 were generated by adding the antagonist to the organ bath in log unit intervals from 1 nM to 1 μM (DMSO vehicle was administered in equal volumes to a parallel set of aortae to serve as a ‘time control’).

Drugs and materials

SB-706375 (2-bromo-4,5-dimethoxy-*N*-[3-(*R*)-1-methyl-pyrrolidin-3-yloxy]-4-trifluoro-methyl-phenyl]-benzenesulphonamide HCl), SB-657510 (2-bromo-*N*-[4-chloro-3-((*R*)-1-methyl-pyrrolidin-3-yloxy)-phenyl]-4,5-dimethoxybenzenesulphonamide HCl) and [³H]SB-657510 (three [³H] incorporated into the pyrrolidine *N*-methyl group; specific activity 87 Ci mmol⁻¹) were synthesized at GlaxoSmithKline (King of Prussia, PA, U.S.A.; Dhanak *et al.*, 2002). The human isoform of U-II was synthesized by California Peptide Research Inc. (Napa, CA, U.S.A.) whereas goby, rat and mouse U-II were from Phoenix Pharmaceuticals Inc. (Mountain View, CA, U.S.A.). Porcine U-II_{A/B} isoforms were synthesized at GlaxoSmithKline. [¹²⁵I]hU-II (Tyr⁹ monoiodinated) was custom-synthesized by Amersham (Arlington Heights, IL, U.S.A.; specific activity 2000 Ci mmol⁻¹). Angiotensin II, carbachol, indomethacin, noradrenaline, phenylephrine and prazosin were from Sigma (St Louis, MO, U.S.A.). All other reagents used were of analytical grade unless otherwise stated. Reagents were made up freshly daily and stored in light-tight containers.

Data analysis

Unless stated to the contrary (pK_b determinations, *vide infra*), all values are expressed as mean ± standard error of the mean (s.e.m.). *n* represents the total number of individual experiments performed and statistical comparisons were made using paired, two-tailed *t*-tests (differences were considered significant when *P* ≤ 0.05) or ANOVA (Dunnett’s multiple comparisons).

Equilibrium binding affinities (*K*_D) and maximum binding site densities (*B*_{max}) were determined by interactive nonlinear curve-fitting using GraphPad Prism (v3.0; GraphPad Software Inc., San Diego, CA, U.S.A.). Unless stated to the contrary (see section ‘Reversibility of binding in HEK293 cells stably expressing the monkey recombinant UT receptor’; *vide infra*), all *K*_D estimates were determined by saturation binding analysis (estimated by exposing HEK293-UT cell membranes to increasing concentrations of radiolabel, that is [¹²⁵I]hU-II or [³H]SB-657510). Hill coefficients (*n*_H) were assumed to approximate unity (see Scatchard plot, Figure 4a; Ames *et al.*, 1999; Elshourbagy *et al.*, 2002; Aiyar *et al.*, 2005):

$$B = \frac{B_{\max}[A]}{K_D + [A]}$$

where *B* represents specific binding, [*A*] the concentration of ligand studied, *B*_{max} the maximum number of binding sites and *K*_D the equilibrium dissociation constant.

*K*_is were calculated by competition binding in recombinant HEK293 cell membranes (see section ‘Competition binding

studies in recombinant HEK293-UT cell membranes' for [¹²⁵I]hU-II binding and 'Inhibition of [³H]SB-657510 binding to recombinant HEK293-UT cell membranes' for [³H]SB-657510 binding) and SJRH30 cells (see section 'Analysis of SB-706375 binding to native UT receptors in human intact SJRH30 cells' for [¹²⁵I]hU-II binding) using the equation by Cheng & Prusoff (1973):

$$K_i = \frac{IC_{50}}{1 + [A]/K_D}$$

where [A] represents the concentration of competing ligand (nonpeptide antagonist or peptide agonist), IC₅₀ the concentration of competing ligand that inhibits radiolabel binding by 50% and K_D the equilibrium dissociation constant of the radioligand.

Agonist contractile response curves generated in rat isolated aortae and Ca²⁺-mobilization studies were analysed using the Hill equation (a logistic equation described previously; Douglas *et al.*, 1995):

$$R = \frac{R_{max}[C]^{n_H}}{EC_{50}^{n_H} + [C]^{n_H}}$$

where *R* is the contractile response, [C] the concentration of agonist, EC₅₀ the concentration of agonist required to produce a half contractile response, n_H the Hill coefficient and R_{max} the maximum contractile response.

Antagonist affinity determinations (pK_b and associated 95% CI) for Ca²⁺-mobilization studies and isolated aortic contraction assays (see sections 'SB-706375 as a competitive, mammalian UT receptor antagonist in recombinant HEK293-UT cells' and 'Selectivity of SB-706375 in rat isolated aortae') were determined by nonlinear regression (Clark) analysis using the method of Lew & Angus (1997):

$$pEC_{50} = -\log([B] + 10^{-pK_b}) - \log c$$

where [B] is the antagonist concentration and the constant log *c* is the difference between the antagonist pK_b and the agonist control curve pEC₅₀.

Finally, in one series of experiments (see section 'Reversibility of binding in HEK293 cells stably expressing the monkey recombinant UT receptor'), K_D was determined by homologous competition binding (whereby multiple concentrations of 'cold' hU-II are used to compete for binding with a fixed concentration of [¹²⁵I]hU-II). Data were analysed by LIGAND (assuming radioligand and competitor both bound reversibly to a single binding site; MacLigand, Version 4.97, NIH, Bethesda, MD, U.S.A.):

$$B = \frac{B_{max}[\text{hot ligand}]}{K_D + [\text{hot ligand}] + [\text{cold ligand}]}$$

where *B* represents specific binding, [hot ligand] the single concentration of [¹²⁵I]hU-II studied, [cold ligand] the concentration of unlabelled hU-II competing with the radiolabel for UT receptor binding, B_{max} the maximum number of binding sites and K_D the equilibrium dissociation constant (the equation was solved where the 'cold ligand' IC₅₀ = [hot ligand] + K_D).

Results

SB-706375 inhibits [¹²⁵I]hU-II binding in mammalian recombinant HEK293-UT cells

[¹²⁵I]hU-II (200 pM) binding to mouse, rat, cat, monkey and human recombinant UT receptors was specific (>90% total binding), saturable (B_{max} ~ 325–1000 fmol mg⁻¹ protein) and of high affinity (K_D 0.2–0.6 nM). Binding was not evident in control ('vector-transfected') HEK293 cell membranes. Goby, human, rat, mouse, pig (A/B isoforms) and human U-II all displaced radioligand from all five UT receptor isoforms with comparable affinities (K_i 0.5–5.7 nM; Table 1).

SB-706375 inhibited [¹²⁵I]hU-II binding to recombinant UT receptors in a 'pan-species' manner with inhibitory potencies (K_i) ranging from 4 to 20 nM across the five mammalian UT isoforms tested (Figure 2 and Table 1). Inhibition was concentration-dependent and the Hill slope (approximating unity) indicated the presence of a homogeneous, single population of binding sites.

Reversibility of SB-706375 binding in mammalian recombinant HEK293-UT cells

Relative to vehicle (DMSO)-treated control cells, prior exposure to 1 μM SB-706375 for 30 min did not alter [¹²⁵I]hU-II binding affinity or density at the monkey recombinant UT receptor (Table 2) consistent with a reversible mode of SB-706375 binding.

High-affinity SB-706375 binding to native human UT receptors expressed in SJRH30 rhabdomyosarcoma cells

In accord with those K_is determined for the human UT receptor using recombinant HEK293 cell membranes,




Figure 2 SB-706375 inhibits [¹²⁵I]hU-II binding to HEK293 membranes stably transfected with mammalian recombinant UT receptors. Membranes isolated from HEK293 cells transfected with the mouse, rat, cat, monkey or human UT receptor were incubated with SB-706375 (0.1 nM–1 μM) and [¹²⁵I]hU-II (0.3 nM) for 60 min at 25°C. Nonlinear regression analysis revealed high-affinity non-cooperative interactions with a single class of binding sites (each point represents the mean of duplicate determinations and is representative of three to six independent experiments; K_is are shown in Table 1).

Table 1 Pharmacological characterization of mammalian recombinant UT receptors with [¹²⁵I]hU-II (saturation binding), SB-706375 and U-II isopeptides (competition binding)

	K_D (pM)					B_{max} (fmol mg ⁻¹)				
	Mouse	Rat	Cat	Monkey	Human	Mouse	Rat	Cat	Monkey	Human
<i>Saturation binding</i>										
[¹²⁵ I]hU-II	654 ± 154 ^a	580 ± 70 ^b	267 ± 25	214 ± 65 ^a	430 ± 10 ^b	1011 ± 125 ^a	325 ± 80 ^b	790 ± 60	497 ± 68 ^a	447 ± 58 ^b
	K_i (nM)					n_H				
	Mouse	Rat	Cat	Monkey	Human	Mouse	Rat	Cat	Monkey	Human
<i>Competition binding</i>										
SB-706375	19.1 ± 1.0	20.7 ± 3.6	4.7 ± 1.5	5.3 ± 1.5	9.3 ± 1.0	0.91 ± 0.12	1.75 ± 0.46	1.02 ± 0.01	0.99 ± 0.04	0.97 ± 0.06
Human U-II	5.7 ± 2.8	2.1 ± 0.2	2.3 ± 0.1	2.5 ± 0.7	2.5 ± 0.3	1.10 ± 0.04	0.94 ± 0.02	1.05 ± 0.06	0.77 ± 0.06	0.84 ± 0.04
Goby U-II	3.2 ± 1.4	1.6 ± 0.2	1.9 ± 0.1	2.0 ± 0.6	2.4 ± 0.3	0.90 ± 0.05	1.02 ± 0.08	0.82 ± 0.04	1.00 ± 0.09	0.79 ± 0.07
Mouse U-II	3.6 ± 1.7	1.6 ± 0.4	3.3 ± 0.2	2.1 ± 0.5	3.2 ± 1.4	1.16 ± 0.04	0.91 ± 0.01	1.11 ± 0.05	0.83 ± 0.05	0.91 ± 0.05
Rat U-II	2.5 ± 0.8	1.4 ± 0.2	2.7 ± 0.6	1.8 ± 0.4	2.9 ± 1.2	0.89 ± 0.05	0.91 ± 0.03	0.79 ± 0.03	0.86 ± 0.03	0.85 ± 0.02
Porcine U-II _A	2.4 ± 1.1	0.8 ± 0.3	5.0 ± 1.8	1.5 ± 0.4	1.6 ± 0.2	1.16 ± 0.17	0.82 ± 0.04	1.41 ± 0.38	0.75 ± 0.07	0.95 ± 0.06
Porcine U-II _B	1.9 ± 1.0	0.5 ± 0.2	1.8 ± 0.6	1.3 ± 0.3	0.9 ± 0.1	1.09 ± 0.07	0.92 ± 0.05	1.19 ± 0.23	0.84 ± 0.09	1.10 ± 0.08

All values, represented as mean ± s.e.m. ($n = 3-6$, duplicate determinations), were determined using membranes isolated from HEK293 cells stably transfected with the mouse, rat, cat, monkey or human UT receptor. U-II isopeptides displayed monophasic competition curves.

^aData are from Elshourbagy *et al.* (2002) and are included for ease of comparison.

^bData are from Ames *et al.* (1999) and are included for ease of comparison.

SB-706375 was a potent inhibitor (K_i 5 nM) of [¹²⁵I]hU-II binding to native human UT receptors expressed by intact SJRH30 cells. Inhibition was monophasic, indicative of an interaction with a homogeneous population of binding sites. Both human and goby U-II also inhibited radioligand binding in this native human cell line in a concentration-dependent manner with K_i 0.5–0.6 nM (Figure 3 and Table 3).

In vitro selectivity profile of SB-706375

SB-706375 was inactive when cross-screened against a diverse range of 86 distinct (see section 'In vitro selectivity profile of SB-706375' mentioned earlier) G-protein-coupled receptors, ion channels, enzymes, transporters and nuclear hormone receptor assays (K_i/IC_{50} were all > 1 μM). As such, SB-706375 functioned as a selective UT receptor antagonist with ≥ 100-fold selectivity for the human UT receptor isoform (K_i 9 nM).

Inhibition of [³H]SB-657510 binding to recombinant HEK293-UT cell membranes

Specific [³H]SB-657510 (10 nM) binding (95 ± 4% total binding) rapidly reached steady state (≤ 10 min at 25°C) in monkey UT-HEK293 membranes and was maintained for ≥ 2 h. The rate of association and dissociation of [³H]SB-657510 binding was fast, reversible and specific for UT-HEK293 membranes. [³H]SB-657510 kinetic rate constants for both association (for a single concentration of 10 nM [³H]SB-657510) and dissociation (made by adding excess cold ligand at equilibrium [30 min]) were estimated to be $t_{1/2}$ 2.0 and 1.2 min, respectively. [³H]SB-657510 binding was saturable, and Scatchard analysis suggested the presence of a single class of high-affinity binding sites ($K_D \sim 2$ nM and $B_{max} \sim 0.9$ pmol mg⁻¹; Figure 4a and Table 4). No specific binding was observed to other GPCRs, and no displacement of [³H]SB-657510 from the UT receptor was evident following exposure to endothelin-1, neuromedin-U, somatostatin or CGRP. [³H]SB-657510 binding was

Table 2 Reversible nature of SB-706375 binding to monkey recombinant UT receptor

	K_D (nM)	B_{max} (pmol mg ⁻¹ protein)
Control (vehicle-treated)	3.1 ± 0.4	3.1 ± 1.0
Pretreated with SB-706375	5.8 ± 0.9	2.8 ± 0.8

Prior exposure to SB-706375 did not alter [¹²⁵I]hU-II radioligand binding site density (B_{max}) or affinity (K_D). Competition binding was performed using membranes pretreated with either vehicle or SB-706375 (1 μM for 30 min at 25°C followed by extensive washing) from HEK293 cells stably transfected with the monkey UT receptor. Data are expressed as mean ± s.e.m. ($n = 4$, determined in duplicate).




Figure 3 Inhibition of [¹²⁵I]hU-II binding to native human UT receptors expressed by SJRH30 cells (200 pM at 37°C for 30 min) by goby and human U-II and SB-706375. Each point represents the mean of duplicate determinations and is representative of results from one of four independent experiments. Binding parameters (K_D , B_{max} and K_i) were determined by nonlinear regression analysis and were consistent with non-cooperative interaction with a single class of binding sites (Table 3).

Table 3 SB-706375 inhibits [¹²⁵I]hU-II binding at the native human UT receptor endogenously expressed in the rhabdomyosarcoma cell line SJRH30

	K_i (nM)	n_H
<i>Competition binding</i>		
SB-706375	5.4 ± 0.4	0.88 ± 0.07
Human U-II	0.5 ± 0.1	0.98 ± 0.16
Goby U-II	0.6 ± 0.2	0.89 ± 0.08

All values are represented as mean \pm s.e.m. ($n=4$, duplicate determinations).




Figure 4 (a) [³H]SB-657510 (0.2–25 nM) saturation binding to monkey UT-HEK293 cell membranes (nonspecific binding defined using 1 μ M hU-II). Scatchard plot analysis (inset) revealed [³H]SB-657510 binding to a single class of receptors. (b) Inhibition of 10 nM [³H]SB-657510 binding (25°C for 30 min) to the monkey recombinant UT receptor by nonpeptidic (SB-706375 and SB-657510) and peptidic (human U-II) ligands. Each figure is representative of $n=3$ independent experiments run in duplicate. Binding parameters (K_D , B_{max} and K_i) were determined by nonlinear regression analysis (Table 4).

inhibited by SB-706375 and ‘cold’ SB-657510 and by human and goby U-II with K_i s ranging from 4 to 17 nM (Figure 4b and Table 4), values consistent with those generated previously using [¹²⁵I]hU-II as the radioligand (Table 1).

Table 4 Inhibition of [³H]SB-657510 binding to monkey UT-HEK293 cell membranes by nonpeptide sulphonamide antagonists (SB-706375 and SB-657510) and native U-II isopeptides

	K_D (nM)	B_{max} (fmol mg ⁻¹ protein)
<i>Saturation binding</i>		
[³ H]SB-657510	2.6 ± 0.4	860 ± 120
K_i (nM) n_H		
<i>Competition binding</i>		
SB-706375	4.6 ± 1.3	0.99 ± 0.05
SB-657510 K_i	17.6 ± 5.4	0.89 ± 0.03
Human U-II K_i	11.4 ± 3.3	1.16 ± 0.15
Goby U-II K_i	14.6 ± 4.3	0.96 ± 0.12

All values are represented as mean \pm s.e.m. ($n=3-4$, duplicate determinations).

Competitive, mammalian UT receptor antagonist in recombinant HEK293-UT cells

As previously reported (Ames *et al.*, 1999; Elshourbagy *et al.*, 2002), hU-II was a potent agonist at mouse, rat, monkey and human UT receptors stably expressed in HEK293 cells ($[Ca^{2+}]_i$ -mobilization EC_{50} s 0.10 ± 0.00 , 1.33 ± 0.41 , 0.13 ± 0.04 and 0.50 ± 0.14 nM, respectively; $n=3$). SB-706375 (0.1–30 μ M) produced a concentration-dependent inhibition of hU-II-induced Ca^{2+} -mobilization in all four recombinant cell lines tested (Figure 5). The resultant rightward, parallel shifts in the hU-II concentration–response curves were consistent with a competitive, surmountable mode of action. In agreement with this, Clark analysis revealed pK_b s for SB-706375 that ranged from 7.29 to 8.00 ($\sim 10-50$ nM; Figure 6 and Table 5). Exposure to SB-706375 did not result in any change in basal $[Ca^{2+}]_i$ levels in HEK293 cells expressing mammalian recombinant UT receptors.

SB-706375 pK_b determination in the rat isolated aorta

Exposure of rat isolated aortic rings to SB-706375 (0.3–3.0 μ M) resulted in concentration-dependent, rightward, parallel shifts in the hU-II concentration–response curves (Figure 7). Inhibition was surmountable inasmuch as the maximal contractile responses (R_{max}) to hU-II remained unaltered (97–122% KCl response; Table 6). Global nonlinear regression (Clark) analysis revealed a pK_b of 7.47 (7.25–7.69 [95% confidence interval]), a value corresponding to an affinity of ~ 33 nM. Pretreatment of rat isolated aortae with SB-706375 did not result in any change in basal tone, that is, SB-706375 was devoid of any intrinsic activity and did not induce a contractile response in this tissue.

Selective inhibition of hU-II-induced contraction in the rat isolated aorta

Exposure to 1 μ M SB-706375, a concentration ~ 30 -fold over the pK_b for hU-II in this vessel, did not alter the contractile responses induced by either (a) membrane depolarization (with KCl) or (b) contractile mechanisms involving ‘non-U-II’ G-protein-coupled receptors, for example, phenylephrine




Figure 5 SB-706375 is a competitive antagonist of hU-II-induced $[Ca^{2+}]_i$ -mobilization (FLIPR) in HEK293 cells stably expressing the (a) mouse, (b) rat, (c) monkey or (d) human UT receptor. Concentration–response curves to hU-II (10 pM–3.3 μ M for mouse and monkey UT receptors, 30 pM–10 μ M for rat and human UT receptors) were rightward-shifted in a parallel manner by SB-706375 (0.1–30 μ M) without any significant suppression of the maximal response to hU-II consistent with competitive antagonism ($n = 3$).

(α_1 -adrenoceptor), angiotensin II (AT_1) and endothelin-1 (ET_A ; Table 7 and Figure 8).

Reversibility of SB-706375-induced inhibition in the rat isolated aorta

The degree of SB-706375-induced inhibition of hU-II-mediated contraction was attenuated by removal of SB-706375 from the organ bath by repeated washing (Figure 9a and Table 8), that is, the inhibitory effects of SB-706375 were reversible. Relative to vehicle-treated tissues, a 30 min preincubation with 1 μ M SB-706375 shifted the hU-II concentration–response curve \sim 30-fold to the right provided SB-706375 was left in contact with the tissues (Table 8). However, repeatedly rinsing the organ baths with normal, ‘antagonist-free’ Krebs solution for 5–60 min following an initial 30 min preincubation period with 1 μ M SB-706375 resulted in time-dependent, leftward shifts of the hU-II concentration–response curves back towards the control (vehicle-treated) hU-II concentration–response curve. Similar data were obtained with phenylephrine/prazosin (Figure 9b and Table 8). The ‘reversibility/washout’ rate constants for SB-706375 and

prazosin were similar, estimated to be 0.022 min^{-1} ($r = 0.998$) and 0.029 min^{-1} ($r = 0.883$), respectively (Figure 9c).

SB-706375-induced inhibition of hU-II established tone in rat isolated aortae

SB-706375 readily attenuated (100% reversal) tone established in the rat aorta by prior exposure to hU-II with $88.6 \pm 10.2 \text{ nM}$ IC_{50} (Figure 10). Application of the Cheng–Prusoff equation (assuming a 3 nM EC_{50} for hU-II) yielded an estimated K_{app} of $\sim 20 \text{ nM}$, consistent with the 7.47 p K_b (33 nM) reported above (see section ‘SB-706375 p K_b determination in the rat isolated aorta’). Established hU-II-induced tone was reduced by 50% within $12.1 \pm 1.7 \text{ min}$ of exposure to 10 nM SB-706375.

Discussion

In recent years, U-II has emerged as a putative cardiorenal target of considerable interest within the pharmaceutical and medical communities (Douglas & Ohlstein, 2000; Maguire & Davenport, 2002; Russell, 2004). This interest has been driven,




Figure 6 SB-706375 is a competitive antagonist of hU-II-induced $[Ca^{2+}]_i$ -mobilization in HEK293 cells stably expressing the (a) mouse, (b) rat, (c) monkey or (d) human UT receptor. Clark plot analysis (global nonlinear regression) revealed pK_b values of 7.29, 7.94, 7.81 and 8.00 (7.20–7.38, 7.51–8.38, 7.64–7.97 and 7.83–8.17, 95% CI). The unity lines shown through the data are included for display purposes only.

Table 5 Potency determination (pK_b) for SB-706375 in intact HEK293 cells stably expressing mammalian recombinant UT receptors using an hU-II-induced $[Ca^{2+}]_i$ -mobilization (FLIPR) assay

UT receptor isoform	pK_b
Mouse	7.29 (7.20–7.38)
Rat	7.94 (7.51–8.38)
Cat ^a	—
Monkey	7.81 (7.64–7.97)
Human	8.00 (7.83–8.17)

pK_b values were determined by Clark analysis (global nonlinear regression; Lew & Angus, 1997) and are represented as mean values with 95% confidence intervals in parentheses ($n = 3$).

^aAntagonism was not determined at the cat UT receptor.

in large part, by the pronounced vasopressor activities exerted by U-II in intact mammals such as the cat (Behm *et al.*, 2004a), monkey (Ames *et al.*, 1999) and man (Böhm & Pernow, 2002; Lim *et al.*, 2004). Further to this, several clinical studies have

noted an emerging association between elevated U-II levels or genotype and cardiorenal diseases such as hypertension, heart failure, atherosclerosis, renal failure and diabetes (see Gilbert *et al.*, 2004; Richards & Charles, 2004). Nevertheless, the precise (patho)physiological significance of U-II remains ambiguous given the fact that the haemodynamic actions of U-II vary significantly between species (in contrast to the cat and primate, bolus i.v. U-II lacks any haemodynamic activity in sheep and induces a ‘paradoxical’ vasodilation in the rat; Hasegawa *et al.*, 1992; Watson *et al.*, 2003; Gardiner *et al.*, 2004).

Although plethysmographic (Böhm & Pernow, 2002), iontophoretic (Lim *et al.*, 2004) and direct intradermal injection (Leslie *et al.*, 2000) studies demonstrate that local U-II administration elevates regional vascular resistance in healthy humans and heart failure patients, at least one well-respected group has failed to demonstrate any such response using almost identical techniques (Affolter *et al.*, 2002; Wilkinson *et al.*, 2002). Further, and in contrast to several other reports (MacLean *et al.*, 2000; Maguire *et al.*, 2000; 2004;




Figure 7 SB-706375 is a competitive inhibitor of hU-II-induced contraction in the rat isolated aorta. (a) SB-706375 (0.3–3.0 μM) produces concentration-dependent rightward, parallel shifts in the hU-II concentration–response curve with no suppression of the maximal response to hU-II consistent with a competitive mode of antagonism ($n=12$). (b) Clark plot (global nonlinear regression analysis) revealed 7.47 pK_b (7.25–7.69 95% CI). The unity line shown through the data is included for display purposes only.

Table 6 SB-706375 is a competitive antagonist of hU-II-induced contraction of the rat isolated aorta

Treatment	EC_{50} (nM)	R_{max} (% 60 mM KCl)	n_H	Dose ratio
Vehicle	2.3 ± 0.4	97 ± 10	1.90 ± 0.15	—
0.3 μM SB-706375	24.0 ± 7.7	122 ± 9	1.93 ± 0.20	13.1 ± 3.5
1.0 μM SB-706375	79.3 ± 15.1	122 ± 7	1.97 ± 0.19	47.6 ± 9.2
3.0 μM SB-706375	182.6 ± 43.4	114 ± 8	1.68 ± 0.18	166.2 ± 66.3

SB-706375 (0.3–3.0 μM) caused concentration-dependent, parallel rightward shifts in the hU-II concentration–response curve. Values are expressed as mean \pm s.e.m. ($n=12$). Neither the maximum contractility (R_{max}) nor Hill slope (n_H) was altered by prior exposure to SB-706375 (parallel rightward shifts in the U-II concentration–response curve) as assessed by ANOVA (analysis for repeated measures; Dunnett's multiple comparisons post-test) consistent with surmountable, competitive mode of action in the rat isolated aorta.

Russell *et al.*, 2001; Paysant *et al.*, 2001; Camarda *et al.*, 2002b), Hillier *et al.* (2001) failed to observe any significant contractile activity with U-II in human isolated arteries in a series of carefully controlled *in vitro* experiments. Clearly, the need for additional studies to help address these inconsisten-

Table 7 SB-706375 (1 μM) is a selective hU-II antagonist in the rat isolated aorta

Treatment	EC_{50} (nM)	R_{max} (% 60 mM KCl)	n_H
<i>Phenylephrine</i>			
Vehicle-treated	19.8 ± 3.1	146 ± 6	2.19 ± 0.41
SB-706375-treated	23.6 ± 2.7	138 ± 7	2.30 ± 0.35
<i>Angiotensin II</i>			
Vehicle-treated	6.3 ± 1.5	38 ± 6	2.52 ± 1.17
SB-706375-treated	8.9 ± 1.9	40 ± 12	1.85 ± 0.53
<i>Endothelin-1</i>			
Vehicle-treated	8.5 ± 1.1	153 ± 6	2.28 ± 0.35
SB-706375-treated	8.6 ± 1.3	151 ± 7	1.84 ± 0.10
<i>KCl</i> ^a			
Vehicle-treated	15.4 ± 0.9	101 ± 7	8.27 ± 1.73
SB-706375-treated	16.9 ± 0.9	95 ± 6	10.51 ± 1.09

Pretreatment of rat isolated aortae with SB-706375 (1 μM) does not affect contractions invoked in the rat isolated aorta by KCl, endothelin-1, angiotensin II or phenylephrine. Values are expressed as mean \pm s.e.m. ($n=4$).

^aContractile responses to KCl are expressed as mM (EC_{50}) and % response to 10 μM noradrenaline (R_{max}). SB-706375 treatment did not alter EC_{50} , R_{max} or n_H values of any spasmogen studied (paired *t*-test).

cies is evident. While observations of SNP associations and elevated plasma U-II levels might be suggestive of a causative role for U-II in the pathogenesis of cardiorenal diseases, such a proposition lacks the affirmation afforded by specific UT receptor antagonists. However, the dearth of such inhibitors has, until now, been prohibitive. It is hoped that the findings of the present study, which details the pharmacodynamic characterization of the novel UT receptor antagonist SB-706375 (see Table 9), will mitigate this issue and facilitate the delineation of the (patho)physiological actions of U-II.

The existing UT receptor ligands described in the literature (Dhanak *et al.*, 2003; Douglas *et al.*, 2004a) are of limited utility as 'tool antagonists' due to a combination of pharmacodynamic and pharmacokinetic considerations. Firstly, many exhibit poor affinities for UT receptor isoforms from non-human, preclinical species for example, SB-710411 lacks potency at the rat UT receptor (Behm *et al.*, 2002). Indeed, the affinities of the majority of such putative UT receptor antagonists are rarely (if ever) reported at non-rat/non-human UT receptor isoforms (i.e. cat and mouse UT receptors). This is an important consideration since, in addition to the rat, the mouse (Vergura *et al.*, 2004) and cat (Behm *et al.*, 2004a) represent two important species for studying the pharmacodynamic actions of U-II. Further, with the exception of the quinolinylurea palosuran (ACT-058362; Clozel *et al.*, 2004), all UT receptor ligands published to date are peptidic and, as such, are inherently unsuitable for chronic enteric administration. In addition, several rat UT receptor antagonists (e.g. lanreotide, SB-710411, [Orn⁸]hU-II and urantide) possess intrinsic activity in mouse (partial agonism) and primate (full agonism) UT receptor systems, thus further complicating the interpretation of the pharmacological actions and hence restricting their use (Camarda *et al.*, 2002a; 2004; Herold *et al.*, 2002; Behm *et al.*, 2004b). Finally, relatively little




Figure 8 SB-706375 (1 μM) was a selective hU-II antagonist in the rat aorta since 1 μM concentration of the ligand (a concentration some 30-fold above its pK_b against hU-II in this tissue) failed to inhibit the contractile actions of (a) KCl, (b) phenylephrine, (c) angiotensin II or (d) endothelin-1 ($n=4$) in the same tissue.

is known about the selectivity profiles of such putative antagonists at present (significant 'secondary interactions' have been established for SB-710411, BIM-23127 and lanreotide/octreotide; Behm *et al.*, 2002; Herold *et al.*, 2002; 2003). As will be detailed below, the limitations described above do not apply to SB-706375, a potent and selective UT receptor antagonist active across mammalian species. As such, the discovery of SB-706375 represents a significant advance in the U-II field.

SB-706375 possesses high affinity for all five of the mammalian UT receptors that have been cloned, including the recently identified cat receptor (Aiyar *et al.*, 2005). In addition to being a potent ligand at recombinant rodent, feline and primate UT receptors (4–20 nM K_i), SB-706375 also inhibits [125 I]hU-II binding with high affinity at endogenous human UT receptors expressed in native human SJRH30 cells (5 nM K_i). As such, SB-706375 is approximately an order of magnitude more potent than the other known nonpeptide U-II antagonist at endogenous human UT receptors (46 nM IC_{50} in human rhabdomyosarcoma TE671 cells; Clozel *et al.*, 2004). Therefore, SB-706375 is suitable for studying the actions of U-II at both recombinant and native UT receptors across species.

In accord with the radioligand binding studies, SB-706375 acted as a surmountable antagonist of hU-II across all species tested, antagonizing hU-II-induced [Ca^{2+}] $_i$ -mobilization in

intact HEK293 cells with pK_b s ranging from around 10 to 50 nM (Hill slopes approximating unity). Such activity is also evident in native tissues (competitive, surmountable inhibition in rat aortae with pK_b 33 nM). Thus, it is concluded that SB-706375 constitutes an agent suitable for attenuating the actions of U-II in cells and tissues from a diverse range of mammalian species.

The 'pan-species' antagonistic properties of SB-706375, coupled with the associated low nanomolar affinity for rodent, cat and primate UT receptors, clearly differentiate this arylsulphonamide from other putative UT receptor ligands. To date, no other antagonists (e.g. BIM-23127, [Orn 8]hU-II, urantide, etc.) have been profiled against monkey or cat UT receptors. Indeed, of this list, only [Orn 8]hU-II has been studied in the mouse (Vergura *et al.*, 2004). The potency of SB-706375 is also striking. Relative to SB-706375 (9 nM rat UT receptor K_i), a compound such as palosuran lacks appreciable affinity at the rat recombinant UT receptor (IC_{50} cited as 1500 and >10,000 nM at the rat UT receptor depending on whether CHO cell membranes or intact cells were used; Clozel *et al.*, 2004). Based on these values, SB-706375 is in excess of 100- to >1000-fold more potent than palosuran as a rat UT receptor ligand. While SB-706375 is only four- to seven-fold more potent at the rat UT receptor than the peptides SB-710411 and




Figure 9 The inhibitory properties of (a) SB-706375 (hU-II-induced contraction) and (b) prazosin (phenylephrine-induced contraction) are reversible by 'washout' in the rat isolated aorta. Following a 30 min preincubation with antagonist (1 μM SB-706375 or 3 nM prazosin) or vehicle, cumulative concentration-response curves to agonists (hU-II or phenylephrine) were generated. Repeatedly washing tissues to remove antagonist from the organ bath for 5, 15, 30 or 60 min prior to initiating the hU-II or phenylephrine concentration-response curves attenuated the degree of antagonism (i.e. rightward shift) observed ($n = 5$). (c) The rate at which SB-706375 and prazosin were washed out of the isolated aorta (0.022 and 0.029 min^{-1} , respectively) was estimated by plotting the degree of antagonism observed ('dose ratio', based on agonist potency in vehicle-treated tissues; see Table 8) against the duration of repeated washing (5–60 min of 'washout').

BIM-23127 (32 and 63 nM rat UT receptor K_i , respectively; Herold *et al.*, 2003; Behm *et al.*, 2004b), it is distinguished from these somatostatin/neuromedin B analogues by superior pharmacodynamic selectivity (*vide infra*).

Although SB-706375 and urantide (5 nM rat UT receptor K_i ; Patacchini *et al.*, 2003) are essentially equipotent human UT receptor ligands, only the former functions as an antagonist at

the human UT receptor. A recent study has revealed that urantide is actually a potent partial agonist at the human UT receptor (Camarda *et al.*, 2004) with pEC_{50} 8.11 and an intrinsic activity $[\alpha] \sim 0.8$ relative to hU-II for $[Ca^{2+}]_i$ -mobilization (consistent with these findings, urantide is also a potent and efficacious cat and monkey artery spasmogen with $EC_{50} \sim 10$ nM and intrinsic activities $[\alpha] \sim 0.5-0.8$; D. Behm & S. Douglas, unpublished observation). Similar observations have also been made with other peptidic moieties such as SB-710411, lanreotide, BIM-23127 and $[Orn^8]hU-II$, which all behave as partial agonists at the mouse UT receptor and/or full agonists at monkey and human UT receptors (Camarda *et al.*, 2002a; 2004; Herold *et al.*, 2002). SB-710411, for example, is an efficacious antagonist at the rat UT receptor, yet it promotes phosphoinositide hydrolysis in monkey UT-HEK293 cells and induces contraction of the monkey isolated carotid artery (Behm *et al.*, 2004b). Such liabilities are not, however, a characteristic shared by SB-706375, which does not exhibit any intrinsic activity at any of the mammalian recombinant UT receptor isoforms tested or in isolated blood vessels.

Ultimately, it will be necessary to study the pharmacodynamic effects of UT receptor antagonists in appropriate cell, tissue and whole animal systems before the true (patho)physiological significance of U-II in mammals becomes apparent. In order to be able to interpret data generated under such conditions unambiguously, a compound must be selective for the UT receptor and without any appreciable 'off target' effects. Regrettably, relatively little is known about the selectivity and specificity of existing putative UT receptor antagonists. Indeed, the limited selectivity information that is available suggests that caution should be used when interpreting the observed biological consequences of antagonist administration. For example, BIM-23127 is a relatively potent neuromedin B ligand (Herold *et al.*, 2002; 2003), whereas lanreotide, octreotide and SB-710411 all exhibit direct/indirect interactions with somatostatin and endothelin-1 receptors at reasonable concentrations (Behm *et al.*, 2002; Herold *et al.*, 2002). Although a handful of antagonists (e.g. $[Pen^5, Orn^8]hU-II[4-11]$, urantide and palosuran) have been shown to have no effect on contractile and relaxant responses elicited in the rat isolated aorta by noradrenaline, 5-HT, endothelin-1 and acetylcholine (at 10 μM, <2-fold greater than the pD_2' of palosuran for U-II in the rat aorta), such studies are not in themselves rigorous tests of pharmacodynamic selectivity. As such, knowledge of the specificity of such agents is currently incomplete. Consequently, it becomes difficult to interpret data generated *in vivo* with such agents with any degree of certainty. However, compare the selectivity window defined for palosuran with that established for SB-706375, for example. Based on the ratio of an antagonist's potency in the rat isolated aorta against U-II (6 μM and 33 nM, respectively) compared to the highest concentration of antagonist studied (1 or 10 μM) that is known not to inhibit 'non-U-II' vasoactive factors (e.g. endothelin-1) in this same tissue, palosuran's selectivity is defined as ≥ 1.6 -fold in rat isolated aorta (Clozel *et al.*, 2004) vs ≥ 30 -fold for SB-706375 in the same tissue. Not only does SB-706375 have a larger selectivity window, this parameter has been defined against a much larger list of diverse molecular targets ($>1 \mu\text{M}$ IC_{50} defined against a total of 86 distinct G-protein-coupled receptors, hormone receptors, tyrosine kinase receptors, ion

Table 8 Inhibitory actions of 1 μM SB-706375 (hU-II-induced contraction) and 3 nM prazosin (phenylephrine-induced contraction) are reversible by repeated washing (5–60 min) of the rat isolated aorta

Treatment	Washout time (min)	hU-II EC_{50} (nM)	hU-II R_{max} (% 60 mM KCl)	Dose ratio	n_H
Vehicle	—	6 \pm 2	113 \pm 15	—	1.47 \pm 0.21
SB-706375	0	148 \pm 21	90 \pm 12	32.3 \pm 6.6	1.68 \pm 0.08
SB-706375	5	40 \pm 11	109 \pm 15	10.9 \pm 4.5	1.69 \pm 0.12
SB-706375	15	38 \pm 9	122 \pm 11	9.2 \pm 2.9	1.36 \pm 0.13
SB-706375	30	29 \pm 8	116 \pm 12	6.4 \pm 1.9	1.29 \pm 0.10
SB-706375	60	14 \pm 3	115 \pm 14	3.3 \pm 1.0	1.66 \pm 0.24

Treatment	Washout time (min)	Phenylephrine EC_{50} (nM)	Phenylephrine R_{max} (% 60 mM KCl)	Dose ratio	n_H
Vehicle	—	13 \pm 2	146 \pm 8	—	2.03 \pm 0.18
Prazosin	0	868 \pm 319	120 \pm 9	70.1 \pm 26.6	1.48 \pm 0.24
Prazosin	5	243 \pm 52	140 \pm 3	22.8 \pm 6.4	1.68 \pm 0.20
Prazosin	15	257 \pm 74	130 \pm 8	25.1 \pm 9.9	2.08 \pm 0.36
Prazosin	30	82 \pm 13	149 \pm 5	7.3 \pm 1.0	1.43 \pm 0.12
Prazosin	60	60 \pm 10	137 \pm 5	5.5 \pm 1.0	2.14 \pm 0.22

Following a 30 min preincubation period with either vehicle or SB-706375 (1 μM), vessels were exposed to increasing concentrations of hU-II. Removal of SB-706375 from the organ bath by repeated washing of the vessels for 5–60 min significantly attenuated the parallel rightward shift seen in the hU-II concentration–response curve, indicating that the inhibitory effects of SB-706375 were reversible. Data were similar to those obtained with an unrelated, pharmacologically distinct antagonist, 3 nM prazosin (phenylephrine [α_1 -adrenoceptor]-induced contraction). All values are expressed as mean \pm s.e.m. ($n = 5$). Neither the maximum contractility (R_{max}) nor Hill slope (n_H) determined for U-II and phenylephrine was altered by prior exposure to either SB-706375 or prazosin (parallel rightward shifts in the agonist concentration–response curves) as assessed by ANOVA (analysis for repeated measures) consistent with surmountable, competitive modes of action for both antagonists in the rat isolated aorta.




Figure 10 SB-706375 reverses hU-II-induced tone established in the rat aorta. Tissues were contracted with 10 nM hU-II and, once induced tone had reached a plateau, aortae were exposed to either vehicle or SB-706375 added to the organ bath in a cumulative manner. SB-706375 reversed the established tone to baseline (100% reversal; $n = 4$).

channels, enzymes and neurotransmitter re-uptake systems). While not exhaustive, this level of selectivity profiling for SB-706375 is extensive relative to the lack of comparable data generated with the other available UT receptor antagonists. This should enable SB-706375 to be used *in vivo* in a more meaningful and less ambiguous manner than existing tool compounds.

The ambiguity that poor or incompletely defined selectivity profiling creates can be seen with studies describing the purported renoprotective effects of systemic palosuran administration (10 mg kg⁻¹ h⁻¹ i.v. infusion) in a rat renal artery ligation–reperfusion model. In this study, Clozel *et al.* (2004) failed to demonstrate equivocally that this palosuran dosage regimen blocks the pharmacodynamic actions of U-II (i.e. it is unknown if this dosage regimen of palosuran inhibits the haemodynamic response evoked by an exogenous U-II challenge). Indeed, the dose used *in vivo* was selected on the basis of palosuran's ability to augment the reactive hyperemia observed upon release of the renal artery clamp, an effect that might well have little to do with UT receptor occupancy. As such, it is difficult to associate the renoprotective effects reported with UT receptor blockade in an equivocal manner. This is of particular concern since plasma palosuran levels attained during this study ($\sim 5 \mu\text{M}$) fell below the affinity of this compound for the rat UT receptor expressed in either the rat isolated aorta or in intact rat recombinant UT-CHO cells (pD_2' 6 μM and $IC_{50} > 10 \mu\text{M}$, respectively; Clozel *et al.*, 2004). Although such considerations might be less of an issue when palosuran is studied in man (due to increased potency at the human UT receptor), it is an issue when using this compound in the rat.

SB-706375 functioned as a reversible UT receptor antagonist. The inhibition observed in the rat isolated aorta could be readily 'washed out', an action consistent with a competitive mode of antagonism. The reversibility seen with SB-706375 was similar to that seen with the α_1 -adrenoceptor antagonist prazosin in this preparation where washout rate constants were in the order of ~ 2 –3% min⁻¹ for both agents. In agreement with this, the reversibility of UT receptor/SB-706375 binding was also demonstrated using monkey UT-HEK293 membranes. Exposure to 1 μM SB-706375 for 30 min immediately followed by washout did not alter [¹²⁵I]U-II binding affinity or density.

Table 9 Synopsis of affinity estimates for SB-706375 at mammalian (mouse, rat, cat, monkey and human) UT receptors as determined in radioligand binding studies (native and recombinant UT receptor assays) and in functional assays using both intact cells (recombinant UT receptor) and isolated arteries (native UT receptor)

	Mouse	Rat	Cat	Monkey	Human
<i>Radioligand binding K_i (nM) estimates (competition binding)</i>					
HEK293-UT membranes ($[^{125}\text{I}]\text{hU-II}$)	19.1	20.7	4.7	5.3	9.3
HEK293-UT membranes ($[^3\text{H}]\text{SB-657510}$)	—	—	—	4.6	—
Intact human SJRH30 cells ($[^{125}\text{I}]\text{hU-II}$)	—	—	—	—	5.4
<i>Functional pK_b estimates</i>					
Intact HEK293-UT cells (inhibition of $[\text{Ca}^{2+}]_i$ -mobilization)	7.29	7.94	—	7.81	8.00
Isolated aortic rings (inhibition of contraction)	—	7.47	—	—	—

hU-II is classified as a 'pseudo-irreversible' ligand at the UT receptor, that is to say that it has an extremely slow dissociation rate from native UT receptors (<15% at 90 min from native hUT receptors expressed in SJRH30 cells; Douglas *et al.*, 2004b). Similar radioligand binding data have been recorded at the recombinant UT receptor (Onan, 2004). Consistent with this observation, the contractile actions of U-II are extremely 'resistant to washout' in mammalian isolated arteries (the time required for a contractile response to return to baseline values approaches ~1 h in the rat and ≥ 8 h in the monkey; Douglas *et al.*, 2000). Despite this profile of U-II, SB-706375 was readily able to reverse 'established hU-II-induced tone' in the rat aorta with $K_{app} \sim 20$ nM. This observation is critical as it indicates that it is not necessary to use SB-706375 in a 'prophylactic' manner in order to antagonize the actions of hU-II in an *in vivo* model. To date, such an action has not been confirmed for other UT receptor antagonists (with the exception of BIM-23127; Herold *et al.*, 2003).

The present report also describes for the first time the characterization of a nonpeptidic UT receptor radiolabel, namely the tritiated radiotracer $[^3\text{H}]\text{SB-657510}$. The development of a high-affinity nonpeptide radioligand such as $[^3\text{H}]\text{SB-657510}$ offers several distinct advantages over currently available UT receptor radiotracers, that is, iodinated fish, amphibian and mammalian U-II isoforms. Antagonists such as $[^3\text{H}]\text{SB-657510}$ are generally preferred as radiolabels over agonists such as $[^{125}\text{I}]\text{hU-II}$ for the characterization of receptors for several reasons. Firstly, antagonist binding does not result in receptor activation and subsequent desensitization/downregulation (SB-657510 is devoid of any intrinsic activity at the UT receptor [pK_b , 7.25 in rat isolated aorta for

'cold' SB-657510; S. Douglas & D. Behm, unpublished data]). Further, antagonist:receptor binding is insensitive to guanine nucleotides and, as such, radiolabels are able to recognize multiple receptor affinity states (note that, to date, the regulation of U-II agonist binding to the UT receptor by guanine nucleotides has not been demonstrated) and, by virtue of its nonpeptidic nature, $[^3\text{H}]\text{SB-657510}$ is refractory to protease degradation. The present study demonstrated that $[^3\text{H}]\text{SB-657510}$ binding was specific (95% of total) and of high affinity ($K_D \sim 2$ nM). The rapid dissociation of $[^3\text{H}]\text{SB-657510}$ binding from the UT receptor (upon addition of cold hU-II) was consistent with a reversible mode of binding. SB-706375 and human and goby hU-II interacted with a common binding site labelled by $[^3\text{H}]\text{SB-657510}$. Consequently, $[^3\text{H}]\text{SB-657510}$ can be considered a useful alternative to $[^{125}\text{I}]\text{hU-II}$ for the pharmacological characterization of the U-II receptor.

In summary, the present report describes the identification and characterization of the arylsulphonamide SB-706375 as a potent, surmountable, reversible and selective mammalian UT receptor antagonist (Table 9). This compound offers several distinct advantages over currently available peptidic and nonpeptidic UT receptor antagonists. Since SB-706375 exhibits high-affinity cross-species antagonism, this nonpeptidic, small molecule inhibitor will serve as a useful agent in delineating the (patho)physiological actions of U-II in both preclinical species and man.

We thank L. Gordon, V. Holland, A. Garner, L. Campbell and K. Winborn (Assay Development and Compound Profiling, Glaxo-SmithKline, Harlow, U.K.) for their expert technical assistance in assessing the selectivity of SB-706375 in a series of radioligand binding studies described herein.

References

- ABOUD, R., SHAFII, M. & DOCHERTY, J.R. (1993). Investigation of the subtypes of alpha 1-adrenoceptor mediating contractions of rat aorta, vas deferens and spleen. *Br. J. Pharmacol.*, **109**, 80–87.
- AFFOLTER, J.T., NEWBY, D.E., WILKINSON, I.B., WINTER, M.J., BALMENT, R.J. & WEBB, D.J. (2002). No effect on central or peripheral blood pressure of systemic urotensin II infusion in humans. *Br. J. Clin. Pharmacol.*, **54**, 617–621.
- AIYAR, N.V., JOHNS, D.G., AO, Z., DISA, J., BEHM, D.J., FOLEY, J.J., BUCKLEY, P.T., SARAU, H.M., VAN DER KEYL, H.K., ELSHOUBAGY, N.A. & DOUGLAS, S.A. (2005). Cloning and pharmacological characterization of the cat urotensin-II receptor (UT). *Biochem. Pharmacol.*, **69**, 1069–1079.
- AMES, R.S., SARAU, H.M., CHAMBERS, J.K., WILLETTE, R.N., AIYAR, N.V., ROMANIC, A.M., LOUDEN, C.S., FOLEY, J.J., SAUERMEHL, C.F., COATNEY, R.W., AO, Z., DISA, J., HOLMES, S.D., STADEL, J.M., MARTIN, J.D., LIU, W.S., GLOVER, G.I., WILSON, S., McNULTY, D.E., ELLIS, C.E., ELSHOUBAGY, N.A., SHABON, U., TRILL, J.J., HAY, D.W., OHLSTEIN, E.H., BERGSMAN, D.J. & DOUGLAS, S.A. (1999). Human urotensin-II is a potent vasoconstrictor and agonist for the orphan receptor GPR14. *Nature*, **401**, 282–286.
- BEHM, D.G., HEROLD, C.L., OHLSTEIN, E.H., KNIGHT, S.D., DHANAK, D. & DOUGLAS, S.A. (2002). Pharmacological characterization of SB-710411 (Cap-c[D-Cys-Pal-D-Trp-Lys-Val-Cys]-Cpa-amide), a novel peptidic urotensin-II receptor antagonist. *Br. J. Pharmacol.*, **137**, 449–458.

- BEHM, D.J., DOE, C.P., JOHNS, D.G., MANISCALCO, K., STANKUS, G.P., WIBBERLEY, A., WILLETTE, R.N. & DOUGLAS, S.A. (2004a). Urotensin-II: a novel systemic hypertensive factor in the cat. *Naunyn Schmiedeberg's Arch. Pharmacol.*, **369**, 274–280.
- BEHM, D.J., HEROLD, C.L., CAMARDA, V., AIYAR, N.V. & DOUGLAS, S.A. (2004b). Differential agonistic and antagonistic effects of the urotensin-II ligand SB-710411 at rodent and primate UT receptors. *Eur. J. Pharmacol.*, **25**, 113–116.
- BÖHM, F. & PERNOW, J. (2002). Urotensin II evokes potent vasoconstriction in humans *in vivo*. *Br. J. Pharmacol.*, **135**, 25–27.
- BOUSETTE, N., PATEL, L., DOUGLAS, S.A., OHLSTEIN, E.H. & GIAID, A. (2004). Increased expression of urotensin II and its cognate receptor GPR14 in atherosclerotic lesions of the human aorta. *Atherosclerosis*, **176**, 117–123.
- CAMARDA, V., GUERRINI, R., KOSTENIS, E., RIZZI, A., CALO', G., HATTENBURGER, A., ZUCCHINI, M., SALVADORI, S. & REGOLI, D. (2002a). A new ligand for the urotensin II receptor. *Br. J. Pharmacol.*, **137**, 311–314.
- CAMARDA, V., RIZZI, A., CALO', G., GENDRON, G., PERRON, S.I., KOSTENIS, E., ZAMBONI, P., MASCOLI, F. & REGOLI, D. (2002b). Effects of human urotensin II in isolated vessels of various species: comparison with other vasoactive agents. *Naunyn Schmiedeberg's Arch. Pharmacol.*, **365**, 141–149.
- CAMARDA, V., SONG, W., MARZOLA, E., SPAGNOL, M., GUERRINI, R., SALVADORI, S., REGOLI, D., THOMPSON, J.P., ROWBOTHAM, D.J., BEHM, D.J., DOUGLAS, S., CALO', G. & LAMBERT, D.G. (2004). Urotensin II mimics urotensin-II induced calcium release in cells expressing recombinant UT receptors. *Eur. J. Pharmacol.*, **498**, 83–86.
- CHEUNG, B.M., LEUNG, R., MAN, Y.B. & WONG, L.Y. (2004). Plasma concentration of urotensin II is raised in hypertension. *J. Hypertens.*, **22**, 1341–1344.
- CHENG, Y.C. & PRUSOFF, W.H. (1973). Relationship between the inhibition constant (K_i) and the concentration of inhibitor which causes 50 percent inhibition (IC_{50}) of an enzymatic reaction. *Biochem. Pharmacol.*, **22**, 357–366.
- CLOZEL, M., BINKERT, C., BIRKER-ROBACZEWSKA, M., BOUKHADRA, C., DING, S.S., FISCHLI, W., HESS, P., MATHYS, B., MORRISON, K., MULLER, C., MULLER, C., NAYLER, O., QIU, C., REY, M., SCHERZ, M.W., VELKER, J., WELLER, T., XI, J.F. & ZILTENER, P. (2004). Pharmacology of the urotensin-II receptor antagonist palosuran (ACT-058362: 1-[2-(4-benzyl-4-hydroxy-piperidin-1-yl)-ethyl]-3-(2-methyl-quinolin-4-yl)-urea sulfate salt): first demonstration of a pathophysiological role of the urotensin system. *J. Pharmacol. Exp. Ther.*, **311**, 204–212.
- DHANAK, D., GALLAGHER, T.F. & KNIGHT, S.D. (2002). Preparation of sulfonamides as antagonists of urotensin II. *PCT Int. Appl.*, WO 2002089792.
- DHANAK, D., NEEB, M.J. & DOUGLAS, S.A. (2003). Urotensin-II receptor modulators. *Ann. Rep. Med. Chem.*, **38**, 99–110.
- DOUGLAS, S.A. (2003). Human urotensin-II as a novel cardiovascular target: 'heart' of the matter or simply a fishy 'tail'? *Curr. Opin. Pharmacol.*, **3**, 159–167.
- DOUGLAS, S.A., BECK, G.R., ELLIOTT, J.D. & OHLSTEIN, E.H. (1995). Pharmacological evidence for the presence of three distinct functional endothelin receptor subtypes in the rabbit lateral saphenous vein. *Br. J. Pharmacol.*, **114**, 1529–1540.
- DOUGLAS, S.A., DHANAK, D. & JOHNS, D.G. (2004a). From 'gills to pills': urotensin-II as a regulator of mammalian cardiorenal function. *Trends Pharmacol. Sci.*, **25**, 76–85.
- DOUGLAS, S.A., NASELSKY, D., AO, Z., DISA, J., HEROLD, C.L., LYNCH, F. & AIYAR, N.V. (2004b). Identification and pharmacological characterization of native, functional human urotensin-II receptors in rhabdomyosarcoma cell lines. *Br. J. Pharmacol.*, **142**, 921–932.
- DOUGLAS, S.A. & OHLSTEIN, E.H. (2000). Human urotensin-II, the most potent mammalian vasoconstrictor identified to date, as a therapeutic target for the management of cardiovascular disease. *Trends Cardiovasc. Med.*, **10**, 229–237.
- DOUGLAS, S.A., SULPIZIO, A.C., PIERCY, V., SARAU, H.M., AMES, R.S., AIYAR, N.V., OHLSTEIN, E.H. & WILLETTE, R.N. (2000). Differential vasoconstrictor activity of human urotensin-II in vascular tissue isolated from the rat, mouse, dog, pig, marmoset and cynomolgus monkey. *Br. J. Pharmacol.*, **131**, 1262–1274.
- DOUGLAS, S.A., TAYARA, L., OHLSTEIN, E.H., HALAWA, N. & GIAID, A. (2002). Congestive heart failure and expression of myocardial urotensin II. *Lancet*, **359**, 1990–1997.
- ELSHOURBAGY, N.A., DOUGLAS, S.A., SHABON, U., HARRISON, S., DUDDY, G., SECHLER, J.L., AO, Z., MALEEFF, B.E., NASELSKY, D., DISA, J. & AIYAR, N.V. (2002). Molecular and pharmacological characterization of genes encoding urotensin-II peptides and their cognate G-protein-coupled receptors from the mouse and monkey. *Br. J. Pharmacol.*, **136**, 9–22.
- GARDINER, S.M., MARCH, J.E., KEMP, P.A. & BENNETT, T. (2004). Bolus injection of human UII in conscious rats evokes a biphasic haemodynamic response. *Br. J. Pharmacol.*, **143**, 422–430.
- GILBERT, R.E., DOUGLAS, S.A. & KRUM, H. (2004). Urotensin-II as a novel therapeutic target in the clinical management of cardiorenal disease. *Curr. Opin. Invest. Drugs*, **5**, 276–282.
- HASEGAWA, K., KOBAYASHI, Y. & KOBAYASHI, H. (1992). Vasodepressor effects of urotensin II in rats. *Neuroendocrinol. Lett.*, **14**, 357–363.
- HEROLD, C.L., BEHM, D.J., BUCKLEY, P.T., FOLEY, J.J. & DOUGLAS, S.A. (2002). The peptidic somatostatin analogs lanreotide, BIM-23127 and BIM-23042 are urotensin-II receptor ligands. *Pharmacologist*, **44**, 170–171.
- HEROLD, C.L., BEHM, D.J., BUCKLEY, P.T., FOLEY, J.J., WIXTED, W.E., SARAU, H.M. & DOUGLAS, S.A. (2003). The neuromedin B receptor antagonist, BIM-23127, is a potent antagonist at human and rat urotensin-II receptors. *Br. J. Pharmacol.*, **139**, 203–207.
- HILLIER, C., BERRY, C., PETRIE, M.C., O'DWYER, P.J., HAMILTON, C., BROWN, A. & McMURRAY, J. (2001). Effects of urotensin II in human arteries and veins of varying caliber. *Circulation*, **103**, 1378–1381.
- JOHNS, D.G., AO, Z., NASELSKY, D., HEROLD, C.L., MANISCALCO, K., SAROV-BLAT, L., STEPLEWSKI, K., AIYAR, N. & DOUGLAS, S.A. (2004). Urotensin-II-mediated cardiomyocyte hypertrophy: effect of receptor antagonism and role of inflammatory mediators. *Naunyn Schmiedeberg's Arch. Pharmacol.*, **370**, 238–250.
- KOMPA, A.R., THOMAS, W.G., SEE, F., TZANIDIS, A., HANNAN, R.D. & KRUM, H. (2004). Cardiovascular role of urotensin II: effect of chronic infusion in the rat. *Peptides*, **25**, 1783–1788.
- LANGHAM, R.G., KELLY, D.J., GOW, R.M., ZHANG, Y., DOWLING, J.K., THOMSON, N.M. & GILBERT, R.E. (2004). Increased expression of urotensin II and urotensin II receptor in human diabetic nephropathy. *Am. J. Kidney Dis.*, **44**, 826–831.
- LAPP, H., BOERRIGTER, G., COSTELLO-BOERRIGTER, L.C., JAEKEL, K., SCHEFFOLD, T., KRAKAU, I., SCHRAMM, M., GUELKER, H. & STASCH, J.P. (2004). Elevated plasma human urotensin-II-like immunoreactivity in ischemic cardiomyopathy. *Int. J. Cardiol.*, **94**, 93–97.
- LESLIE, S.J., DENVIR, M. & WEBB, D.J. (2000). Human urotensin II causes vasoconstriction in the human skin microcirculation. *Circulation*, **102**, II-113.
- LEW, M.J. & ANGUS, J.A. (1997). An improved method for analysis of competitive agonist/antagonist interactions by non-linear regression. *Ann. N.Y. Acad. Sci.*, **812**, 179–181.
- LIM, M., HONISSETT, S., SPARKES, C.D., KOMESAROFF, P., KOMPA, A. & KRUM, H. (2004). Differential effect of urotensin II on vascular tone in normal subjects and patients with chronic heart failure. *Circulation*, **109**, 1212–1214.
- MACLEAN, M.R., ALEXANDER, D., STIRRAT, A., GALLAGHER, M., DOUGLAS, S.A., OHLSTEIN, E.H., MORECROFT, I. & POLLAND, K. (2000). Contractile responses to human urotensin-II in rat and human pulmonary arteries: effect of endothelial factors and chronic hypoxia in the rat. *Br. J. Pharmacol.*, **130**, 201–204.
- MAGUIRE, J.J. & DAVENPORT, A.P. (2002). Is urotensin-II the new endothelin? *Br. J. Pharmacol.*, **137**, 579–588.
- MAGUIRE, J.J., KUC, R.E. & DAVENPORT, A.P. (2000). Orphan-receptor ligand human urotensin II: receptor localization in human tissues and comparison of vasoconstrictor responses with endothelin-1. *Br. J. Pharmacol.*, **131**, 441–446.
- MAGUIRE, J.J., KUC, R.E., WILEY, K.E., KLEINZ, M.J. & DAVENPORT, A.P. (2004). Cellular distribution of immunoreactive urotensin-II in human tissues with evidence of increased expression in atherosclerosis and greater constrictor response of small compared to large coronary arteries. *Peptides*, **25**, 1767–1774.

- MATSUSHITA, M., SHICHIRI, M., IMAI, T., IWASHINA, M., TANAKA, H., TAKASU, N. & HIRATA, Y. (2001). Co-expression of urotensin II and its receptor (GPR14) in human cardiovascular and renal tissues. *J. Hypertens.*, **19**, 2185–2190.
- NG, L.L., LOKE, I., O'BRIEN, R.J., SQUIRE, I.B. & DAVIES, J.E. (2002). Plasma urotensin in human systolic heart failure. *Circulation*, **106**, 2877–2880.
- ONAN, D. (2004). Cardiovascular and regulatory aspects of the urotensin-II receptor. PhD Thesis, Monash University, Melbourne, Australia.
- ONAN, D., PIPOLO, L., YANG, E., HANNAN, R.D. & THOMAS, W.G. (2004). Urotensin II promotes hypertrophy of cardiac myocytes via mitogen-activated protein kinases. *Mol. Endocrinol.*, **18**, 2344–2354.
- PATACCHINI, R., SANTICIOLI, P., GIULIANI, S., GRIECO, P., NOVELLINO, E., ROVERO, P. & MAGGI, C.A. (2003). Urantide: an ultrapotent urotensin II antagonist peptide in the rat aorta. *Br. J. Pharmacol.*, **140**, 1155–1158.
- PAYSANT, J., RUPIN, A., SIMONET, S., FABIANI, J.N. & VERBEUREN, T.J. (2001). Comparison of the contractile responses of human coronary bypass grafts and monkey arteries to human urotensin-II. *Fund. Clin. Pharmacol.*, **15**, 227–231.
- RICHARDS, A.M. & CHARLES, C. (2004). Urotensin II in the cardiovascular system. *Peptides*, **25**, 1795–1802.
- RICHARDS, A.M., NICHOLLS, M.G., LAINCHBURY, J.G., FISHER, S. & YANDLE, T.G. (2002). Plasma urotensin II in heart failure. *Lancet*, **360**, 545–546.
- RUSSELL, F.D. (2004). Emerging roles of urotensin-II in cardiovascular disease. *Pharmacol. Ther.*, **103**, 223–243.
- RUSSELL, F.D., MEYERS, D., GALBRAITH, A.J., BETT, N., TOTH, I., KEARNS, P. & MOLENAAR, P. (2003). Elevated plasma levels of human urotensin-II immunoreactivity in congestive heart failure. *Am. J. Physiol.*, **285**, H1576–H1581.
- RUSSELL, F.D., MOLENAAR, P. & O'BRIEN, D.M. (2001). Cardiostimulant effects of urotensin-II in human heart *in vitro*. *Br. J. Pharmacol.*, **132**, 5–9.
- SHENOUDA, A., DOUGLAS, S.A., OHLSTEIN, E.H. & GIAID, A. (2002). Localization of urotensin-II immunoreactivity in normal human kidneys and renal carcinoma. *J. Histochem. Cytochem.*, **50**, 885–889.
- SONG, W., ASHTON, N. & BALMENT, R.J. (2003). Effects of single bolus urotensin II injection in anaesthetised rats. *J. Physiol.*, **552P**, P107.
- TOTSUNE, K., TAKAHASHI, K., ARIHARA, Z., SONE, M., ITO, S. & MURAKAMI, O. (2003). Increased plasma urotensin II levels in patients with diabetes mellitus. *Clin. Sci.*, **104**, 1–5.
- TOTSUNE, K., TAKAHASHI, K., ARIHARA, Z., SONE, M., MURAKAMI, O., ITO, S., KIKUYA, M., OHKUBO, T., HASHIMOTO, J. & IMAI, Y. (2004). Elevated plasma levels of immunoreactive urotensin II and its increased urinary excretion in patients with Type 2 diabetes mellitus: association with progress of diabetic nephropathy. *Peptides*, **25**, 1809–1814.
- TOTSUNE, K., TAKAHASHI, K., ARIHARA, Z., SONE, M., SATOH, F., ITO, S., KIMURA, Y., SASANO, H. & MURAKAMI, O. (2001). Role of urotensin II in patients on dialysis. *Lancet*, **358**, 810–811.
- TZANIDIS, A., HANNAN, R.D., THOMAS, W.G., ONAN, D., AUTELITANO, D.J., SEE, F., KELLY, D.J., GILBERT, R.E. & KRUM, H. (2003). Direct actions of urotensin II on the heart: implications for cardiac fibrosis and hypertrophy. *Circ. Res.*, **93**, 246–253.
- VERGURA, R., CAMARDA, V., RIZZI, A., SPAGNOL, M., GUERRINI, R., CALO', G., SALVADORI, S. & REGOLI, D. (2004). Urotensin II stimulates plasma extravasation in mice via UT receptor activation. *Naunyn Schmiedeberg's Arch. Pharmacol.*, **370**, 347–352.
- WATANABE, T., PAKALA, R., KATAGIRI, T. & BENEDICT, C.R. (2001). Synergistic effect of urotensin II with mildly oxidized LDL on DNA synthesis in vascular smooth muscle cells. *Circulation*, **104**, 16–18.
- WATSON, A.M.D., LAMBERT, G.W., SMITH, K.J. & MAY, C.N. (2003). Urotensin II acts centrally to increase epinephrine and ACTH release and cause potent inotropic and chronotropic actions. *Hypertension*, **42**, 373–379.
- WENYI, Z., SUZUKI, S., HIRAI, M., HINOKIO, Y., TANIZAWA, Y., MATSUTANI, A., SATOH, J. & OKA, Y. (2003). Role of urotensin II gene in genetic susceptibility to Type 2 diabetes mellitus in Japanese subjects. *Diabetologia*, **46**, 972–976.
- WILKINSON, I.B., AFFOLTER, J.T., DE HAAS, S.L., PELLEGRINI, M.P., BOYD, J., WINTER, M.J., BALMENT, R.J. & WEBB, D.J. (2002). High plasma concentrations of human urotensin II do not alter local or systemic hemodynamics in man. *Cardiovasc. Res.*, **53**, 341–347.
- ZOU, Y., NAGAI, R. & YAMAACKI, T. (2001). Urotensin II induces hypertrophic responses in cultured cardiomyocytes from neonatal rats. *FEBS Lett.*, **508**, 57–60.

(Received January 24, 2005)

Revised March 7, 2005

Accepted March 10, 2005

Published online 25 April 2005