

Perfluoroalkyl and Polyfluoroalkyl Substances in the Environment: Terminology, Classification, and Origins

Robert C Buck,[†] James Franklin,^{*‡} Urs Berger,[§] Jason M Conder,^{||} Ian T Cousins,[§] Pim de Voogt,[#]
Allan Astrup Jensen,^{††} Kurunthachalam Kannan,^{‡‡} Scott A Mabury,^{§§} and Stefan PJ van Leeuwen^{|||}

[†]E.I. du Pont de Nemours & Co., Inc., DuPont Chemicals and Fluoroproducts, Wilmington, Delaware, USA

[‡]CLF-Chem Consulting, 3 Clos du Châtaignier, BE-1390 Grez-Doiceau, Belgium

[§]Department of Applied Environmental Science (ITM), Stockholm University, Stockholm, Sweden

^{||}ENVIRON International, Irvine, California, USA

[#]Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, Amsterdam, The Netherlands

^{††}Nordic Institute for Product Sustainability, Environmental Chemistry and Toxicology (NIPSECT), Frederiksberg, Denmark

^{‡‡}Wadsworth Center, New York State Department of Health, and Department of Environmental Health Sciences, School of Public Health, State University of New York at Albany, Albany, New York, USA

^{§§}Department of Chemistry, University of Toronto, Toronto, Ontario, Canada

^{|||}RIKILT—Institute of Food Safety, Wageningen, The Netherlands

(Submitted 2 May 2011; Returned for Revision 26 May 2011; Accepted 5 July 2011)

ABSTRACT

The primary aim of this article is to provide an overview of perfluoroalkyl and polyfluoroalkyl substances (PFASs) detected in the environment, wildlife, and humans, and recommend clear, specific, and descriptive terminology, names, and acronyms for PFASs. The overarching objective is to unify and harmonize communication on PFASs by offering terminology for use by the global scientific, regulatory, and industrial communities. A particular emphasis is placed on long-chain perfluoroalkyl acids, substances related to the long-chain perfluoroalkyl acids, and substances intended as alternatives to the use of the long-chain perfluoroalkyl acids or their precursors. First, we define PFASs, classify them into various families, and recommend a pragmatic set of common names and acronyms for both the families and their individual members. Terminology related to fluorinated polymers is an important aspect of our classification. Second, we provide a brief description of the 2 main production processes, electrochemical fluorination and telomerization, used for introducing perfluoroalkyl moieties into organic compounds, and we specify the types of byproducts (isomers and homologues) likely to arise in these processes. Third, we show how the principal families of PFASs are interrelated as industrial, environmental, or metabolic precursors or transformation products of one another. We pay particular attention to those PFASs that have the potential to be converted, by abiotic or biotic environmental processes or by human metabolism, into long-chain perfluoroalkyl carboxylic or sulfonic acids, which are currently the focus of regulatory action. The Supplemental Data lists 42 families and subfamilies of PFASs and 268 selected individual compounds, providing recommended names and acronyms, and structural formulas, as well as Chemical Abstracts Service registry numbers. Integr Environ Assess Manag 2011;7:513–541. © 2011 SETAC

Keywords: Perfluoroalkyl Polyfluoroalkyl Terminology Acronyms PFAS

INTRODUCTION

“Fluorinated substances” is a general, nonspecific name that describes a universe of organic and inorganic substances that contain at least 1 F atom, with vastly different physical, chemical, and biological properties (Banks et al. 1994). Synonyms include “fluorochemicals” and “fluorinated chemicals.” A subset of fluorinated substances is the highly fluorinated aliphatic substances that contain 1 or more C atoms on which all the H substituents (present in the nonfluorinated analogues from which they are notionally derived) have been replaced by F atoms, in such a manner that they contain the perfluoroalkyl moiety $C_nF_{2n+1}-$. These compounds are hereafter referred to as “perfluoroalkyl and polyfluoroalkyl substances” and denoted by the acronym PFASs, justification for the choice of which is provided below.

Since 1950, PFASs and surfactants and polymers made with the aid of PFASs have been widely used in numerous industrial and commercial applications (Kissa 2001). The C–F bond is extremely strong and stable (Smart 1994). The chemical and thermal stability of a perfluoroalkyl moiety, in addition to its hydrophobic and lipophobic nature, lead to highly useful and enduring properties in surfactants and polymers into which the perfluoroalkyl moiety is incorporated (Kissa 1994, 2001). Polymer applications include textile stain and soil repellents and grease-proof, food-contact paper (Rao and Baker 1994). Surfactant applications that take advantage of the unparalleled aqueous surface tension-lowering properties include processing aids for fluoropolymer manufacture, coatings, and aqueous film-forming foams (AFFFs) used to extinguish fires involving highly flammable liquids (Kissa 1994; Taylor 1999; Kissa 2001). Numerous additional applications have been described (3M Company 1999; Kissa 2001).

As a consequence of the widespread use of PFASs and their resulting emissions, a broad range of these substances have been detected in the environment, wildlife, and humans. The global extent of such contamination was first demonstrated

All Supplemental Data may be found in the online version of this article.

* To whom correspondence may be addressed: james.franklin@skynet.be

Published online 25 July 2011 in Wiley Online Library

(wileyonlinelibrary.com).

DOI: 10.1002/ieam.258

for perfluorooctane sulfonic acid, $C_8F_{17}SO_3H$ (PFOS) in wildlife by Giesy and Kannan (2001). (It should be noted that, throughout this article, we refer to all PFASs containing an acid functionality as “acids,” regardless of whether or not they are likely to be highly or completely ionized in environmental or human matrices). At about the same time as the study by Giesy and Kannan, Hansen et al. (2001) discovered that PFOS, perfluorooctanoic acid (PFOA, $C_7F_{15}COOH$), and other PFASs were present in numerous samples of human blood purchased from biological supply companies. This latter study suggested that PFASs were responsible for a substantial fraction of the organic F detected in human serum in earlier pioneering studies on individuals not occupationally exposed to PFASs (e.g., Taves 1968; Belisle 1981). The blood of a group of fluorochemical industry workers had already been confirmed to contain PFOA (Ubel et al. 1980). The relative significance of various human exposure pathways for PFOS, PFOA, and related substances, i.e., via food, food-contact materials, drinking water, breast milk, airborne dust, air, and so forth, is a crucially important question that has been the focus of much research, reviewed recently by D'Hollander et al. (2010). Another important research topic, directly related to exposure of humans and wildlife, is the question of how and how fast PFOS and PFOA, as well as their homologues and precursors, are transported away from their emission sources over long distances in air and/or water (Armitage et al. 2006; Prevedouros et al. 2006; Wallington et al. 2006; Yarwood et al. 2007; Wania 2007; Schenker et al. 2008; Armitage et al. 2009a, 2009b; Stemmler and Lammel 2010).

The global regulatory community is specifically interested in “long-chain” perfluoroalkyl sulfonic acids ($C_nF_{2n+1}SO_3H$, $n \geq 6$, PFASs) and perfluoroalkyl carboxylic acids ($C_nF_{2n+1}COOH$, $n \geq 7$, PFCAs) and their corresponding anions (USEPA 2009; OECD 2011), which have been shown to be more bioaccumulative than their short-chain analogues (Martin et al. 2003a, 2003b; Conder et al. 2008; Olsen et al. 2009). PFOS and PFOA are the 2 “long-chain” perfluoroalkyl acids most often reported and discussed in the scientific literature.

As explained, for example, by Paul et al. (2009) and Prevedouros et al. (2006), the presence of PFOS, PFOA, and similar substances in the environment originates from the industrial use and environmental release of these substances, from use and disposal of consumer products that may contain them as an impurity, and from the abiotic or biotic degradation of larger functional derivatives and polymers that contain a perfluoroalkyl moiety and degrade in the environment to form PFOS, PFOA, and similar substances. These precursor substances are more commonly used commercially and may be released to the environment from industrial raw materials and products and from consumer products and articles.

Concerns about the potential environmental and toxicological impact of long-chain PFASs and PFCAs have led to: 1) the phase-out of production of PFOS and related compounds and PFOA by their major global manufacturer in 2000 to 2002 (3M Company 2000a; USEPA 2000); 2) the conclusion of a stewardship agreement between the US Environmental Protection Agency (USEPA) and 8 leading global companies to reduce emissions and product content of PFOA and related chemicals by 95% by 2010 and to work toward their elimination by 2015 (USEPA 2006b); 3) a similar agreement

between the Canadian environmental and health authorities and 5 companies to restrict PFCAs in products (Environment Canada 2010); 4) a European Union Marketing and Use Directive restricting the use of “perfluorooctane sulfonates” in the European Union (European Parliament 2006b); 5) the inclusion of PFOS in the Stockholm Convention on Persistent Organic Pollutants as an Annex B substance, i.e., restricted in its use (UNEP 2009); and 6) other regulatory and voluntary initiatives intended to reduce environmental emissions of this family of compounds.

The concern over potential environmental and human health impacts of PFASs has led to the launching of several large research programs to elucidate their environmental origin, fate, and impact, funded by various authorities in, for example, the European Union (de Voogt et al. 2006; de Voogt 2009), the United States (USEPA 2010), and Canada (INAC 2009). Moreover, alternative PFASs intended to be replacements for the long-chain PFASs and PFCAs have been developed and implemented in certain cases (Visca et al. 2003; Higuchi et al. 2005; Hintzer et al. 2005; Brothers et al. 2008; Ishikawa et al. 2008; Peschka et al. 2008; Gordon 2011).

Since the first reports revealing the widespread global occurrence of PFOS in wildlife (Giesy and Kannan 2001) and the frequent detection of PFASs in human blood (Hansen et al. 2001) were published a decade ago, the scientific literature on the environmental and toxicological aspects of PFASs has burgeoned rapidly, and the rate of publication currently exceeds 400 articles per year. In the existing body of literature, including governmental reports, authors have created terminology, names, and acronyms to describe these substances. Unfortunately, inconsistencies have inevitably arisen between various groups of authors. In the absence of any concerted effort between scientists to agree on a common terminology to designate the substances, a given compound has often been denoted by a variety of different names and acronyms, or a given acronym has been used to represent different substances. In addition, names to describe broad groups of substances have proliferated that in some instances mistakenly portray substances that are very different from one another as being the same. As a result, the scientific literature for these substances has at times become confusing. There is a need for harmonized terminology, names, and acronyms that clearly and specifically describe PFASs.

OBJECTIVES

The primary aim of this article is to recommend clear, specific, and descriptive terminology, names, and acronyms for PFASs, so as to promote a sound, unified understanding among all players in the PFAS industry, the environmental science related to it, and the bodies responsible for the regulation of chemicals, hence facilitating meaningful communication among all concerned.

A particular emphasis is placed on the long-chain perfluoroalkyl acids, substances related to the long-chain perfluoroalkyl acids, and substances intended as alternatives to the use of the long-chain perfluoroalkyl acids or their precursors. We trust that the terminology, names, and acronyms suggested will be broadly adopted by the “perfluoroalkyl and polyfluoroalkyl substances community” at large, leading to harmonized usage and the avoidance of

misnomers. We have nevertheless refrained from creating an all new nomenclature but have retained—as far as possible—the most popular terms and acronyms used by authors to date. In other words, our proposals result from a pragmatic compromise among textbook and/or International Union for Pure and Applied Chemistry (IUPAC) chemical nomenclature, universal consistency, and frequently adopted “legacy” usage.

It is important to note that the substance terminology, names and acronyms proposed in this article are in no way intended to compete with or supplant IUPAC or Chemical Abstracts Service (CAS) nomenclature. The latter names are the designations of choice when a specific substance needs to be unequivocally identified, e.g., in official regulatory documents. Our intention is to provide terminology, names, and acronyms for pragmatic everyday use within the scientific community. Thus, for example, the IUPAC name for the substance $\text{C}_8\text{F}_{17}\text{SO}_2\text{N}(\text{C}_2\text{H}_5)\text{CH}_2\text{CH}_2\text{OH}$ is “N-ethyl-1,1,2,2,3,3,4,4,5,5,6,6,7,7,8,8,8-heptafluorooctane-1-sulfonamide,” but it is more convenient to use the less rigorous but shorter designation “N-ethyl perfluorooctane sulfonamidoethanol” (or the corresponding acronym EtFOSE) for use in publications aimed at specialist readers. Rigor can always be ensured by appending the appropriate CAS Registry Number when each compound is first mentioned in a publication. We encourage this practice and provide CAS numbers for many commonly discussed compounds in the Supplemental Data.

In addition to recommending terminology, names, and acronyms, this article provides a brief review of certain topics useful for understanding the occurrence of and relationships between various families of PFASs in the environment. First, we describe the major commercial processes for synthesizing perfluoroalkyl moieties and the resulting compositions, including formation of isomers and/or homologues of the targeted main products. Second, we present the interrelationships between families of PFASs that may be precursors to or products of one another as a result of abiotic or biotic transformations that may occur under industrial, environmental, or metabolic conditions.

A large number of PFASs have been commercially produced (OECD 2007), and not all are covered here. We have included the main families, individual compounds, and their degradation products that have been detected in environmental and human samples related to long-chain perfluoroalkyl acids, precursors to these substances, and their short-chain fluorinated alternatives. We provide literature references for studies that demonstrate how one family of PFASs may be transformed into another under abiotic or biotic conditions, and/or report the presence of the various families in the environment or humans. Nevertheless, given the vast number of publications on the most common PFASs, such as the perfluoroalkyl sulfonic and carboxylic acids and their anions and salts, the reader is referred to published reviews and extensive surveys for comprehensive literature compilations for these compounds (e.g., Kannan et al. 2004; Houde et al. 2006; Lau et al. 2007; van Leeuwen and de Boer 2007; Jahnke and Berger 2009; Loos et al. 2009; Pistocchi and Loos 2009; Rayne and Forest 2009b; Butt, Berger, et al. 2010; de Voogt 2010; Kwok et al. 2010; Loos et al. 2010; Sturm and Ahrens 2010; Ahrens 2011; Houde et al. 2011). Furthermore, because an emphasis here is on how the various categories of PFASs are interrelated, our citations on transformation

processes and environmental presence often refer to families of substances, so the reader should consult the original publications for details on individual substances.

It should be noted that in this article, the terms “substance,” “compound,” “chemical,” and “species” are used interchangeably for designating a given molecular structure, although it is recognized that in other contexts their meanings may not be identical. For example, in the European REACH legislation (European Parliament 2006a), a “substance” may include impurities and stabilizers in addition to the main constituent.

KEY TERMINOLOGY AND USAGE ASSOCIATED WITH PERFLUOROALKYL AND POLYFLUOROALKYL SUBSTANCES

Perfluoroalkyl and polyfluoroalkyl substances and perfluorocarbons defined

As defined above, PFASs are aliphatic substances containing one or more C atoms on which all the H substituents present in the nonfluorinated analogues from which they are notionally derived have been replaced by F atoms, in such a manner that PFASs contain the perfluoroalkyl moiety $\text{C}_n\text{F}_{2n+1}-$. More explicitly, we recommend that the family of compounds denoted by the acronym PFAS should encompass:





- Perfluoroalkyl substances, which are defined as aliphatic substances for which *all* of the H atoms attached to C atoms in the nonfluorinated substance from which they are notionally derived have been replaced by F atoms, except those H atoms whose substitution would modify the nature of any functional groups present. This usage is consistent with the definition of “perfluoro” and “perfluorinated” provided by Banks et al. (1994, p. 2).
- Polyfluoroalkyl substances, defined here as aliphatic substances for which all H atoms attached to at least one (but not all) C atoms have been replaced by F atoms, in such a manner that they contain the perfluoroalkyl moiety $\text{C}_n\text{F}_{2n+1}-$ (e.g., $\text{C}_8\text{F}_{17}\text{CH}_2\text{CH}_2\text{OH}$). Thus, whereas the general chemical concept of “polyfluorination” embraces compounds containing “scattered” multiple F atoms (such as in $\text{CH}_2\text{FCH}_2\text{FCH}_2\text{FCH}_2\text{OH}$), as well as “grouped” ones (such as in $\text{CF}_3\text{CF}_2\text{CH}_2\text{COOH}$), we consider that only those polyfluorinated substances having at least one perfluoroalkyl moiety $\text{C}_n\text{F}_{2n+1}-$ belong to the PFAS family.

The differences between perfluoroalkyl and polyfluoroalkyl substances are illustrated by 2 concrete examples in Table 1.

Polyfluoroalkyl substances have the potential (i.e., the demonstrated or theoretical capability under appropriate conditions) to be transformed abiotically or biotically into perfluoroalkyl substances. For example, $\text{C}_n\text{F}_{2n+1}\text{SO}_2\text{NHCH}_2\text{CH}_2\text{OH}$ (a polyfluoroalkyl substance) may degrade in the environment to $\text{C}_n\text{F}_{2n+1}\text{SO}_3\text{H}$ (a perfluoroalkyl substance).

The general term “perfluoroalkyl(ated) substance,” with the acronym PFAS, was the first to be defined and widely used to describe the broad class of highly fluorinated substances observed in the environment (Hekster et al. 2002; Hekster et al. 2003). It has been employed by the groups of scientists who collaborated in the finalized European Union PERFORCE project (de Voogt et al. 2006) and others who have

Table 1. Examples of the correct and incorrect (or undesirable) uses of the proposed nomenclature for perfluoroalkyl and polyfluoroalkyl substances (PFASs)

Example	Example statements	
	Correct	Incorrect or <i>undesirable</i>
	<ul style="list-style-type: none"> Both are PFASs, within the family of perfluoroalkyl and polyfluoroalkyl substances 	<ul style="list-style-type: none"> Both are: <ul style="list-style-type: none"> Perfluoroalkyl substances, chemicals, compounds Perfluorinated substances, chemicals, compounds Polyfluoroalkyl substances Polyfluorinated substances Fluorocarbons Perfluorocarbons Fluorinated substances, chemicals, compounds Perfluorochemicals Perfluorinated chemicals Both contain fluorocarbons
	<ul style="list-style-type: none"> Both are carboxylic acids 	
	<ul style="list-style-type: none"> All H atoms on all C atoms in the alkyl chain attached to the carboxylic acid functional group are replaced by F This is a: PFAS, perfluoroalkyl acid (PFAA), perfluoroalkyl carboxylic acid (PFCA) Specifically, this is perfluorooctanoic acid, CAS number 335-67-1 	<ul style="list-style-type: none"> This is a: <ul style="list-style-type: none"> Perfluorinated substance, chemical, compound Fluorinated substance, chemical, compound Fluorocarbon Perfluorocarbon
	<ul style="list-style-type: none"> The alkyl chain attached to the carboxylic acid functional group is polyfluorinated This is a: PFAS, polyfluoroalkyl acid, polyfluoroalkyl carboxylic acid Specifically, this is 2,2,3,3,4,4,5,5,7,7,8,8- tridecafluorooctanoic acid 	<ul style="list-style-type: none"> This is a: <ul style="list-style-type: none"> Polyfluorinated substance, chemical, compound Fluorinated substance, chemical, compound Perfluorinated substance, chemical, compound A portion of this compound is perfluorinated

followed their example. Soon thereafter, many authors began using the acronym PFC and have defined it in many different ways. As a result, the meaning of the acronym PFC is unclear and not well defined. Moreover, we consider this choice to have been an unfortunate and inappropriate one, given that the acronym PFC has been used in official Kyoto Protocol documents since its adoption in 1997 to specifically designate perfluorocarbons (United Nations 1998), one of the families of greenhouse gases regulated by this important multilateral international agreement. Clearly, a given acronym may legitimately be used in different spheres of activity to denote different concepts, provided these activities are sufficiently disconnected from each other. However, both PFCs and PFASs belong to the overall family of fluorinated chemicals and, hence, are too closely related to share a common acronym. We, therefore, strongly urge the community to adopt henceforth the use of the term PFASs (singular PFAS) as an acronym for “perfluoroalkyl and polyfluoroalkyl substances” and the term PFCs (singular PFC) exclusively for “perfluorocarbons.” PFCs are notionally derived from hydrocarbons by replacing all H atoms by F atoms, so that they contain only the elements C and F, and functional groups are absent. Examples of PFCs are tetrafluoromethane (CF₄), hexafluoroethane (C₂F₆), octafluorocyclobutane (c-C₄F₈),

and perfluorodecalin (C₁₀F₁₈). Those PFCs that contain a C_nF_{2n+1} moiety are, by definition, members of the PFAS family, but PFCs are chemically very stable substances, and it is uncertain whether any of them can actually degrade in the environment (e.g., in the upper atmosphere) to give functionalized PFASs such as PFCAAs that might ultimately be deposited to the Earth’s surface.

“Fluorinated polymers” and “fluoropolymers” defined

We recommend using the broad generic term “fluorinated polymers” to encompass all polymers for which one or more of the monomer units contains the element F, in the backbone and/or in side chains. Fluorinated polymers may or may not be PFASs, depending on whether they contain perfluoroalkyl moieties.

In compliance with time-honored usage within the industry, we recommend further that the term “fluoropolymers” be applied only to a distinct subset of fluorinated polymers, namely, those made by (co)polymerization of olefinic monomers, at least one of which contains F bound to one or both of the olefinic C atoms, to form a carbon-only polymer backbone with F atoms directly attached to it, e.g., polytetrafluoroethylene.

Chain length terminology

PFASs, especially the perfluoroalkyl acids and their anions, are frequently referred to as “long-chain” or “short-chain.” To avoid any subjectivity associated with these adjectives, we urge scientists to adopt the definition provided by the Organisation for Economic Co-operation and Development (OECD 2011), which stipulates that “long-chain” refers to:

- perfluoroalkyl carboxylic acids with eight carbons and greater (i.e., with 7 or more perfluorinated carbons) and,
- perfluoroalkane sulfonates with six carbons and greater (i.e., with 6 or more perfluorinated carbons).

The “long-chain” definitions for PFCAs and PFSA are different in number of C atoms because a PFSA (e.g., PFHxS, $C_6F_{13}SO_3H$) with a given number of carbons (6 in the example given) has a greater tendency to bioconcentrate and/or bioaccumulate than a PFCA with the same number of C atoms (e.g., PFHxA, $C_5F_{11}COOH$) (Martin et al. 2003a, 2003b). Although the OECD definition does not include perfluoroalkyl substances other than carboxylates and sulfonates, one may consider that a perfluoroalkyl chain with 7 or more C atoms, e.g., $C_7F_{15}-$, is, in any case, “long.”

Linear and branched terminology

Many PFASs exist as families of isomers due to branching of the main C backbone (Alsmeyer et al. 1994). Linear isomers, for which there can only be 1 congener per C_n homologue group, are composed of carbons that are bonded to only 1 or 2 other C atoms. Branched isomers, for which there can be several or many congeners per C_n homologue group, are composed of C atoms that may be bound to more than 2 C atoms, resulting in a branching of the C backbone. For example, PFOS is routinely present in many environmental samples as a mixture of the linear isomer and 10 branched isomers (Riddell et al. 2009), whereas 89 congeners are theoretically possible (Rayne et al. 2008). To address the characterization of the numerous isomers and homologues arising during the electrochemical fluorination process (see below), a systematic numbering system for unequivocally identifying the linear and branched congeners of several families of PFASs has been proposed (Rayne et al. 2008). In the following text and in the Supplemental Data, we will designate perfluoroalkyl moieties, in general, by the formula $C_nF_{2n+1}-$, thereby including both linear and branched structures, even for substances that, given their manufacturing process (see discussion below), may be presumed to be predominantly linear, so that $C_nF_{2n+1}-$ is equivalent to $F(CF_2)_n-$.

The mixture of linear and branched isomers presents challenges in providing an accurate quantification of many PFASs in environmental matrices (Riddell et al. 2009). Nevertheless, the study of linear and branched isomers is useful for understanding sources of PFASs (De Silva and Mabury 2004, 2006; De Silva et al. 2009; Benskin, De Silva, et al. 2010; Benskin, Yeung, et al. 2010), because the production of isomers varies by manufacturing process. The telomerization process produces primarily or exclusively linear PFASs, whereas the electrochemical fluorination process produces a mixture of branched and linear isomers, as discussed below.

Use of acronyms for acids and their anions

Many PFASs are acids and may be present as protonated or anionic forms, or a mixture of both, depending on the pH of the environmental matrix and the compound's acid dissociation constant (pK_a). The pK_a values for many of the PFASs (e.g., PFOA) are under review or are unknown, and for simplicity, we will refer to all PFASs with an acid functionality as “acids,” rather than as carboxylates, sulfonates, and so forth, although recognizing that the dissociated forms may well predominate in environmental and human matrices. Furthermore, given that these acids are generally analyzed as their anions (Larsen and Kaiser 2007), we recommend using the same substance acronym to cover both the protonated and ionized forms. However, an exception is made to this general rule when it is essential to make a distinction between the protonated acid form and the anionic form, such as when reporting physicochemical properties or modeling environmental fate and transport (Armitage et al. 2009b; Webster et al. 2010). In these cases, it is recommended to designate PFCA anions by removing the “A” from the individual substance acronym (e.g., PFO for perfluorooctanoate), maintain the original abbreviation for the acid (e.g., PFOA for perfluorooctanoic acid), and refer to both chemical forms using a collective abbreviation involving parentheses surrounding the “A,” e.g., PFO(A) for combined perfluorooctanoate and perfluorooctanoic acid. In the case of PFSA, it is suggested to add the prefix “H-” to the generic substance acronym to form the abbreviation for the neutral species. This leads, for example, to the abbreviations H-PFOS, PFOS, and (H-)PFOS for the protonated, anionic, and combined forms of the 8-C PFSA, respectively.

Surfactant terminology

Many PFASs are used as surfactants. Traditional surfactants comprise a water-soluble hydrophilic portion and a water-insoluble hydrophobic portion. Surfactants lower the surface tension of a liquid, or the interfacial tension between 2 liquids, or between a liquid and a solid. In fluorinated surfactants, the hydrophobic portion contains F bound to C, often as a perfluoroalkyl moiety. The extent of fluorination and location of the F atoms affect the surfactant properties. PFAS surfactants, often referred to as “fluorinated surfactants,” “fluorosurfactants,” “fluorinated tensides,” or “fluorotensides,” are superior in their aqueous surface tension reduction at very low concentrations and are useful as wetting and leveling agents, emulsifiers, foaming agents, or dispersants (Kissa 1994; Taylor 1999; Kissa 2001). The term “tenside” is encountered most frequently in publications of German origin, and the synonym “surfactant” is preferred in English. Examples of fluorinated surfactants are $NH_4^+ C_7F_{15}CO_2^-$ and $Na^+ C_6F_{13}CH_2CH_2SO_3^-$.

Terminology describing direct and indirect sources of PFASs to the environment

The sources of PFAS (e.g., PFOS or PFOA) emissions to the environment are from their purposeful manufacture, use, and disposal, from their being present as impurities in substances that are emitted to the environment or from precursor substances that degrade abiotically or biotically in the environment. Harmonizing the terminology for describing “sources” is needed. We recommend that the term “direct”

emission sources should refer to emissions of a specific PFAS as such, throughout its product life cycle from manufacture to use and disposal, including emissions from a product in which the PFAS is present as an impurity. On the other hand, the term “indirect” emissions should apply to formation of a specific PFAS by transformation of precursor substances in the environment, wildlife, or humans, such as PFOA formed from the biotransformation of 8:2 fluorotelomer alcohol (FTOH), or C_4F_9COOH from the atmospheric degradation of perfluorobutane sulfonamidoethanol. These definitions depart somewhat from those of Prevedouros et al. (2006) who considered emissions of impurities present in a product to be “indirect.” These alternative definitions do not create large differences in the emissions allocated to direct and indirect sources in the case of PFOA, because the majority of direct emissions are derived from manufacturing sources.

MANUFACTURING PROCESSES

For a better understanding of the environmental occurrence and behavior of PFASs, as well as the relationships between families of PFASs, it is useful to describe briefly the 2 principal manufacturing processes used to produce compounds containing perfluoroalkyl chains.

Electrochemical fluorination

Electrochemical fluorination (ECF) is a technology in which an organic raw material (e.g., octane sulfonyl fluoride [OSF], $C_8H_{17}SO_2F$) undergoes electrolysis in anhydrous HF, leading to the replacement of all the H atoms by F atoms (Alsmeyer et al. 1994). The free-radical nature of the process leads to C chain rearrangement and breakage, resulting in a mixture of linear and branched perfluorinated isomers and homologues of the raw material, as well as PFCs and other species (Alsmeyer et al. 1994). The ratio of linear to branched

perfluorinated C chains formed in the ECF process varies depending on how the process is controlled but is roughly 70% to 80% linear and 20% to 30% branched in the case of the synthesis of PFOS and PFOA (3M Company 1999; Reagen et al. 2007; Lehmler 2009; Benskin, De Silva, et al. 2010). The ECF of $C_8H_{17}SO_2F$ yields 1) perfluorooctane sulfonyl fluoride (POSF, $C_8F_{17}SO_2F$), which is the major raw material used to manufacture PFOS (Figure 1a); 2) a series of functional raw materials such as sulfonamides, sulfonamido alcohols, and sulfonamido acrylate monomers; and 3) a family of surfactants and polymers derived therefrom (3M Company 1999; Lehmler 2005). Likewise, the ECF of octanoyl fluoride, $C_7H_{15}COF$, is the major historic process used to manufacture perfluorooctanoyl fluoride, $C_7F_{15}COF$, which is further reacted to make PFOA and its salts (Figure 1b) (Kissa 1994). The major global historic manufacturer using the ECF process produced 6-, 8-, and (to a lesser extent) 10-carbon perfluoroalkane sulfonyl derivatives and products therefrom (3M Company 2000c). In 2001, the company announced it would no longer manufacture these substances or PFOA. Others continued to use the ECF process to make these substances and there are now new manufacturers of both PFOS and PFOA. The major historic manufacturer is now making alternative products using the ECF process based on perfluorobutane, rather than perfluorooctane, sulfonyl chemistry (Renner 2006; Olsen et al. 2009; Ritter 2010).

Telomerization

Telomerization (Figure 2), which is a second important process for manufacturing perfluoroalkyl substances, is a technology in which a perfluoroalkyl iodide, $C_mF_{2m+1}I$ (PFAI), most commonly pentafluoroethyl (or perfluoroethyl) iodide, C_2F_5I (PFEI), is reacted with tetrafluoroethylene, $CF_2=CF_2$ (TFE) to yield a mixture of perfluoroalkyl iodides




Figure 1. Synthesis, by electrochemical fluorination, of building blocks leading to PFOS, PFOA, and derivatives.




Figure 2. Synthesis, by telomerization, of building blocks leading to fluorotelomer alcohols.

with longer perfluorinated chains $\text{C}_m\text{F}_{2m+1}(\text{CF}_2\text{CF}_2)_n\text{I}$. The starting iodide is referred to as the “telogen” and the TFE as the “taxogen.” The product perfluoroalkyl iodide mixture is often then reacted further, in a 2nd process step, where ethylene is inserted, to give $\text{C}_m\text{F}_{2m+1}(\text{CF}_2\text{CF}_2)_n\text{CH}_2\text{CH}_2\text{I}$. The perfluoroalkyl iodides, $\text{C}_m\text{F}_{2m+1}(\text{CF}_2\text{CF}_2)_n\text{I}$, commonly known as Telomer A, resulting from telomerization, the 1st step, and the “fluorotelomer iodides,” $\text{C}_m\text{F}_{2m+1}(\text{CF}_2\text{CF}_2)_n\text{CH}_2\text{CH}_2\text{I}$, commonly known as Telomer B, formed in the 2nd step, are raw material intermediates used to produce additional building blocks that are further reacted to create a family of “fluorotelomer-based” surfactant and polymer products. This process is illustrated in Figure 2 for the synthesis of a fluorotelomer alcohol (FTOH), whereas Figure 3 shows how a range of products can be synthesized from the perfluoroalkyl iodide intermediate (exemplified for a starting PFAI with 8 C atoms).

It should be noted that, in the “X:Y” designation, e.g., 8:2 fluorotelomer alcohol ($\text{C}_8\text{F}_{17}\text{CH}_2\text{CH}_2\text{OH}$, 8:2 FTOH), used for naming fluorotelomer-based substances, X is the number of perfluorinated C atoms and Y is the number of non-fluorinated C atoms that originate from the commercial synthesis. As with products derived from ECF, the major global fluorotelomer manufacturers are making available alternative shorter-chain products, in this case based on 6 (rather than 8) perfluoroalkyl C atoms (Renner 2006; Ritter 2010).

The most widely used commercial telomerization process uses PFEI and TFE. When a linear telogen and taxogen are employed in the telomerization process, the resulting perfluoroalkyl iodides have exclusively linear perfluoroalkyl chains. If a branched and/or odd C number telogen, e.g., $(\text{CF}_3)_2\text{CFI}$, is employed and reacted with TFE, the resulting product mixture will be branched and/or will contain an odd number of C atoms, despite the incorporation of an even number of taxogen $-\text{CF}_2-$ units from the TFE. The extent to which branched and/or odd C number telogens may have been actually used in commercial practice is unclear. Such telogens have been described in patents (e.g., Katsushima et al. 1964; Millauer 1971; Grottenmüller et al. 2000), but

this does not necessarily mean that they have been employed commercially. Nevertheless, in certain environmental samples, “isopropyl branched PFCA isomers,” i.e., ones with a terminal $(\text{CF}_3)_2\text{CF}-$ group, have been observed, albeit at low levels compared to their linear counterparts, whereas other branched isomers were either absent or present at much lower levels. This is the case, inter alia, for PFCAs with 9, 11, or 13 C atoms, i.e., perfluorononanoic, perfluoroundecanoic, and perfluorotridecanoic acids (PFNA, PFUnDA, and PFTrDA, respectively), which are believed to be manufactured by the ozonation of a mixture of fluorotelomer olefins (FTOs, $\text{C}_n\text{F}_{2n+1}\text{CH}=\text{CH}_2$) (Ukihashi et al. 1977; Aoyama and Chiba 1997) and which may be formed by the environmental transformation of telomer-derived precursor PFASs. The isopropyl branched isomers of these PFCAs observed in the environment (Furdui et al. 2008; De Silva et al. 2009; Benskin, De Silva, et al. 2010; Zushi et al. 2010) may therefore originate from the use of branched telogens for manufacturing specific isomers of PFNA, PFUnDA, and PFTrDA or their precursors. Nevertheless, the interpretation of branched-to-linear isomer concentration ratios is not straightforward, because certain environmental samples were found to contain up to 3 other PFNA isomers (for example) in addition to the linear and isopropyl branched forms (De Silva and Mabury 2006; Benskin et al. 2007; De Silva et al. 2009). Furthermore, the fact that individual isomers have different physicochemical properties means the patterns in the environment and biota will be transformed relative to the pattern in the emission source.

FAMILIES OF PERFLUOROALKYL AND POLYFLUOROALKYL SUBSTANCES

There are numerous families of PFASs (Figure 4), each with many individual homologous members and isomers thereof (Tables 2, 3, and 4). This section provides a hierarchical overview of the common substance names, acronyms, and chemical formulas of those families of compounds and selected individual substances that have been detected in environmental and human matrices. The discussion includes references to manufacturing processes and




Figure 3. Perfluoroalkyl carboxylic acids and fluorotelomer (FT) derivatives synthesized from perfluoroalkyl iodides (PFAIs), exemplified for a starting PFAI with 8 C atoms. N.B. Names and acronyms for substance families are indicated. Those for the specific compounds shown can be found in the Supplemental Data.

uses for individual PFASs, as well as their environmental occurrence, for a better understanding of their environmental origin and how certain families and substances are related to one another. Another key point of the discussion is the likelihood that any or all members of PFAS groups have the ability to transform to the long-chain perfluorinated acids, provided, of course, that they have a long enough perfluoroalkyl moiety. A more comprehensive compilation of individual substances is given in the Supplemental Data, which also includes CAS registry numbers when assigned.

First, we choose to make a fundamental distinction in substances by dividing them into 2 primary categories: nonpolymers and polymers (Figure 4). It is well accepted that polymers generally have very different physical, chemical, and biological properties than discrete chemical substances of low molecular weight (e.g., methyl methacrylate versus poly[methyl methacrylate]). There are various definitions of a polymer, but the basic concept describes a substance consisting of molecules characterized by the sequence of one or more types of monomer unit. Precise criteria for distinguishing polymers from nonpolymers have been established, for instance, under the European Union REACH legislation (ECHA 2008).

Nonpolymer perfluoroalkyl and polyfluoroalkyl substances

Perfluoroalkyl acids. Perfluoroalkyl acids (PFAAs) occupy a prominent place in the literature on PFASs. The family of PFAAs includes perfluoroalkyl carboxylic, sulfonic, sulfinic, phosphonic, and phosphinic acids (Table 2). PFAAs are important both because they are highly persistent substances that have been directly emitted to the environment or are formed indirectly from the environmental degradation or

metabolism of precursor substances, and because they (or their salts) are or have been used in a wide variety of industrial and consumer applications. Depending on their acid strength (pK_a value), PFAAs will dissociate to a greater or lesser extent to their anions in aqueous environmental media, soils, or sediments. The protonated and anionic forms have very different physicochemical properties. For instance, the perfluorooctanoate anion is highly water-soluble and has negligible vapor pressure, whereas perfluorooctanoic acid has very low water solubility and sufficient vapor pressure to partition out of water into air (Kaiser et al. 2005; Kaiser et al. 2006; Webster and Ellis 2010; Webster et al. 2010). However, for perfluoroalkyl carboxylic acids, there is an ongoing debate regarding what is the environmentally relevant pK_a , with measured and estimated values varying by several log units for PFOA (Burns et al. 2008; Goss 2008; Cheng et al. 2009; Rayne and Forest 2010a).

Perfluoroalkyl carboxylic acids: Perfluoroalkyl carboxylic acids (PFCAs; Table 2), also known as perfluorocarboxylic acids or perfluoroalkanoic acids, have the general chemical formula $C_nF_{2n+1}COOH$. The most frequently discussed PFCA is PFOA, $C_7F_{15}COOH$. The ammonium salt of PFOA, ammonium perfluorooctanoate (APFO, $NH_4^+ C_7F_{15}COO^-$) has been used for many decades as an essential “processing aid” in the manufacture of fluoropolymers such as polytetrafluoroethylene, by the dispersion (or emulsion) process (Kissa 1994; Fluoropolymer Manufacturing Group 2001). A chemically inert perfluorinated surfactant is chosen for this application to avoid reaction of the growing free-radical polymer chains with the processing aid, which would lead to a lowering of the molecular weight of the polymer produced. APFO and derivatives of it were also produced and marketed for fluorosurfactant use (3M Company 2000a). Between 1947




Figure 4. Classification hierarchy of environmentally relevant perfluoroalkyl and polyfluoroalkyl substances (PFASs).

and 2002, APFO was manufactured by multiple companies around the world, probably mainly or exclusively by ECF of octanoyl fluoride. In 2002, the major global historic APFO manufacturer ceased its production (3M Company 2000a, 2000c). Thus, in addition to continued ECF-based APFO production from the remaining ECF producers, a process in which linear perfluorooctyl iodide (PFOI) synthesized by telomerization is converted into PFOA was brought on-stream in late 2002 to meet the need for this critical raw material (Prevedouros et al. 2006). This new telomerization-based process leads to only linear PFOA, whereas the ECF process produces a mixture of linear (70%–80%) and branched PFOA isomers.

Perfluorononanoic acid, $C_8F_{17}COOH$ (PFNA) has also been manufactured and used (from 1975 onward) as its ammonium salt, $NH_4^+ C_8F_{17}COO^-$ (APFN), principally for producing fluoropolymer dispersions, especially polyvinylidene fluoride (PVDF) (Prevedouros et al. 2006). It has also been marketed for general use as a fluorinated surfactant. A sample of commercial “APFN,” known as Surflon[®] S-111, has been analyzed and shown to contain significant propor-

tions of the ammonium salts of longer PFCA homologues, especially those with 11 (PFUnDA) and 13 (PFTrDA) C atoms, which amounted to 20 and 5 weight percent of the mixture, respectively (Prevedouros et al. 2006; in the supporting information). The presence of these homologues with 2 and 4 additional C atoms, as confirmed by an industrial user (van der Putte et al. 2010), indicates that Surflon[®] S-111 is derived from a mixture of fluorotelomer-based precursors and, hence, suggests that it is constituted, predominantly or exclusively, of linear isomers. These conclusions are consistent with patents that claim manufacture of PFNA from telomer-based raw materials, namely by the oxidation of 8:2 fluorotelomer olefin, $C_8F_{17}CH=CH_2$ (Ukihashi et al. 1977; Aoyama and Chiba 1997) or by the carboxylation of $C_8F_{17}I$ (Nagasaki et al. 1988). The APFN commercial mixture has its own CAS Registry Number: 72968-38-8. Several publications report toxicological studies on the blend corresponding to this number, but do not provide information on the proportions or linearity of the homologues present (Mundt et al. 2007; Stump et al. 2008; Mertens et al. 2010).

Table 2. Hierarchical overview of the nonpolymer perfluoroalkyl substances, compounds for which all H atoms on all C atoms in the alkyl chain attached to the functional group have been replaced with F

Classification and chemical structure	$C_nF_{2n+1}R$, where R =	Examples	Uses
Perfluoroalkyl acids (PFAAs)			
Perfluoroalkyl carboxylic acids (PFCAs) ^a	-COOH	Perfluorooctanoic acid (PFOA), $C_7F_{15}COOH$	Surfactant
Perfluoroalkyl carboxylates (PFCAs) ^a	-COO ⁻	Perfluorooctanoate (PFOA), $C_7F_{15}COO^-$	
Perfluoroalkane sulfonic acids (PFSA) ^b	-SO ₃ H	Perfluorooctane sulfonic acid (PFOS), $C_8F_{17}SO_3H$	Surfactant
Perfluoroalkane sulfonates (PFSA) ^b	-SO ₃ ⁻	Perfluorobutane sulfonic acid (PFBS), $C_4F_9SO_3H$	
		Perfluorooctane sulfonate (PFOS), $C_8F_{17}SO_3^-$	
		Perfluorobutane sulfonate (PFBS), $C_4F_9SO_3^-$	
Perfluoroalkane sulfinic acids (PFSIAs) ^b	-SO ₂ H	Perfluorooctane sulfinic acid (PFOSI), $C_8F_{17}SO_2H$	Intermediate environmental transformation product
Perfluoroalkyl phosphonic acids (PFPAAs) ^c	-P(=O)(OH) ₂	Perfluorooctyl phosphonic acid (C8-PFPA) $C_8F_{17}P(=O)(OH)_2$	Surfactant
Perfluoroalkyl phosphinic acids (PFPIAs) ^c	-P(=O)(OH)(C _m F _{2m+1})	Bis(perfluorooctyl) phosphinic acid (C8/C8-PFPIA) $C_8F_{17}P(=O)(OH)(C_8F_{17})$	Surfactant
Perfluoroalkane sulfonyl fluorides (PASFs) ^b	-SO ₂ F	Perfluorooctane sulfonyl fluoride (POSF), $C_8F_{17}SO_2F$	Major raw material for surfactant and surface protection products
		Perfluorobutane sulfonyl fluoride (PBSF), $C_4F_9SO_2F$	
Perfluoroalkane sulfonamides (FASAs) ^b	-SO ₂ NH ₂	Perfluorooctane sulfonamide (FOSA), $C_8F_{17}SO_2NH_2$	Major raw material for surfactant and surface protection products
Perfluoroalkanoyl fluorides (PAFs) ^b	-COF	Perfluorooctanoyl fluoride (POF), $C_7F_{15}COF$	Major raw material for PFOA made by the ECF process; raw material for surfactant and surface protection products
Perfluoroalkyl iodides (PFAIs) (Telomer A) ^c	-I	Perfluorohexyl iodide (PFHxI), $C_6F_{13}I$	Major raw material for surfactant and surface protection products
Perfluoroalkyl aldehydes (PFALs) and aldehyde hydrates (PFAL-H ₂ O) ^c	-CHO and -CH(OH) ₂	Perfluorononanal (PFNAL), $C_8F_{17}CHO$	Intermediate environmental transformation product

^aSubstances originating by either electrochemical fluorination (ECF) or fluorotelomer processes;^bSubstances originating by the ECF process;^cSubstances originating by the fluorotelomer process.

Table 3. Hierarchical overview of the nonpolymer polyfluoroalkyl substances: compounds for which all H atoms on at least one (but not all) C atoms have been replaced with F

	Classification and chemical structure	$C_nF_{2n+1}R$, where $R =$	Examples	Uses
Perfluoroalkane sulfonamido substances ^a	<i>N</i> -Alkyl perfluoroalkane sulfonamides (MeFASAs, EtFASAs, BuFASAs)	$-SO_2NH(R')$ where $R' = C_mH_{2m+1}$ ($m = 1, 2, 4$)	<i>N</i> -Methyl perfluorooctane sulfonamide (MeFOSA), $C_8F_{17}SO_2N(CH_3)H$	Major raw material for surfactant and surface protection products
			<i>N</i> -Ethyl perfluorobutane sulfonamide (EtFBSA), $C_4F_9SO_2N(C_2H_5)H$	
			<i>N</i> -Butyl perfluorooctane sulfonamide (BuFOSA), $C_8F_{17}SO_2N(C_4H_9)H$	
	Perfluoroalkane sulfonamidoethanols (FASEs) and <i>N</i> -alkyl perfluoroalkane sulfonamidoethanols (MeFASEs, EtFASEs, BuFASEs)	$-SO_2N(R')CH_2CH_2OH$ where $R' = C_mH_{2m+1}$ ($m = 0, 1, 2, 4$)	Perfluorooctane sulfonamidoethanol (FOSE), $C_8F_{17}SO_2NHCH_2CH_2OH$	Major raw material for surfactant and surface protection products
			<i>N</i> -Ethyl perfluorobutane sulfonamidoethanol (EtFBSE), $C_4F_9SO_2N(C_2H_5)CH_2CH_2OH$	
	<i>N</i> -Alkyl perfluoroalkane sulfonamidoethyl acrylates and methacrylates (MeFAS(M)ACs, EtFAS(M)ACs, BuFAS(M)ACs)	$-SO_2N(R')CH_2CH_2O-C(O)CH=CH_2$ and $-SO_2N(R')CH_2CH_2O-C(O)C(CH_3)=CH_2$ where $R' = C_mH_{2m+1}$ ($m = 1, 2, 4$)	<i>N</i> -Ethyl perfluorooctane sulfonamidoethyl acrylate (EtFOSAC), $C_8F_{17}SO_2N(C_2H_5)CH_2CH_2O-C(O)CH=CH_2$	Major raw material for surfactant and surface protection products
	Perfluoroalkane sulfonamidoacetic acids (FASAAAs) and <i>N</i> -alkyl perfluoroalkane sulfonamidoacetic acids (MeFASAAAs, EtFASAAAs, BuFASAAAs)	$-SO_2N(R')CH_2COOH$ where $R' = C_mH_{2m+1}$ ($m = 0, 1, 2, 4$)	<i>N</i> -Ethyl perfluorooctane sulfonamidoacetic acid (EtFOSAAA), $C_8F_{17}SO_2N(C_2H_5)CH_2CO_2H$	Intermediate environmental transformation product
Fluorotelomer substances ^b	Semifluorinated <i>n</i> -alkanes (SFAs) and alkenes (SFAenes)	$-(CH_2)_mH$ and $-CH=CH(CH_2)_{m-2}H$, with $m = 2-16$ and $n = 6-16$	Perfluorohexylhexadecane (F_6H_{16}), $F(CF_2)_6(CH_2)_{16}H$	Ski wax; medical applications
	<i>n</i> :2 Fluorotelomer iodides (<i>n</i> :2 FTIs) (Telomer B)	$-CH_2CH_2I$	8:2 Fluorotelomer iodide (8:2 FTI), $C_8F_{17}CH_2CH_2I$	Major raw material for surfactant and surface protection products
	<i>n</i> :2 Fluorotelomer olefins (<i>n</i> :2 FTOs)	$-CH=CH_2$	6:2 Fluorotelomer olefin (6:2 FTO), $C_6F_{13}CH=CH_2$	Raw material for surfactant and surface protection products
	<i>n</i> :2 Fluorotelomer alcohols (<i>n</i> :2 FTOHs)	$-CH_2CH_2OH$	10:2 Fluorotelomer alcohol (10:2 FTOH), $C_{10}F_{21}CH_2CH_2OH$	Major raw material for surfactant and surface protection products
	<i>n</i> :2 Unsaturated fluorotelomer alcohols (<i>n</i> :2 FTUOHs)	$-CF=CHCH_2OH$	8:2 Unsaturated fluorotelomer alcohol (8:2 FTUOH), $C_7F_{15}CF=CHCH_2OH$	Intermediate environmental transformation product
	<i>n</i> :2 Fluorotelomer acrylates (<i>n</i> :2 FTACs) and methacrylates (<i>n</i> :2 FTMACs)	$-CH_2CH_2OC(O)CH=CH_2$ and $-CH_2CH_2OC(O)C(CH_3)=CH_2$	8:2 Fluorotelomer acrylate (8:2 FTAC), $C_8F_{17}CH_2CH_2OC(O)CH=CH_2$ 6:2 Fluorotelomer methacrylate (6:2 FTMAC), $C_6F_{13}CH_2CH_2O-C(O)C(CH_3)=CH_2$	Major raw material for fluorotelomer-based polymers used in surface protection products
	<i>n</i> :2 Polyfluoroalkyl phosphoric acid esters, polyfluoroalkyl phosphates, fluorotelomer phosphates (PAPs)	$-(CH_2CH_2O)_xP(=O)(OH)_{3-x}$ where $x = 1$ or 2	8:2 Fluorotelomer phosphate monoester (8:2 monoPAP), $C_8F_{17}CH_2CH_2OP(=O)(OH)_2$ 8:2 Fluorotelomer phosphate diester (8:2 diPAP), $(C_8F_{17}CH_2CH_2O)_2P(=O)OH$	Surfactant and surface protection products

(Continued)

Table 3. (Continued)

	Classification and chemical structure	$C_nF_{2n+1}R$, where R =	Examples	Uses
	n:2 Fluorotelomer aldehydes (n:2 FTALs) and unsaturated aldehydes (n:2 FTUALs)	$-CH_2CHO$ and $-CF=CHCHO$	8:2 Fluorotelomer aldehyde (8:2 FTAL), $C_8F_{17}CH_2CHO$ 8:2 Fluorotelomer unsaturated aldehyde (8:2 FTUAL), $C_7F_{15}CF=CHCHO$	Intermediate environmental transformation product
	n:2 Fluorotelomer carboxylic acids (n:2 FTCAs) and unsaturated carboxylic acids (n:2 FTUCAs)	$-CH_2COOH$ and $-CF=CHCOOH$	8:2 Fluorotelomer carboxylic acid (8:2 FTCA), $C_8F_{17}CH_2COOH$ 8:2 Fluorotelomer unsaturated carboxylic acid (8:2 FTUCA), $C_7F_{15}CF=CHCOOH$	Intermediate environmental transformation product
	n:3 Saturated acids (n:3 Acids) and n:3 Unsaturated acids (n:3 UAcids)	$-CH_2CH_2COOH$ and $-CH=CHCOOH$	7:3 Acid, $C_7F_{15}CH_2CH_2COOH$ 7:3 UAcid, $C_7F_{15}CH=CHCOOH$	Intermediate environmental transformation product
	n:2 Fluorotelomer sulfonic acids (n:2 FTSAs)	$-CH_2CH_2SO_3H$	8:2 Fluorotelomer sulfonic acid (8:2 FTSa), $C_8F_{17}CH_2CH_2SO_3H$	Surfactant and environmental transformation product
Miscellaneous	Polyfluoroalkyl ether carboxylic acids	For example: $-O(C_mF_{2m})OCHF(C_pF_{2p})COOH$	4,8-Dioxa-3H-perfluorononanoate, $CF_3OCF_2CF_2CF_2OCHFCF_2COOH$	Alternative fluoropolymer processing aid (as ammonium salt)

^aSubstances originating by electrochemical fluorination (ECF) process;

^bSubstances originating by fluorotelomer process.

In addition to their major commercial use as fluoropolymer processing aids and numerous industrial and consumer applications (Kissa 2001; Prevedouros et al. 2006), PFCAs are also the terminal degradation products from abiotic and biotic degradation of certain precursor PFASs. Such precursors include fluorotelomer alcohols (Hagen et al. 1981; Dinglasan et al. 2004; Ellis et al. 2004; Hurley et al. 2004; Wang et al. 2009; Liu et al. 2010), fluorotelomer acrylates (Butt et al. 2009; Butt et al. 2010b), fluorotelomer iodides (Young et al. 2008), fluorotelomer olefins (Nakayama et al. 2007), N-alkyl perfluoroalkane sulfonamides (Tomy, Tittlemier, et al. 2004; Martin et al. 2006; Plumlee et al. 2009), N-alkyl perfluoroalkane sulfonamidoethanols (D'eon et al. 2006; Plumlee et al. 2009), and polyfluoroalkyl phosphates (D'eon and Mabury 2007; Lee et al. 2010). Short-chain PFCAs (e.g., trifluoroacetic and pentafluoropropionic acids) may also be formed in the atmospheric degradation of certain hydrochlorofluorocarbons, hydrofluorocarbons, and fluorinated anesthetics (Boutonnet et al. 1999; Young and Mabury 2010) and perfluoro-2-methyl-3-pentanone (Jackson et al. 2011), as well as in the oxidative thermolysis of fluorinated polymers (Ellis et al. 2001). Yet, the quantitative attribution of sources of these short-chain PFCAs in the environment remains uncertain, and it is quite possible that further precursors will be identified. PFCA yields and rates of formation vary depending on the precursor substance and degradation conditions. Moreover, PFCAs and potential PFCA precursors, such as residual raw materials, may be present as impurities in commercial PFAS-based products (Washburn et al. 2005; Berger and Herzke 2006; Dinglasan-Panlilio and Mabury 2006; Larsen et al. 2006; Prevedouros et al. 2006; Schulze and Norin 2006; D'eon and Mabury 2007; Jensen et al. 2008; Fiedler et al. 2010). It was estimated that the majority (~80%) of PFCAs have been released to the

environment from fluoropolymer manufacture and use (Prevedouros et al. 2006). This percentage is, however, an overall value, heavily weighted toward the PFCAs with the greatest emissions, namely PFOA and (to a much lesser extent) PFNA. PFCAs with shorter or longer chain lengths are not known to arise primarily from fluoropolymer manufacture and use. Although in the same study (Prevedouros et al. 2006), indirect sources of PFOA and PFNA were estimated to be much less important than direct sources, there were larger uncertainties associated with the calculations for indirect sources and some recently identified precursors (e.g., polyfluoroalkyl phosphates) were excluded.

In 2006, 8 major global companies signed on to the USEPA "2010/2015 PFOA Stewardship Program" (USEPA 2006b) with commitments first to reduce emissions and product content of PFOA, higher homologues and precursors by 95% by 2010 and second to work toward the elimination of PFOA, higher homologues, and precursors by 2015. Companies have reported significant progress toward achieving these goals (Ritter 2010). Interestingly, coincident with these changes, there have been reports showing significantly increased levels of perfluorobutanoic acid (PFBA) in water (Möller et al. 2010) and air (Weinberg et al. 2011b) that are most likely associated with the conversion to shorter chain perfluoroalkyl products.

Perfluoroalkane (or -alkyl) sulfonic acids: Perfluoroalkyl sulfonic acids, $C_nF_{2n+1}SO_3H$ (PFSAs, Table 2), are the 2nd major PFAA family of significance. The alternative name perfluoroalkane sulfonic acid has been used most commonly in the literature, in line with IUPAC recommendations, and we will adopt it here. Perfluorooctane sulfonic acid $C_8F_{17}SO_3H$ (PFOS), is the PFSA that has commanded greatest attention beginning when it was first detected

Table 4. Hierarchical overview of fluoropolymers, perfluoropolyethers, and side-chain-fluorinated polymers

		Example(s)	Uses
Fluoropolymers: Carbon-only polymer backbone with F directly attached to backbone C atoms		$-(CF_2CF_2)_n-$ Polytetrafluoroethylene (PTFE) $-(CH_2CF_2)_n-$ Polyvinylidene fluoride (PVDF) $-(CH_2CHF)_n-$ Polyvinyl fluoride (PVF) $-(CF_2CF_2)_n-(CF(CF_3)CF_2)_m-$ Fluorinated ethylene propylene (FEP)	Plastics
Perfluoropolyethers (PFPEs): Ether polymer backbone with F atoms directly attached		Examples: $F-(C_mF_{2m}O)_nCF_3$ $HOCH_2O-[C_mF_{2m}O]_nCH_2OH$ -where $C_mF_{2m}O$ represents $-CF_2O-$, $-CF_2CF_2O-$, and/or $-CF(CF_3)CF_2O-$ units distributed randomly along the polymer backbone	Functional fluids, surfactants, and surface protection products
Side-chain-fluorinated polymers: Nonfluorinated polymer backbone with fluorinated side chains, ending in $-C_nF_{2n+1}$	Fluorinated acrylate and methacrylate polymers	Acrylate: Backbone-CH-C(O)O-X- C_nF_{2n+1} Methacrylate: Backbone-C(CH ₃)-C(O)O-X- C_nF_{2n+1} -where X is $-CH_2CH_2N(R')SO_2-$ with $R' = -C_nH_{2n+1}$ ($n = 0, 1, 2, 4$) or $-CH_2CH_2-$	Surfactants and surface protection products
	Fluorinated urethane polymers	Backbone-NHC(O)O-X- C_nF_{2n+1} -where X is either $-CH_2CH_2N(R')SO_2-$ with $R' = -C_nH_{2n+1}$ ($n = 0, 1, 2, 4$) or $-CH_2CH_2-$	Surfactants and surface protection products
	Fluorinated oxetane polymers	Backbone-CH ₂ OCH ₂ -R -where R = $-CF_3$, $-C_2F_5$ or $-CH_2C_4F_9$	Surfactants and surface protection products

globally in biota (Giesy and Kannan 2001) and humans (Hansen et al. 2001). Subsequently, as stated above, the production of PFOS, perfluorohexane sulfonic acid (PFHxS), perfluorodecane sulfonic acid (PFDS), and the precursors of these PFSA, was phased out by the major manufacturer in 2002 (3M Company 2000c; USEPA 2000). Nevertheless, PFOS and its derivatives are still manufactured in China (Han 2009), with a production of more than 200 tons of its precursor, perfluorooctane sulfonyl fluoride, in 2006 (Yue 2008). PFOS and related compounds have been the subject of a European Union directive restricting their production and use (European Parliament 2006b). Furthermore, PFOS has been classified as a persistent, bioaccumulative, and toxic substance (OECD 2002) and was recently added to Annex B (requiring use restrictions) of the Stockholm Convention list of persistent organic pollutants (UNEP 2009). Formerly, PFOS had a number of industrial and commercial applications (3M Company 1999; Kissa 2001; Brooke et al. 2004; Paul et al. 2009). However, the environmental and toxicological significance of PFOS, ubiquitous in the global environment, also results from its presence as an impurity in and formation from perfluorooctane sulfonamido precursor substances (3M Company 1999, 2000a; Lange 2000, 2001; Xu et al. 2004; Boulanger et al. 2005; D'eon et al. 2006; Rhoads et al. 2008; Xie et al. 2009) used in vastly greater quantities (Brooke et al. 2004; Paul et al. 2009). The global commercial production of PFOS and related compounds has, to our knowledge, been based essentially or perhaps exclusively on ECF. In this

process, the electrolysis of a solution of octane sulfonyl fluoride in anhydrous HF leads to perfluorooctane sulfonyl fluoride, $C_8F_{17}SO_2F$ (POSF), the key intermediate from which all PFOS-related products are subsequently produced (3M Company 1999; Brooke et al. 2004; Lehmler 2005; Paul et al. 2009). The resulting PFOS, the precursor POSF and other derivatives manufactured from it, e.g., perfluorooctane sulfonamido derivatives such as amides, ethanol-substituted amides, and surfactant and polymeric products therefrom, may contain up to 30% branched isomers (Reagen et al. 2007), as well as additional C chain length homologues. For example, samples of the K salt of PFOS taken from the same 3M commercial lot were analyzed by 2 laboratories and found to have a purity of only 85% to 87% (representing the sum of all K-PFOS isomers), on the account of the presence mainly of C_2 - C_{10} PFSA homologues, but also of a range of PFCAs and other impurities (Seacat et al. 2003; Arsenault et al. 2008). Shorter perfluoroalkyl chain length products, notably perfluorobutane sulfonyl-based products, have been introduced as alternatives to the previously used compounds with 6 or more perfluorinated carbons, because these shorter chain length substances do not bioaccumulate due to their rapid elimination in multiple organisms tested (Olsen et al. 2009). This substitution is a consequence of the voluntary phase-out and/or subsequent regulatory restriction of PFOS-related substances and certain homologues with 5 to 7 and 9 or 10 perfluorinated C atoms (3M Company 2000b; Federal Register 2006b). Coincident with these changes, reports have

shown significantly increased levels of perfluorobutane sulfonic acid (PFBS) in environmental waters, no doubt as a consequence of the conversion to 4-C ECF-derived perfluorobutane sulfonyl products (Eschauzier et al. 2010; Möller et al. 2010).

Perfluoroalkane (or -alkyl) sulfinic acids: Perfluoroalkane sulfinic acids, $C_nF_{2n+1}SO_2H$ (PFSIAs; Table 2), are degradation products from commercial precursor compounds containing the $C_nF_{2n+1}SO_2N<$ moiety (e.g., perfluoroalkane sulfonamido ethanols, $C_nF_{2n+1}SO_2N(R)CH_2CH_2OH$) (Lange 2000, 2001; Boulanger et al. 2005; Rhoads et al. 2008). PFSIAs have been detected in wastewater treatment plant (WWTP) effluents and in the environment (Ahrens et al. 2009b; Ahrens, Siebert, et al. 2009; Ahrens, Xie, et al. 2010).

Perfluoroalkyl phosphonic and phosphinic acids: Perfluoroalkyl phosphonic acids, $O=P(OH)_2C_nF_{2n+1}$ (PFPAs; Table 2), and perfluoroalkyl phosphinic acids, $O=P(OH)(C_nF_{2n+1})(C_mF_{2m+1})$ (PFPIAs; Table 2), are commercial surfactants manufactured and offered for a range of consumer and industrial uses (USEPA 2006a; Mason Chemical 2011). Blends of C_6 – C_{12} PFPAs and similar PFPIA blends, with CAS numbers 68412-68-0 and 68412-69-1, respectively, have been reported to have had annual production volumes in the range of tonnes to hundreds of tonnes in 1998 and 2002 (Howard and Muir 2010), but only recently have PFPAs been widely detected in environmental waters (D'eon et al. 2009b; D'eon and Mabury 2010) and PFPIAs in WWTP sludge (D'eon and Mabury 2010) and human serum (Lee and Mabury 2011).

Fluorotelomer-based products. The term “fluorotelomer-based products” describes a family of raw material building blocks, surfactant and polymeric products, and degradation products that all originate from the starting fluorotelomer raw material, perfluoroalkyl iodides (PFAIs), as depicted in Figures 2 and 3. As reviewed below, the degradation of fluorotelomer-based products is a potential source of PFCAs in the environment.

Perfluoroalkyl iodides, fluorotelomer iodides, and fluorotelomer olefins: Perfluoroalkyl iodides, $C_nF_{2n+1}I$ (PFAIs; Table 2), and $n:2$ fluorotelomer iodides, $C_nF_{2n+1}CH_2CH_2I$ ($n:2$ FTIs; Table 3), are the first 2 raw materials that lead to the family of polyfluoroalkyl “fluorotelomer-based” products. Both PFAIs and $n:2$ FTIs have recently been detected in air and soil near a fluorotelomer manufacturing facility in China (Ruan et al. 2010). Fluorotelomer olefins, $C_nF_{2n+1}CH=CH_2$ (FTOs; Table 3) are synthesized by dehydrohalogenation of FTIs and may also be formed as an impurity in synthesizing fluorotelomer alcohols (FTOHs) from FTIs (Prevedouros et al. 2006). As stated above, processes for manufacturing PFNA by oxidation of 8:2 FTO have been patented and may have been used industrially. FTOs are hydrosilylated to create silanes that are used in a number of applications. FTOs have been detected in the atmosphere (Barber et al. 2007; Jahnke et al. 2007; Piekartz et al. 2007), where they degrade completely and rapidly, but are expected to form low yields of PFCAs (Young and Mabury 2010). The degradation scheme proceeds via a $C_nF_{2n+1}CHO$ perfluoroalkyl aldehyde (PFAL; Table 2) intermediate (Vésine et al. 2000; Nakayama

et al. 2007). The atmospheric transformation of FTIs probably is comparable to FTOs in the ultimate outcome, mineralization with low yield of PFCAs (typically 1%–10%), and involves both fluorotelomer aldehyde $C_nF_{2n+1}CH_2CHO$ (FTAL; Table 3) and PFAL intermediates, together with the fluorotelomer carboxylic acids, $C_nF_{2n+1}CH_2COOH$ (FTCAs; Table 3) (Young et al. 2008). FTIs may hydrolyze in natural waters (Rayne and Forest 2010c), and this transformation process would presumably lead to fluorotelomer alcohols and, hence, their degradation products, as discussed below.

Fluorotelomer alcohols and their acrylic, methacrylic, and phosphoric esters: The $n:2$ fluorotelomer alcohols, $C_nF_{2n+1}CH_2CH_2OH$ ($n:2$ FTOHs; Table 3), are key raw materials in the production of $n:2$ fluorotelomer acrylates, $C_nF_{2n+1}CH_2CH_2OC(O)CH=CH_2$ ($n:2$ FTACs) and $n:2$ fluorotelomer methacrylates, $C_nF_{2n+1}CH_2CH_2OC(O)C(CH_3)=CH_2$ ($n:2$ FTMACs) (Table 3 and Figure 3). The FT(M)AC monomers are copolymerized in an aqueous emulsion polymerization with a host of non-fluorinated acrylates and other monomers to manufacture fluorotelomer-based polymers (Rao and Baker 1994). These polymers provide water, oil, and stain repellency to textiles, leather, and paper substrates. There is extensive scientific literature on the environmental occurrence of FTOHs, particularly (but not exclusively) in air (Martin et al. 2002; Oono, Harada, et al. 2008; Oono, Matsubara, et al. 2008; Strynar and Lindstrom 2008; Jahnke et al. 2009; Mahmoud et al. 2009; Dreyer et al. 2010; Langer et al. 2010; Shoeib et al. 2010; Yoo et al. 2010; Ahrens et al. 2011; Haug et al. 2011; Shoeib et al. 2011; Yoo et al. 2011). Likewise, some FTACs (Piekartz et al. 2007; Oono, Harada, et al. 2008; Oono, Matsubara, et al. 2008; Dreyer, Weinberg, et al. 2009; Mahmoud et al. 2009; Dreyer et al. 2010; Langer et al. 2010; Weinberg et al. 2011a, 2011b) and FTMACs (Oono, Matsubara, et al. 2008) have also been detected in environmental samples. The chain lengths of these fluorotelomer derivatives may vary over a broad range. For example, FTOHs with up to 18 fluorinated C atoms have been reported as detected, but not quantified, in air from an occupational setting (Nilsson et al. 2010).

Fluorotelomer alcohol phosphate esters (Table 3) are commercial fluorinated surfactants that are made by many global suppliers by the same reactions employed for non-fluorinated phosphates and used primarily for their surface tension lowering, wetting, and leveling surfactant properties (Taylor 1999). The terminology we recommend for these substances is polyfluoroalkyl phosphoric acid monoesters (monoPAPs), $(O)P(OH)_2(OCH_2CH_2C_nF_{2n+1})$, and diesters (diPAPs), $(O)P(OH)(OCH_2CH_2C_nF_{2n+1})(OCH_2CH_2C_mF_{2m+1})$. They may also be called $n:2$ fluorotelomer monophosphates and diphosphates. These compounds have been used as grease-proofing agents for food-contact paper (D'eon and Mabury 2007; Begley et al. 2008; FDA 2009; Lee et al. 2010; Lee and Mabury 2011), often as blends of varying perfluoroalkyl chain length and as salts (e.g., of diethanolamine). One specific use of monoPAPs and diPAPs that has led to their widespread presence in the environment is as an approved defoaming adjuvant in pesticide formulations. Approval for this use has now been rescinded (Federal Register 2006a). Recently, diPAPs have been reported detected in human serum at concentrations in some cases comparable to those of PFOA and in WWTP sludge at much greater levels than PFOA (D'eon et al. 2009a; Lee and Mabury 2011).

Semifluorinated alkanes and alkenes: Diblock semifluorinated n -alkanes (SFAs), $F(CF_2)_n(CH_2)_mH$ (or, briefly, F_nH_m ; Table 3), are a class of chemicals that are manufactured with a wide variety of chain lengths, depending on the intended use, by adding an olefin to a perfluoroalkyl iodide followed by reductive dehalogenation (Napoli 1996). These reactions also lead to semifluorinated n -alkenes (SFAenes), $F(CF_2)_nCH=CH(CH_2)_{m-2}H$ (or, briefly, F_nH_{m-1} ene), as byproducts (Coe and Milner 1972). Since the 1990s, industrial mixtures of long-chain SFAs (≥ 22 C atoms) have been applied in ski waxes, because they reduce friction and repel dirt due to their extremely low surface tension (Rogowski et al. 2007). Shorter-chain SFAs are used in medicinal applications (e.g., Kirchhof et al. 2002). In fluorinated ski waxes, up to 15% of SFAs are mixed with normal paraffins. The presence of SFAs in snow and soil samples from a ski area in Sweden has recently been demonstrated (Plassmann and Berger 2010).

Degradation products of fluorotelomer alcohols and their esters: Fluorotelomer aldehydes and acids, perfluoroalkyl aldehydes, perfluoroalkyl carboxylic acids, and so forth: The aerobic biodegradation and metabolic degradation pathways for fluorotelomer alcohols have been well studied (Frömel and Knepper 2010). A general overview of the 8:2 FTOH aerobic biodegradation pathways is presented in Figure 5. The pathways and yields of transformation products depend on the matrix in which the environmental microbial degradation (e.g., sludge, soil) or metabolism (rat, mouse, in vivo, in vitro) takes place and the length of the perfluoroalkyl chain in the fluorotelomer alcohol (Hagen et al. 1981; Dinglasan et al. 2004; Martin et al. 2005; Wang et al. 2009; Butt et al. 2010a; Liu et al. 2010; Brandsma et al. 2011). In general, the first step in biodegradation is aerobic oxidation of the starting n :2 fluorotelomer alcohol to form the corresponding n :2 fluorotelomer aldehyde, $C_nF_{2n+1}CH_2CHO$ (n :2 FTAL; Table 3), a short-lived, highly reactive species. The aldehyde is rapidly oxidized to form the corresponding n :2 fluorotelomer carboxylic acid, $C_nF_{2n+1}CH_2COOH$ (n :2 FTCA; Table 3). Next, dehydrohalogenation of the acid occurs to form the corresponding n :2 unsaturated carboxylic acid, $C_{n-1}F_{2n-1}CF=CHCOOH$ (n :2 FTUCA; Table 3). The dehydrohalogenation of the starting n :2 fluorotelomer alcohol to form the n :2 unsaturated fluorotelomer alcohol, $C_{n-1}F_{2n-1}CF=CHCH_2OH$ (Table 3), and oxidation to yield the n :2 unsaturated fluorotelomer aldehyde, $C_{n-1}F_{2n-1}CF=CHCHO$ (n :2 FTUAL; Table 3), have also been observed. Thereafter, a host of transient and stable transformation products, including PFCAs, have been identified. A unique transformation product identified is a polyfluorinated carboxylic acid with the same number of total C atoms as the parent n :2 FTOH where the 2 F atoms of the $-CF_2-$ group directly adjacent to the $-CH_2CH_2-$ moiety have been replaced with H atoms, $C_{n-1}F_{2n-1}CH_2CH_2COOH$, and a corresponding unsaturated acid, $C_{n-1}F_{2n-1}CH=CHCOOH$ (Table 3) (Martin et al. 2005; Wang et al. 2005; Fasano et al. 2006; Wang et al. 2009; Butt et al. 2010a). For these substances, we suggest for simplicity that either the formal name of the acid be used or the simple acronyms x :3 Acid and x :3 UAcid, where the x ($= n - 1$) designates the number of perfluorinated carbons and “3” the number of nonfluorinated C atoms. For the remaining transformation products, we suggest adopting the naming given to these substances by the authors (e.g., Martin et al. 2005; Wang et al. 2009; Butt et al. 2010a;

Liu et al. 2010). In a sediment–water microcosm, the degradation products observed from n :2 FTCA substrates were the corresponding PFCAs, whereas n :2 FTUCAs also led to $(n - 1)$:3 Acids (Myers and Mabury 2010).

In mammals, the metabolic pathways for 8:2 and 6:2 FTOHs have been well studied in vivo in rats and mice and in vitro in rats, mice, and human hepatocytes. In general, the majority of administered FTOH test substance was eliminated rapidly in urine as conjugates. Absorption, distribution, metabolism, and elimination (ADME) studies using [^{14}C]-radiolabeled FTOHs have been conducted. The characteristic degradation products observed in microbial studies, including PFCAs, as well as some of their conjugates, have been reported in urine and at trace levels in organs and tissues (Fasano et al. 2006; Nabb et al. 2007; Fasano et al. 2009). The reader is referred to the articles for greater detail on these studies.

In atmospheric degradation studies, reviewed by Young and Mabury (2010), it has been shown that oxidation of n :2 FTOHs also leads to the formation of n :2 FTALs, n :2 FTCAs, and perfluoroalkyl aldehydes, $C_nF_{2n+1}CHO$ (PFALs; Table 2). Low yields (typically 1%–5%) of PFCAs having the same number of perfluorinated C atoms as the parent FTOH, or fewer, down to CF_3COOH , may be expected in low- NO_x atmospheres. The PFCAs with $n - 2$ or fewer perfluorinated C atoms result from “unzipping” of the perfluoroalkyl chain, by splitting off of $C(O)F_2$ molecules from the intermediate perfluoroalkoxy radicals (Ellis et al. 2004). Nevertheless, complete mineralization to $C(O)F_2$ is the major atmospheric outcome, and the yields of PFCAs decline as atmospheric NO_x levels increase (Ellis et al. 2004; Wallington et al. 2006; Young and Mabury 2010). A simplified scheme, given in Figure 6, shows the key intermediates in the atmospheric degradation of n :2 FTOHs to the products mentioned above, illustrated for $n = 8$. This scheme also includes the atmospheric breakdown pathways for FTIs and FTOs, discussed above, as well as for FTACs (Butt et al. 2009), because all these fluorotelomer derivatives have part of their degradation mechanism in common. This is also likely to be the case for PFALs (Figure 6), assuming they photolyze easily to perfluoroalkyl radicals (which add O_2 to give perfluoroalkylperoxy radicals) in the lower atmosphere, as has been demonstrated for CF_3I (Solomon et al. 1994).

It is worth noting here that the PFALs will probably exist in cloud and surface waters largely as their *gem*-diol hydrates, $C_nF_{2n+1}CH(OH)_2$ (PFAL- H_2O s; Table 2), unlike the FTALs for which the hydration equilibrium is much less favorable (Rayne and Forest 2010b). With estimated pK_a values of 9 or higher, the PFAL- H_2O s will not be ionized to any great extent under environmental conditions, whereas the corresponding hydrates formed from FTALs are even weaker acids ($pK_a > 12$) (Rayne and Forest 2010b).

The esters of FTOHs may hydrolyze abiotically or biotically to FTOHs and, hence, ultimately lead to the same range of fluorinated transformation products described above. Hydrolysis studies of mono- and polyesters and mono-urethanes containing a fluorotelomer moiety have recently been reported (Dasu et al. 2010). Moreover, as expected, characteristic FTOH degradation products were detected when rainbow trout were exposed to 8:2 FTAC through their diet (Butt et al. 2010b), and when rats were dosed with monoPAPs or diPAPs (D’eon and Mabury 2007, 2011). Both




Figure 5. Aerobic biotransformation pathways for 8:2 fluorotelomer alcohol (8:2 FTOH). Adapted from Wang et al. (2009).

FTOHs and their transformation products were observed in experiments intended to simulate aerobic biodegradation of monoPAPs and diPAPs in WWTPs (Lee et al. 2010). The abiotic hydrolysis of FTACs has been predicted to have half-lives of years in marine systems but possibly only days in landfills (Rayne and Forest 2010c). Hydrolytic stability studies, conducted under OECD 111 Guidelines, on a commercial fluorotelomer-based acrylate polymer (Russell et al. 2008) and a urethane polymer (Russell et al. 2010) showed no discernible hydrolysis. Nevertheless, there is much debate regarding the hydrolysis and biodegradation of commercial fluorotelomer-based polymers (Russell et al. 2008; Koch et al. 2009; Russell et al. 2009; Washington et al. 2009a; Washington et al. 2009b) that future research will illuminate.

A number of reported observations of *n*:2 FTACs and/or *n*:2 FTUCAs have occurred in environmental media and biota such as atmospheric particles (Stock et al. 2007), indoor dust (Barber et al. 2007), precipitation (Loewen et al. 2005; Scott et al. 2006; Taniyasu et al. 2008; Kwok et al. 2010; Scott et al. 2010), surface waters (Stock et al. 2007; Ahrens et al., 2009a; Scott et al. 2010; Zushi et al. 2011), sediments (Stock et al. 2007), WWTP effluent (Sinclair and Kannan 2006; Zushi et al. 2011), sewage sludge (Zhang et al. 2010), landfill leachate (Huset et al. 2011), animal biota (Houde et al. 2005; Taniyasu et al. 2005; Butt, Mabury, et al. 2007; Butt, Muir, et al. 2007; Furdui et al. 2007; Gebbink et al. 2009), human breast milk (So et al. 2006), and foodstuffs (Ostertag et al. 2009). The 7:3 Acid has also been detected in biota (Powley et al. 2008; Peng et al. 2010; Guruge et al. 2011). The various perfluorinated and polyfluorinated aldehydes discussed above

have apparently not yet been found in environmental samples. This is most likely due to their highly reactive nature, because only trapping experiments have qualified their presence thus far in laboratory studies.

Fluorotelomer sulfonic acids: The *n*:2 fluorotelomer sulfonic acids, $C_nF_{2n+1}CH_2CH_2SO_3H$ (FTSAs; Table 3) have been found in groundwater, soil, and biota, especially at military bases, firefighting training sites, and locations where major fires have been extinguished through use of AFFFs (Schultz et al. 2004; Norwegian Pollution Control Authority 2008; Oakes et al. 2010). They have also been detected in WWTP effluents (Huset et al. 2008; Ahrens et al., 2009b), landfill leachate (Eggen et al. 2010; Huset et al. 2011), precipitation and fresh surface waters (Kim and Kannan 2007; Scott et al. 2010; Nguyen et al. 2011), seawater contaminated by AFFFs (Taniyasu et al. 2005), sediments (Zushi et al. 2010), Arctic biota (Miljeteig et al. 2009), and human serum (Lee and Mabury 2011). These FTSAs arise from the degradation of more complex fluorotelomer-based substances containing the $C_nF_{2n+1}CH_2CH_2S-R$ or $C_nF_{2n+1}CH_2CH_2SO_2-R$ moiety (where *R* is a hydrophilic functional group that provides surfactant properties). These precursor compounds may be used as components of firefighting foams (Bertocchio and Foulletier 1970; Falk 1982; Schultz et al. 2004), e.g., the betaine $F(CF_2)_nCH_2CH_2SO_2NHCH_2CH_2N^+(CH_3)_2CH_2CH_2CO_2^-$, or in food packaging applications, e.g., the fluorotelomer mercaptoalkyl phosphate esters (Lee and Mabury 2011; Trier, Granby, et al. 2011; Trier, Nielsen, et al. 2011). FTSAs have been shown to undergo slow aerobic biotransformation to form trace levels of PFCAs (Wang et al.




Figure 6. Simplified atmospheric degradation scheme for 8:2 fluorotelomer derivatives. Free-radical and transient molecular intermediates are shown in boxes with a dashed outline, while the starting compounds, the more stable molecular intermediates, and the final products are shown in boxes with a solid outline, their acronyms being indicated in bold type. An arrow on the chart often implies several elementary steps: i.e., certain intermediates are omitted.

2011). It should be noted that 6:2 FTSA has been referred to in some literature as “tetrahydro PFOS.” Because 6:2 FTSA is both chemically and biologically very different from PFOS (Wang et al. 2011), we strongly discourage this usage and recommend 6:2 FTSA be used in naming this substance.

Perfluoroalkane sulfonamido derivatives: Perfluoroalkane sulfonamides, sulfonamidoethanols, sulfonamidoethyl acrylates, and sulfonamidoethyl methacrylates. In the same way as the perfluoroalkyl iodides and fluorotelomer iodides are important building blocks for a broad range of fluorotelomer derivatives, the perfluoroalkane sulfonyl fluorides, $C_nF_{2n+1}SO_2F$ (PASFs; Table 2) play an analogous role as precursors in the manufacture not only of the PFSA already discussed, but also of a variety of compounds containing

the perfluoroalkane sulfonamido group, $C_nF_{2n+1}SO_2N<$ (Tables 2 and 3). This is illustrated in Figure 7 for the synthesis of several families of perfluoroalkane sulfonamido derivatives, exemplified for a starting PASF with 8 C atoms. PFSA were directly manufactured by hydrolysis of PASFs and the various salt forms (ammonium, diethanolamine, and K and Li salts) were manufactured by neutralization of the acids. The greater part of the production of PASFs (notably POSF), however, was used to produce fluorinated surfactants and high-molecular-weight fluorinated polymeric products (3M Company 1999). The major pathway for conversion of PASFs into commercial derivatives involves reacting them in a first step with a primary amine, generally methylamine or ethylamine, to give *N*-methyl or *N*-ethyl perfluoroalkane sulfonamides, $C_nF_{2n+1}SO_2NH(C_mH_{2m+1})$, where $m = 1$ or 2

(MeFASAs and EtFASAs; Table 3) (3M Company 1999; Lehmler 2005). These *N*-alkyl FASAs are, in some cases, commercial products in their own right, as well as building blocks for further synthesis. For instance, *N*-ethyl perfluorooctane sulfonamide, $C_8F_{17}SO_2NH(C_2H_5)$, or EtFOSEA, is the pesticide sulfluramid. In a 2nd major industrial reaction step, *N*-alkyl FASAs are reacted with ethylene carbonate to give another series of building blocks, the *N*-methyl or *N*-ethyl perfluoroalkane sulfonamido ethanols, $C_nF_{2n+1}SO_2N(C_mH_{2m+1})CH_2CH_2OH$, where $m=1$ or 2 (MeFASEs and EtFASEs; Table 3) (3M Company 1999; Lehmler 2005). These *N*-alkyl FASEs are analogous to FTOHs. Because they are alcohols, they can be converted into acrylates and methacrylates, as well as into phosphates and other derivatives (3M Company 1999) that will not be discussed further here. The *N*-alkyl perfluoroalkane sulfonamidoethyl acrylates, $C_nF_{2n+1}SO_2N(C_mH_{2m+1})CH_2CH_2OC(O)CH=CH_2$, where $m=1$ or 2 (MeFASACs and EtFASACs; Table 3) and the corresponding *N*-alkyl perfluoroalkane sulfonamidoethyl methacrylates, $C_nF_{2n+1}SO_2N(C_mH_{2m+1})CH_2CH_2OC(O)C(CH_3)=CH_2$ (MeFASMACs and EtFASMACs; Table 3) are used in a similar manner to the fluorotelomer acrylates and methacrylates, as comonomers for synthesizing acrylic polymers used in surface protection applications (3M Company 1999).

The (alkyl)-FASA, FASE, FASAC, and FASMAC intermediates were the principal building blocks of many fluorochemical products used in surface treatments, paper packaging protectors, and other specialist applications. A more extensive range of commercial compounds has, however, been produced, as described in industry reports (3M Company 1999). In 2002, the largest historic manufacturer of perfluorooctane sulfonyl derivatives ($n \geq 6$) ceased manufacture (3M Company 2000a; USEPA 2000) and has since introduced products based on perfluorobutane sulfonyl

chemistry (Renner 2006; Ritter 2010). Meanwhile, existing and new manufacturers continue to make PFOS and other long-chain perfluoroalkane sulfonates and products derived from them.

Degradation products of perfluoroalkane sulfonamido derivatives: Published studies on the aerobic biotransformation of the perfluoroalkane sulfonamido derivatives focus on those compounds having 8 perfluorinated C atoms, in particular *N*-ethyl perfluorooctane sulfonamidoethanol (EtFOSE), which is ultimately degraded to PFOS. Various intermediates leading to this perfluoroalkane sulfonic acid have been reported, including the members of the following families (Tables 2 and 3) with $n=8$: *N*-ethyl perfluoroalkane sulfonamidoacetic acids (EtFASAAAs), $C_nF_{2n+1}SO_2N(C_2H_5)CH_2COOH$; *N*-ethyl perfluoroalkane sulfonamides (EtFASAs), $C_nF_{2n+1}SO_2NH(C_2H_5)$; perfluoroalkane sulfonamidoacetic acids (FASAAAs), $C_nF_{2n+1}SO_2NHCH_2COOH$; perfluoroalkane sulfonamides (FASAs), $C_nF_{2n+1}SO_2NH_2$; FASA *N*-glucuronides, and perfluoroalkane sulfonic acids (PFSIAs), $C_nF_{2n+1}SO_2H$ (Lange 2000, 2001; Tomy, Tittlemier, et al. 2004; Xu et al. 2004; Boulanger et al. 2005; Xu et al. 2006; Rhoads et al. 2008; Xie et al. 2009) (Figure 8). There appears to be conflicting evidence as to whether PFOA can be formed in the environment from EtFOSE as a minor end product (Lange 2001; Tomy, Tittlemier, et al. 2004; Boulanger et al. 2005; Rhoads et al. 2008).

The *N*-alkyl perfluoroalkane sulfonamidoethyl acrylates and methacrylates, and polymers based on them, may undergo hydrolysis of the ester linkage in the environment to give *N*-alkyl FASEs (Martin et al. 2010) and, hence, lead to the same perfluoroalkyl biotransformation products. However, there do not appear to have been any published experimental studies that explicitly demonstrate this to be the case.




Figure 7. Perfluoroalkane sulfonamido derivatives synthesized from perfluoroalkane sulfonyl fluorides (PASFs), exemplified for a starting PASF with 8 C atoms. N.B. Names and acronyms for substance families are indicated. Those for the specific compounds shown can be found in the Supplemental Data.

Studies on the hydroxyl-radical-initiated degradation of EtFOSE in the aqueous phase show that some of the intermediates and products observed, including EtFOSAA, EtFOSA, FOSAA, FOSA, and PFOA, are the same as those reported for biodegradation. On the other hand, PFOS and PFOSI were not observed or were present at only trace levels in these abiotic studies (Hatfield 2001; Plumlee et al. 2009) and FOSA was considered to be a stable end product (Plumlee et al. 2009).

Atmospheric degradation pathways have been studied for 2 perfluoroalkane sulfonamido derivatives having 4 perfluorinated C atoms. The breakdown of EtFBSA, $C_4F_9SO_2NH(C_2H_5)$, has been shown to proceed through ketone and aldehyde intermediates to give PFCAs, i.e., PFBA, PFPrA and TFA, as well as COF_2 (Martin et al. 2006). The PFPrA and TFA are formed via chain unzipping of the perfluoroalkoxy radical, as already mentioned above for FTOHs and depicted schematically on Figure 6, so that alkyl-FASAs share part of their degradation scheme with FTOHs. PFBS was not observed to be formed from EtFBSA (Martin et al. 2006). MeFBSE, $C_4F_9SO_2N(CH_3)CH_2CH_2OH$, was observed to degrade to the same PFCAs as EtFBSA, together with PFBS, MeFBSA, and other products (D'eon et al. 2006).

Environmental occurrence of perfluoroalkyl sulfonamido derivatives: Various perfluoroalkyl sulfonamido derivatives have been found in the environment and human samples, whether this is due to industrial or consumer use of these compounds as such, losses during manufacturing operations, presence as “residuals” in other commercial products, or formation as environmental degradation products or metabolites of precursors.

It should be noted that perfluoroalkane sulfonamido derivatives bearing a H on the N atom are acidic in nature and can dissociate to an amide anion, to a greater or lesser extent depending on the ambient environmental or physiological conditions, with the degree of branching of the perfluoroalkyl chain having a significant influence on the pK_a for a given family of compounds (Rayne and Forest 2009a). For FASAs, there is the additional possibility of dissociation of the carboxylic H (more acidic than the amide H), whereas for the *N*-alkyl FASAs, this is the only possible ionization (Rayne and Forest 2009a). The dissociated species are not depicted in the list of compounds provided in the Supplemental Data.

All the families of perfluoroalkane sulfonamido derivatives discussed above and depicted in Tables 2 and 3 have been found in the environment or in human biota. Those with 8 perfluorinated C atoms are, in general, much more abundant than those with other chain lengths. However, more recently, compounds with 4 such C atoms have also been reported. The medium in which they are detected depends on their physical properties and on their likelihood of being formed there from precursors. In atmospheric air and its associated particulate matter, commonly detected compounds are the relatively volatile FOSA, MeFBSA, MeFOSA, Me₂FOSA, EtFOSA, MeFBSE, EtFBSE, MeFOSE, and EtFOSE (Martin et al. 2002; Barber et al. 2007; Piekarz et al. 2007; Stock et al. 2007; Dreyer, Matthias, et al. 2009; Dreyer, Weinberg, et al. 2009; Dreyer et al. 2010; Langer et al. 2010; Shoeib et al. 2010; Haug et al. 2011; Weinberg et al. 2011a, 2011b), whereas house dust has been found to contain FOSA, MeFOSA, EtFOSA, MeFOSE, EtFOSE (Shoeib et al. 2005; Kato et al. 2009; Goosey and Harrad 2011), the acrylate MeFOSAC (Shoeib et al. 2005), and the oxidation products MeFOSAA and EtFOSAA (Kato et al. 2009). FOSA has also




Figure 8. Transformation pathways for perfluoroalkane sulfonamido derivatives. Adapted from Olsen et al. 2002 and Olsen et al. 2005.

been detected in open ocean water, sometimes at levels comparable to those of PFOA (Ahrens, Gerwinski, et al. 2010; Ahrens, Xie, et al. 2010; Busch et al. 2010b; Kirchgeorg et al. 2010), as well as in precipitation (Kim and Kannan 2007; Taniyasu et al. 2008; Kwok et al. 2010), river and lake water (Kim and Kannan 2007; So et al. 2007; Ahrens et al. 2009b; Ahrens, Gerwinski, et al. 2010; Scott et al. 2010; Zushi et al. 2011), groundwater (Murakami, Kuroda, et al. 2009), surface runoff water (Kim and Kannan 2007; Murakami, Shinohara, et al. 2009), landfill leachate (Kallenborn et al. 2004; Busch et al. 2010a; Huset et al. 2011), sewage sludge (Llorca et al. 2011), and drinking water (Ericson et al. 2009). In wildlife, FOSA is often the predominant sulfonamido species, although it is generally present at lower levels than PFOS (Sturm and Ahrens 2010 and references therein), whereas EtFOSA and/or Et₂FOSA (Tomy, Budakowski, et al. 2004; Tittlemier et al. 2005; Tittlemier et al. 2006; Löfstrand et al. 2008; Ahrens, Siebert, et al. 2009; Yeung et al. 2009), MeFOSE (Ahrens and Ebinghaus 2010), FOSAA (Peng et al. 2010), and EtFOSAA (Yoo et al. 2009) have also been reported. FOSA and various *N*-alkyl-FOSAs (Me-, Et-, Me₂-, and Et₂-FOSAs) were detected in foodstuffs (Tittlemier et al. 2005; Tittlemier et al. 2006). WWTP effluents and river, coastal, and ocean waters were found to contain some *N*-alkyl sulfonamido derivatives (MeFBSA, MeFBSE, MeFOSE, EtFOSE, MeFB-SAA, MeFOSAA, and EtFOSAA) as well as FOSA and FOSAA (Ahrens et al. 2009a; Ahrens et al. 2009b; Ahrens, Gerwinski, et al. 2010; Huset et al. 2011; Nguyen et al. 2011; Zushi et al. 2011). In human blood, the sulfonamido derivatives FOSA, FOSAA, MeFOSAA, and EtFOSAA have been quantified (Kannan et al. 2004; Calafat et al. 2007; Olsen et al. 2008; Weihe et al. 2008; Toms et al. 2009; Lee and Mabury 2011). MeFOSAA and/or EtFOSAA have also been detected in precipitation (Taniyasu et al. 2008; Kwok et al. 2010), wildlife (Yoo et al. 2009), sediments (Higgins et al. 2005; Ahrens, Taniyasu, et al. 2010; Zushi et al. 2010) and WWTP influent and effluent (Boulanger et al. 2005). These 2 compounds have also been shown to be among the most abundant PFAS components of municipal WWTP sludge (Higgins et al. 2005; Sepulvado et al. 2011), in which FOSAA has also been detected (Higgins et al. 2005).

Perfluoroalkyl and polyfluoroalkyl ether carboxylic acids. Salts of perfluoroalkyl ether carboxylic acids (not depicted in the tables) and polyfluoroalkyl ether carboxylic acids (Table 3) are widely cited in patents as alternative fluoropolymer processing aids, that are more environmentally and/or toxicologically acceptable alternatives to APFO and APFN. A common feature is that a terminal -COO⁻ group, attached to one or both ends of the fluorinated ether chain, is the common hydrophile, generally with an NH₄⁺ counter-ion (Tsuda et al. 2003; Visca et al. 2003; Higuchi et al. 2005; Hintzer et al. 2005; Brothers et al. 2008; Ishikawa et al. 2008; Gordon 2011). These and/or other alternative surfactants are expected to enable manufacturers to meet the USEPA 2010/15 Stewardship Program goal to eliminate the use of PFOA and higher homologues. Most recently, a toxicological evaluation for one of these substances (ammonium 4,8-dioxa-3*H*-perfluorononanoate; Table 3) has been published (Gordon 2011). Substances based on certain members of this family of compounds have a sufficient number of repeating units (together with other characteristics) to enable them to

be considered to be polymers under the European Union REACH legislation (ECHA 2008).

Fluorinated polymers

The polymers discussed in this section are those: 1) whose synthesis involves the incorporation of one or more PFASs as monomers. In this case, there is some potential (theoretical or demonstrated) for the degradation of the polymer, during or after its useful lifetime, to lead to release of PFASs to the environment; or 2) whose manufacture requires the use of a PFAS as a processing aid.

Fluoropolymers. Fluoropolymers contain F bound to one or both of the olefinic C atoms, to form a perfluorinated C-only polymer backbone with F atoms directly attached to it (Table 4). Examples of fluoropolymers are polytetrafluoroethylene (PTFE); polyvinylidene fluoride (PVDF); polyvinyl fluoride (PVF); copolymers of tetrafluoroethylene (TFE) and hexafluoropropylene (HFP); terpolymers of TFE, vinylidene fluoride, and HFP; and copolymers of TFE and ethylene. Certain grades of fluoropolymers, manufactured by emulsion (or dispersion) polymerization, in order to obtain a fine particle size distribution, require the use of a fluorosurfactant “processing aid.” This additive, used at a level of a few tenths of a percent relative to the amount of polymer produced (Prevedouros et al. 2006), was often traditionally the ammonium salt of PFOA or PFNA. The fluorosurfactant is removed when the fluoropolymer aqueous emulsion is dried for sale as a solid. Similarly, when an aqueous fluoropolymer emulsion is used, the polymer is heated to cure it. High cure temperatures thermally destroy the fluorosurfactant. At low cure temperatures, residual surfactant may remain (Guo et al. 2009). Most producers have discontinued the use of PFOA and PFNA salts as processing aids and have developed and implemented more environmentally acceptable alternatives, as discussed above in the *Perfluoroalkyl and Polyfluoroalkyl Ether Carboxylic Acids* section. It should be emphasized that those grades of fluoropolymers (e.g., PTFE, PVDF) that are made by suspension (rather than emulsion) polymerization do not require a fluorosurfactant to be used as a “processing aid.”






Perfluoropolyethers. Perfluoropolyethers (PFPEs; Table 4) are polymers in whose backbone -CF₂-, -CF₂CF₂-, and possibly -CF(CF₃)CF₂- units are separated by O atoms. For example, the ultraviolet-initiated copolymerization of TFE with O₂ leads to PFPEs with a structure that may be represented symbolically by CF₃O(CF₂CF₂O)_m(CF₂O)_nCF₃, although this overall formula does not show that the -CF₂O- and -CF₂CF₂O- units are generally distributed randomly rather than in blocks (Sianesi et al. 1994). If the photopolymerization is conducted using hexafluoropropylene (HFP) instead of (or together with) TFE, then PFPEs with the overall formula CF₃O(CF₂CF₂O)_m(CF₂O)_n[CF(CF₃)CF₂O]_pCF₃ are obtained. Furthermore, the PFPE -[CF(CF₃)CF₂O]_n- can be synthesized by homopolymerization of HFP (ep)oxide.

Because the repeating units of these PFPEs contain only 2 or 3 perfluorinated C atoms per O atom, their degradation cannot lead to the formation of long-chain PFCAs. The reason for mentioning them in this review is that certain difunctional polymeric perfluoro-polyether products, corresponding to the

overall formula $X-CF_2O(CF_2CF_2O)_m(CF_2O)_nCF_2-X$, where X is a hydrophilic group, are marketed as surface treatments for natural stone, metal, glass, plastic, textiles, leather, and paper and paperboard treatment for food-contact applications. These functionalized PFPEs bring properties such as a low surface energy, high contact angle, reduced coefficient of friction, and high oleo-hydrophobicity (Solvay Solexis 2011), so that they are potential alternatives to the ECF-based polymers, fluorotelomer-based polymers, and fluorinated oxetane polymers described in this review.

Side-chain-fluorinated polymers. In contrast to the polymers described previously, side-chain-fluorinated polymers do not have perfluorinated or polyfluorinated polymer backbones, but are composed of variable composition backbones with polyfluoroalkyl (and possibly perfluoroalkyl) side chains (Table 4). With regard to the sources of long-chain PFAAs, we review 3 groups of side-chain-fluorinated polymers distinguished from one another by the linkage (acrylate and/or methacrylate, urethane, and oxetane) between the polymer backbone and the polyfluoroalkyl (and possibly perfluoroalkyl) side chains. Side chains of each of these polymer types may possess the ability to sever from the polymer chain to become PFASs shown in Tables 2 and 3. It should be noted, however, that this transformation process can occur over long time periods (e.g., >1000 y) and may exhibit low yields of PFASs such that their contribution to the environmental inventory of long-chain PFAAs may be insignificant relative to other historical and current sources. Further research is required to clarify this question.

Fluorinated acrylate polymers: Fluorinated acrylate polymers are made by polymerizing a fluorinated acrylate (or methacrylate) monomer, in which the alcohol moiety is $n:2$ FTOH, $C_nF_{2n+1}CH_2CH_2OH$, or an alkyl-FASE, $C_nF_{2n+1}SO_2N(R)CH_2CH_2OH$, where $R = CH_3$, C_2H_5 , or another alkyl group (Table 4). Some possible structures for the fluorinated acrylate monomers are therefore:



These fluorinated acrylate monomers are copolymerized with one or more nonfluorinated acrylate monomers, and possibly other monomers, to give the final side-chain fluorinated acrylate polymers. These types of polymers are useful as water-, stain- and grease-proofing finishes for textile, leather, and paper surfaces. As stated above, it is not yet clear to what extent such polymers may break down in the environment to give PFAAs, such as PFOA, PFOS, PFBA, and PFBS. Moreover, although we have shown only fluorotelomer and perfluoroalkane sulfonamido (meth)acrylates, the term “side-chain-fluorinated polymer” would encompass many other potential structures and products therefrom that conform to the definition provided.

Fluorinated urethane polymers: Polymeric materials for repelling water and stains may also be based on urethane polymers formed by reacting fluorotelomer alcohols (FTOHs), or perfluoroalkane sulfonamidoethanols (alkyl-FASEs), with polyisocyanato homopolymers, followed by a cross-linking step (Kirchner 1989). The products are polyfluorinated in their side chains (Table 4). They are used mainly in textile applications. In the case of an (8:2) FTOH-based urethane polymer, a recent study has shown that the half-life with respect to biodegradation to PFOA in aerobic soils is on the order of a century (Russell et al. 2010).

Fluorinated oxetane polymers: An alternative fluorinated polymer technology to those described thus far originates from the reaction of polyfluorinated alcohols with oxetanes bearing a $-CH_2Br$ group in their side chains, to create oxetane monomers that can undergo ring-opening polymerization to give side-chain-polyfluorinated polyethers (Figure 9). These fluorinated oxetane polymers (Table 4) are offered in many forms and functionalities primarily as fluorosurfactants and coatings additives (Kausch et al. 2002; Kausch et al. 2003a, 2003b; Thomas 2006; Omnova Solutions 2011).

Commercial articles containing multiple types of fluorinated polymers. It should be noted that there are commercial products that contain both fluoropolymers and side-chain-fluorinated polymers, which can cause confusion about the origin of individual PFASs. In all-weather clothing products, for example, multiple layered materials containing different types of polymers are common. A porous PTFE membrane layer is often used in garments to make the fabric “breathable.” The outer fabric layer may be nylon or polyester




Figure 9. Oxetane-based fluorinated polymers.

treated with a side-chain-fluorinated polymer water repellent. Analyses of all-weather clothing revealed the presence of FTOHs in the outer layer of some all-weather clothing products (Berger and Herzke 2006; Schulze and Norin 2006). The origin of the FTOHs is not the PTFE breathable membrane.

SUMMARY AND FUTURE PROSPECTS

We have provided an overview of PFASs detected in the environment, wildlife, and humans and recommended clear, specific, and descriptive terminology, names, and acronyms for PFASs. We hope the terminology will be widely adopted and used. Future interest in fluorinated substances by the global scientific community is expected to remain high, and continued publications should be numerous. The consistent use of the terminology described here by this community will facilitate clear and coherent communication, understanding, interpretation, and comparison of published studies as well as serve to highlight similarities and acknowledge key differences between PFASs. We strongly discourage the use of broad, poorly defined terms and acronyms in favor of the clear, specific, and descriptive terminology provided here.

SUPPLEMENTAL DATA

Supplemental Data. Names, formulas, acronyms, and CAS numbers for selected perfluoroalkyl and polyfluoroalkyl substances. Terminology decision flow charts.

Acknowledgment—We thank the PlasticsEurope trade association, Fluoropolymers Committee, for initiating this project and for providing J Franklin with funding. We are also grateful to our colleagues and students for their helpful and constructive review of this work.

REFERENCES

- (NOTE: For patents, the years indicated are those of filing of the earliest priority application.)
- [3M] Company. 1999. Fluorochemical use, distribution and release overview. USEPA Administrative Record AR226-0550. Available from: www.regulations.gov as document EPA-HQ-OPPT-2002-0051-0003.
- [3M] Company. 2000a. Voluntary Use and Exposure Information Profile for Perfluorooctanoic Acid and Salts. USEPA Administrative Record AR226-0595. Available from: www.regulations.gov as document EPA-HQ-OPPT-2002-0051-0009.
- [3M] Company. 2000b. Re: CAS Names for PFOS-Based Products. USEPA Administrative Record AR226-0931. Available from: OPPT.NCIC@epamail.epa.gov
- [3M] Company. 2000c. Re: Phase-Out Plan for POSF-based Products. USEPA Administrative Record AR226-0600. Available from: www.regulations.gov, as document EPA-HQ-OPPT-2002-0051-0006.
- Ahrens L. 2011. Polyfluoroalkyl compounds in the aquatic environment: a review of their occurrence and fate. *J Environ Monit* 13:20–31.
- Ahrens L, Ebinghaus R. 2010. Spatial distribution of polyfluoroalkyl compounds in dab (*Limanda limanda*) bile fluids from Iceland and the North Sea. *Mar Pollut Bull* 60:139–151.
- Ahrens L, Siebert U, Ebinghaus R. 2009. Total body burden and tissue distribution of polyfluorinated compounds in harbor seals (*Phoca vitulina*) from the German Bight. *Mar Pollut Bull* 58:520–525.
- Ahrens L, Felizeter S, Ebinghaus R. 2009a. Spatial distribution of polyfluoroalkyl compounds in seawater of the German Bight. *Chemosphere* 76:179–184.
- Ahrens L, Felizeter S, Sturm R, Xie Z, Ebinghaus R. 2009b. Polyfluorinated compounds in waste water treatment plant effluents and surface waters along the River Elbe, Germany. *Mar Pollut Bull* 58:1326–1333.
- Ahrens L, Gerwinski W, Theobald N, Ebinghaus R. 2010. Sources of polyfluoroalkyl compounds in the North Sea, Baltic Sea and Norwegian Sea: Evidence from their spatial distribution in surface water. *Mar Pollut Bull* 60:255–260.
- Ahrens L, Taniyasu S, Yeung LWY, Yamashita N, Lam PKS, Ebinghaus R. 2010. Distribution of polyfluoroalkyl compounds in water, suspended particulate matter and sediment from Tokyo Bay, Japan. *Chemosphere* 79:266–272.
- Ahrens L, Xie Z, Ebinghaus R. 2010. Distribution of perfluoroalkyl compounds in seawater from Northern Europe, Atlantic Ocean, and Southern Ocean. *Chemosphere* 78:1011–1016.
- Ahrens L, Shoeib M, Harner T, Lee SC, Guo R, Reiner EJ. 2011. Wastewater treatment plant and landfills as sources of polyfluoroalkyl compounds to the atmosphere. *Environ Sci Technol* DOI: 10.1021/es1036173
- Alsmeyer YW, Childs WV, Flynn RM, Moore GGI, Smeltzer JC. 1994. Electrochemical fluorination and its applications. In: Banks RE, Smart BE, Tatlow JC, editors. *Organofluorine chemistry: Principles and commercial applications*. New York (NY): Plenum. p 121–144.
- Aoyama H, Chiba Y. 1997. Process of preparing perfluoroalkylcarboxylic acid. Washington (DC): US Patent office 5945562. Daikin.
- Armitage J, Cousins IT, Buck RC, Prevedouros K, Russell MH, MacLeod M, Korzeniowski SH. 2006. Modeling global-scale fate and transport of perfluorooctanoate emitted from direct sources. *Environ Sci Technol* 40:6969–6975.
- Armitage JM, MacLeod M, Cousins IT. 2009a. Comparative assessment of the global fate and transport pathways of long-chain perfluorocarboxylic acids (PFCAs) and perfluorocarboxylates (PFCs) emitted from direct sources. *Environ Sci Technol* 43:5830–5836.
- Armitage JM, MacLeod M, Cousins IT. 2009b. Modeling the global fate and transport of perfluorooctanoic acid (PFOA) and perfluorooctanoate (PFO) emitted from direct sources using a multispecies mass balance model. *Environ Sci Technol* 43:1134–1140.
- Arsenault G, Chittim B, McAlees A, McCrindle R, Riddell N, Yeo B. 2008. Some issues relating to the use of perfluorooctanesulfonate (PFOS) samples as reference standards. *Chemosphere* 70:616–625.
- Banks RE, Smart BE, Tatlow JC. 1994. *Organofluorine chemistry: Principles and commercial applications*. New York (NY): Plenum. 670 p.
- Barber JL, Berger U, Chaemfa C, Huber S, Jahnke A, Temme C, Jones KC. 2007. Analysis of per- and polyfluorinated alkyl substances in air samples from Northwest Europe. *J Environ Monit* 9:530–541.
- Begley TH, Hsu W, Noonan G, Diachenko G. 2008. Migration of fluorochemical paper additives from food-contact paper into foods and food simulants. *Food Addit Contam* 25:384–390.
- Belisle J. 1981. Organic fluorine in human serum: Natural versus industrial sources. *Science* 212:1509–1510.
- Benskin JP, Bataineh M, Martin JW. 2007. Simultaneous characterization of perfluoroalkyl carboxylate, sulfonate, and sulfonamide isomers by liquid chromatography-tandem mass spectrometry. *Anal Chem* 79:6455–6464.
- Benskin JP, De Silva AO, Martin JW. 2010. Isomer profiling of perfluorinated substances as a tool for source tracking: a review of early findings and future applications. *Rev Environ Contam Toxicol* 208:111–160.
- Benskin JP, Yeung LWY, Yamashita N, Taniyasu S, Lam PKS, Martin JW. 2010. Perfluorinated acid isomer profiling in water and quantitative assessment of manufacturing source. *Environ Sci Technol* 44:9049–9054.
- Berger U, Herzke D. 2006. Per- and polyfluorinated alkyl substances (PFAS) extracted from textile samples. *Organohalogen Compd* 68:2023–2026.
- Bertocchio R, Foulletier L. 1970. Amphoteric fluorinated sulfonamides. British Patent 1343990. Ugine Kuhlmann.
- Boulanger B, Vargo JD, Schnoor JL, Hornbuckle KC. 2005. Evaluation of perfluorooctane surfactants in a wastewater treatment system and in a commercial surface protection product. *Environ Sci Technol* 39:5524–5530.
- Boutonnet JC, Bingham P, Calamari D, de Rooij C, Franklin J, Kawano T, Libre J-M, McCulloch A, Malinverno G, Odom JM., et al. 1999. Environmental risk assessment of trifluoroacetic acid. *Hum Ecol Risk Assess* 5:59–124.
- Brandsma SH, Smithwick M, Solomon K, Small J, de Boer J, Muir DCG. 2011. Dietary exposure of rainbow trout to 8:2 and 10:2 fluorotelomer alcohols and

- perfluorooctanesulfonamide: Uptake, transformation and elimination. *Chemosphere* 82:253–258.
- Brooke D, Footitt A, Nwaogu TA. 2004. Environmental risk evaluation report: Perfluorooctane sulphonate (PFOS). London (UK): UK Environment Agency.
- Brothers PD, Gangal SV, Chapman GA, Howell JL, Smith AP. 2008. Aqueous polymerization of fluorinated monomer using a mixture of fluoropolyether acids or salts. Washington (DC): US Patent Office. Patent application 2009/0281241. Du Pont.
- Burns DC, Ellis DA, Li H, McMurdo CJ, Webster E. 2008. Experimental pK_a determination for perfluorooctanoic acid (PFOA) and the potential impact of pK_a concentration dependence on laboratory-measured partitioning phenomena and environmental modeling. *Environ Sci Technol* 42:9283–9288.
- Busch J, Ahrens L, Sturm R, Ebinghaus R. 2010a. Polyfluoroalkyl compounds in landfill leachates. *Environ Pollut* 158:1467–1471.
- Busch J, Ahrens L, Xie Z, Sturm R, Ebinghaus R. 2010b. Polyfluoroalkyl compounds in the East Greenland Arctic Ocean. *J Environ Monit* 12:1242–1246.
- Butt CM, Mabury SA, Muir DCG, Braune BM. 2007. Prevalence of long-chained perfluorinated carboxylates in seabirds from the Canadian Arctic between 1975 and 2004. *Environ Sci Technol* 41:3521–3528.
- Butt CM, Muir DCG, Stirling I, Kwan M, Mabury SA. 2007. Rapid response of Arctic ringed seals to changes in perfluoroalkyl production. *Environ Sci Technol* 41:42–49.
- Butt CM, Young CJ, Mabury SA, Hurley MD, Wallington TJ. 2009. Atmospheric chemistry of 4:2 fluorotelomer acrylate [$C_4F_9CH_2CH_2OC(O)CH=CH_2$]: Kinetics, mechanisms, and products of chlorine-atom- and OH-radical-initiated oxidation. *J Phys Chem A* 113:3155–3161.
- Butt CM, Berger U, Bossi R, Tomy GT. 2010. Review: Levels and trends of poly- and perfluorinated compounds in the arctic environment. *Sci Total Environ* 408:2936–2965.
- Butt CM, Muir DCG, Mabury SA. 2010a. Elucidating the pathways of poly- and perfluorinated acid formation in rainbow trout. *Environ Sci Technol* 44:4973–4980.
- Butt CM, Muir DCG, Mabury SA. 2010b. Biotransformation of the 8:2 fluorotelomer acrylate in rainbow trout. 1. In vivo dietary exposure. *Environ Toxicol Chem* 29:2726–2735.
- Calafat AM, Kuklenyik Z, Reidy JA, Caudill SP, Tully JS, Needham LL. 2007. Serum concentrations of 11 polyfluoroalkyl compounds in the U.S. population: Data from the National Health and Nutrition Examination Survey (NHANES) 1999–2000. *Environ Sci Technol* 41:2237–2242.
- Cheng J, Psillakis E, Hoffmann MR, Colussi AJ. 2009. Acid dissociation versus molecular association of perfluoroalkyl oxoacids: Environmental implications. *J Phys Chem A* 113:8152–8156.
- Coe PL, Milner NE. 1972. Perfluoroalkylcopper(I) compounds. II. Reactions of perfluoroalkylcopper(I) compounds and perfluoroalkyl iodide/copper mixtures with olefins. *J Organomet Chem* 39:395–402.
- Conder JM, Hoke RA, de Wolf W, Russell MH, Buck RC. 2008. Are PFASs bioaccumulative? A critical review and comparison with regulatory criteria and persistent lipophilic compounds. *Environ Sci Technol* 42:995–1003.
- D'eon JC, Mabury SA. 2007. Production of perfluorinated carboxylic acids (PFACs) from the biotransformation of polyfluoroalkyl phosphate surfactants (PAPS): Exploring routes of human contamination. *Environ Sci Technol* 41:4799–4805.
- D'eon JC, Mabury SA. 2010. Uptake and elimination of perfluorinated phosphonic acids in the rat. *Environ Toxicol Chem* 29:1319–1329.
- D'eon JC, Mabury SA. 2011. Exploring indirect sources of human exposure to perfluoroalkyl carboxylates (PFACs): Evaluating uptake, elimination and biotransformation of polyfluoroalkyl phosphate esters (PAPS) in the rat. *Environ Health Perspect* 119:344–350.
- D'eon JC, Hurley MD, Wallington TJ, Mabury SA. 2006. Atmospheric chemistry of *N*-methyl perfluorobutane sulfonamidoethanol, $C_4F_9SO_2N(CH_3)CH_2CH_2OH$: Kinetics and mechanism of reaction with OH. *Environ Sci Technol* 40:1862–1868.
- D'eon JC, Crozier PW, Furdui VI, Reiner EJ, Libelo EL, Mabury SA. 2009a. Observation of a commercial fluorinated material, the polyfluoroalkyl phosphoric acid diesters, in human sera, wastewater treatment plant sludge, and paper fibers. *Environ Sci Technol* 43:4589–4594.
- D'eon JC, Crozier PW, Furdui VI, Reiner EJ, Libelo EL, Mabury SA. 2009b. Perfluorinated phosphonic acids in Canadian surface waters and wastewater treatment plant effluent: Discovery of a new class of perfluorinated acids. *Environ Toxicol Chem* 28:2101–2107.
- D'Hollander W, de Voogt P, De Coen W, Bervoets L. 2010. Perfluorinated substances in human food and other sources of human exposure. *Rev Environ Contam Toxicol* 208:179–215.
- Dasu K, Royer LA, Liu J, Lee LS. 2010. Hydrolysis of fluorotelomer compounds leading to fluorotelomer alcohol production during solvent extractions of soils. *Chemosphere* 81:911–917.
- De Silva AO, Mabury SA. 2004. Isolating isomers of perfluorocarboxylates in polar bears (*Ursus maritimus*) from two geographical locations. *Environ Sci Technol* 38:6538–6545.
- De Silva AO, Mabury SA. 2006. Isomer distribution of perfluorocarboxylates in human blood: Potential correlation to source. *Environ Sci Technol* 40:2903–2909.
- De Silva AO, Muir DCG, Mabury SA. 2009. Distribution of perfluorocarboxylate isomers in select samples from the North American environment. *Environ Toxicol Chem* 28:1801–1814.
- de Voogt P. 2009. The PERFOOD project. [cited 2011 February 8]. Available from: http://www.perfood.eu/index.php?option=com_content&view=article&id=49&Itemid=56
- de Voogt P. 2010. Special foreword: Perfluorinated alkylated substances. *Rev Environ Contam Toxicol* 208:v–vii.
- de Voogt P, Berger U, de Coen W, de Wolf W, Heimstad E, McLachlan M, van Leeuwen S, van Roon A. 2006. PERFORCE: Perfluorinated organic compounds in the European environment. [cited 2011 February 8]. Available from: <http://www.science.uva.nl/perforce/Final%20reportA.pdf>
- Dinglasan MJA, Ye Y, Edwards EA, Mabury SA. 2004. Fluorotelomer alcohol biodegradation yields poly- and perfluorinated acids. *Environ Sci Technol* 38:2857–2864.
- Dinglasan-Panlilio MJA, Mabury SA. 2006. Significant residual fluorinated alcohols present in various fluorinated materials. *Environ Sci Technol* 40:1447–1453.
- Dreyer A, Matthias V, Temme C, Ebinghaus R. 2009. Annual time series of air concentrations of polyfluorinated compounds. *Environ Sci Technol* 43:4029–4036.
- Dreyer A, Weinberg I, Temme C, Ebinghaus R. 2009. Polyfluorinated compounds in the atmosphere of the Atlantic and Southern Oceans: Evidence for a global distribution. *Environ Sci Technol* 43:6507–6514.
- Dreyer A, Shoeib M, Fiedler S, Barber J, Harner T, Schramm K-W, Jones KC, Ebinghaus R. 2010. Field intercomparison on the determination of volatile and semivolatile polyfluorinated compounds in air. *Environ Chem* 7:350–358.
- [ECHA] European Chemicals Agency. 2008. Guidance for monomers and polymers. [cited 2011 February 8]. Available from: http://guidance.echa.europa.eu/docs/guidance_document/polymers_en.pdf
- Eggen T, Moeder M, Arukwe A. 2010. Municipal landfill leachates: A significant source for new and emerging pollutants. *Sci Total Environ* 408:5147–5157.
- Ellis DA, Mabury SA, Martin JW, Muir DCG. 2001. Thermolysis of fluoropolymers as a potential source of halogenated organic acids in the environment. *Nature* 412:321–324.
- Ellis DA, Martin JW, De Silva AO, Mabury SA, Hurley MD, Sulbaek Andersen MP, Wallington TJ. 2004. Degradation of fluorotelomer alcohols: A likely atmospheric source of perfluorinated carboxylic acids. *Environ Sci Technol* 38:3316–3321.
- Environment Canada. 2010. Environmental performance agreement ("agreement") respecting perfluorinated carboxylic acids (PFACs) and their precursors in perfluorochemical products sold in Canada. [cited 2011 February 8]. Available from: <http://ec.gc.ca/epe-epa/default.asp?lang=En&n=81AE80CE-1>
- Ericson I, Domingo JL, Nadal M, Bigas E, Llebaria X, van Bavel B, Lindström G. 2009. Levels of perfluorinated chemicals in municipal drinking water from Catalonia, Spain: Public health implications. *Arch Environ Contam Toxicol* 57:631–638.

- Eschauzier C, Haftka J, Stuyfzand PJ, de Voogt P. 2010. Perfluorinated compounds in infiltrated River Rhine water and infiltrated rainwater in coastal dunes. *Environ Sci Technol* 44:7450–7455.
- European Parliament. 2006a. Regulation (EC) No 1907/2006 of the European Parliament and of the Council of 18 December 2006 concerning the Registration, Evaluation, Authorisation and Restriction of Chemicals (REACH). Official Journal of the EU, 30 December 2006, L396/1–L396/849.
- European Parliament. 2006b. Directive 2006/122/EC of the European Parliament and of the Council of 12 December 2006. Official Journal of the EU, 27 December 2006, L372/32–L372/34.
- Falk RA. 1982. Perfluoroalkyl amphoteric compounds. European Patent 115251. Ciba-Geigy.
- Fasano WJ, Carpenter SC, Gannon SA, Snow TA, Stadler JC, Kennedy GL, Buck RC, Korzeniowski SH, Hinderliter PM, Kemper RA. 2006. Absorption, distribution, metabolism, and elimination of 8-2 fluorotelomer alcohol in the rat. *Toxicol Sci* 91:341–355.
- Fasano WJ, Sweeney LM, Mawn MP, Nabb DL, Szostek B, Buck RC, Gargas ML. 2009. Kinetics of 8-2 fluorotelomer alcohol and its metabolites, and liver glutathione status following daily oral dosing for 45 days in male and female rats. *Chem Biol Interact* 180:281–295.
- [FDA] US Food and Drug Administration. 2009. Indirect food additives: Paper and paperboard components. US FDA, Code of Federal Regulations 21 CFR 176.170.
- Federal Register. 2006a. Inert ingredient; Revocation of the tolerance exemption for mono- and bis-(1H, 1H, 2H, 2H-perfluoroalkyl) phosphates where the alkyl group is even numbered and in the C₆–C₁₂ range. *Fed Reg* 71(153):45408–45411. 9 August 2006.
- Federal Register. 2006b. Perfluoroalkyl sulfonates; Proposed significant new use rule. *Fed Reg* 71 (47): 12311–12324. 10 March 2006. Available from: www.regulations.gov, as document EPA-HQ-OPPT-2005-0015-0001.
- Fiedler S, Pfister G, Schramm K-W. 2010. Poly- and perfluorinated compounds in household consumer products. *Toxicol Environ Chem* 92:1801–1811.
- Fluoropolymer Manufacturing Group. 2001. Guide to the safe handling of fluoropolymer dispersions. Washington (DC): Society of Plastics Industry.
- Frömel T, Knepper TP. 2010. Biodegradation of fluorinated alkyl substances. *Rev Environ Contam Toxicol* 208:161–177.
- Furdui VI, Stock NL, Ellis DA, Butt CM, Whittle DM, Crozier PW, Reiner EJ, Muir DCG, Mabury SA. 2007. Spatial distribution of perfluoroalkyl contaminants in lake trout from the Great Lakes. *Environ Sci Technol* 41:1554–1559.
- Furdui VI, Helm PA, Crozier PW, Lucaciu C, Reiner EJ, Marvin CH, Whittle DM, Mabury SA, Tomy GT. 2008. Temporal trends of perfluoroalkyl compounds with isomer analysis in lake trout from Lake Ontario (1979–2004). *Environ Sci Technol* 42:4739–4744.
- Gebbink WA, Hebert CE, Letcher RJ. 2009. Perfluorinated carboxylates and sulfonates and precursor compounds in herring gull eggs from colonies spanning the Laurentian Great Lakes of North America. *Environ Sci Technol* 43:7443–7449.
- Giesy JP, Kannan K. 2001. Global distribution of perfluorooctane sulfonate in wildlife. *Environ Sci Technol* 35:1339–1342.
- Goosey E, Harrad S. 2011. Perfluoroalkyl compounds in dust from Asian, Australian, European, and North American homes and UK cars, classrooms, and offices. *Environ Int* 37:86–92.
- Gordon SC. 2011. Toxicological evaluation of ammonium 4,8-dioxo-3H-perfluorononanoate, a new emulsifier to replace ammonium perfluorooctanoate in fluoropolymer manufacturing. *Regul Toxicol Pharm* 59:64–80.
- Goss K-U. 2008. The pK_a values of PFOA and other highly fluorinated carboxylic acids. *Environ Sci Technol* 42:456–458.
- Grottenmüller R, Knaup W, Probst A, Dullinger K. 2000. Process for the preparation of perfluorocarboxylic acids. Washington (DC): US Patent Office. Patent 6,515,172. Clariant.
- Guo Z, Liu X, Krebs KA, Roache NF. 2009. Perfluorocarboxylic acid content in 116 articles of commerce. Report EPA/600/R-09/033. Research Triangle Park (NC): US Environmental Protection Agency.
- Guruge KS, Yeung LWY, Li P, Taniyasu S, Yamashita N, Nakamura M. 2011. Fluorinated alkyl compounds including long chain carboxylic acids in wild bird livers from Japan. *Chemosphere* 83:379–384.
- Hagen DF, Belisle J, Johnson JD, Venkateswarlu P. 1981. Characterization of fluorinated metabolites by a gas chromatographic-helium microwave plasma detector. The biotransformation of 1H,1H,2H,2H-perfluorodecanol to perfluorooctanoate. *Anal Biochem* 118:336–343.
- Han W. 2009. PFOS related actions in China. In: International Workshop on Managing Perfluorinated Chemicals and Transitioning to Safer Alternatives; 2009 February 12–13; Geneva, Switzerland. Available from: http://www.chem.unep.ch/unepsaicm/cheminprod_dec08/PFCWorkshop/Presentations/HWenya%20-%20PFOS%20in%20China.pdf
- Hansen KJ, Clemen LA, Ellefson ME, Johnson HO. 2001. Compound-specific, quantitative characterization of organic fluorochemicals in biological matrices. *Environ Sci Technol* 35:766–770.
- Hatfield TL. 2001. Screening studies on the aqueous photolytic degradation of 2-(N-ethylperfluorooctanesulfonamido)-ethyl alcohol (N-ETFOSE alcohol). 3M Company, Project W2783. USEPA Administrative Record AR226-1030a.080. Available from: OPPT.NCIC@epamail.epa.gov
- Haug LS, Huber S, Schlabach M, Becher G, Thomsen C. 2011. Investigation on per- and polyfluorinated compounds in paired samples of house dust and indoor air from Norwegian homes. *Environ Sci Technol* DOI: 10.1021/es103456h
- Hekster FM, de Voogt P, Pijnenburg AMCM, Laane RWPM. 2002. Perfluoroalkylated substances: Aquatic environmental assessment Report RIKZ/2002.043. The Hague (NL): National Institute for Coastal and Marine Management (RIKZ).
- Hekster FM, Laane RWPM, de Voogt P. 2003. Environmental and toxicity effects of perfluoroalkylated substances. *Rev Environ Contam Toxicol* 179:99–121.
- Higgins CP, Field JA, Criddle CS, Luthy RG. 2005. Quantitative determination of perfluorochemicals in sediments and domestic sludge. *Environ Sci Technol* 39:3946–3956.
- Higuchi S, Kamiya H, Hoshikawa J, Matsuoka Y. 2005. Aqueous polytetrafluoroethylene emulsion, and polytetrafluoroethylene fine powder and porous material produced from the same. European Patent 1939222. Asahi Glass.
- Hintzer K, Jurgens M, Kaspar H, Königsmann H, Maurer AR, Schwertfeger W, Zipplies TC. 2005. Fluorinated surfactants for making fluoropolymers. Washington (DC): US Patent Office. Patent application 2007/142541. 3M Innovative Properties Co.
- Houde M, Wells RS, Fair PA, Bossart GD, Hohn AA, Rowles TK, Sweeney JC, Solomon KR, Muir DCG. 2005. Polyfluoroalkyl compounds in free-ranging bottlenose dolphins (*Tursiops truncatus*) from the Gulf of Mexico and the Atlantic Ocean. *Environ Sci Technol* 39:6591–6598.
- Houde M, Martin JW, Letcher RJ, Solomon KR, Muir DCG. 2006. Biological monitoring of polyfluoroalkyl substances: A review. *Environ Sci Technol* 40:3463–3473.
- Houde M, De Silva AO, Muir DCG, Letcher RJ. 2011. Monitoring of perfluorinated compounds in aquatic biota: An updated review. *Environ Sci Technol* DOI: 10.1021/es104326w
- Howard PH, Muir DCG. 2010. Identifying new persistent and bioaccumulative organics among chemicals in commerce. *Environ Sci Technol* 44:2277–2285.
- Hurley MD, Ball JC, Wallington TJ, Sulbaek Andersen MP, Ellis DA, Martin JW, Mabury SA. 2004. Atmospheric chemistry of 4:2 fluorotelomer alcohol (CF₃(CF₂)₃CH₂CH₂OH): Products and mechanism of Cl atom initiated oxidation. *J Phys Chem A* 108:5635–5642.
- Huset C, Chiaia AC, Barofsky DF, Jonkers N, Kohler H-PE, Ort C, Giger W, Field JA. 2008. Occurrence and mass flows of fluorochemicals in the Glatt Valley watershed, Switzerland. *Environ Sci Technol* 42:6369–6377.
- Huset CA, Barlaz MA, Barofsky DF, Field JA. 2011. Quantitative determination of fluorochemicals in municipal landfill leachates. *Chemosphere* 82:1380–1386.
- [INAC] Indian and Northern Affairs Canada. 2009. Indian and Northern Affairs Canada—Blueprints and proposal guidelines: Call for proposals 2010–2011. [cited 2011 February 8]. Available from: <http://www.ainc-inac.gc.ca/nth/ct/ncp/pubs/calprop/cfp1011-eng.pdf>

- Ishikawa T, Tsuda N, Yamamoto Y. 2008. Fluoroethercarboxylic acid and production method thereof, surfactant, method of producing fluoropolymer and aqueous dispersion used thereof. Washington (DC): US Patent Office. Patent application 2009/0176942. Daikin.
- Jackson DA, Young CJ, Hurley MD, Wallington TJ, Mabury SA. 2011. Atmospheric degradation of perfluoro-2-methyl-3-pentanone: Photolysis, hydrolysis and hydration. *Environ Sci Technol* DOI: 10.1021/es104362g
- Jahnke A, Berger U. 2009. Trace analysis of per- and polyfluorinated alkyl substances in various matrices: How do current methods perform? *J Chromatogr A* 1216:410–421.
- Jahnke A, Huber S, Temme C, Kylin H, Berger U. 2007. Development and application of a simplified sampling method for volatile polyfluorinated alkyl substances in indoor and environmental air. *J Chromatogr A* 1164:1–9.
- Jahnke A, Barber JL, Jones KC, Temme C. 2009. Quantitative trace analysis of polyfluorinated alkyl substances (PFAS) in ambient air samples from Mace Head (Ireland): A method intercomparison. *Atmos Environ* 43:844–850.
- Jensen AA, Poulsen PB, Bossi R. 2008. Survey and environmental/health assessment of fluorinated substances in impregnated consumer products and impregnating agents. Danish Environmental Protection Agency. Survey of Chemical Substances in Consumer Products 99.
- Kaiser MA, Larsen BS, Kao C-PC, Buck RC. 2005. Vapor pressures of perfluorooctanoic, -nonanoic, -decanoic, -undecanoic, and -dodecanoic acids. *J Chem Eng Data* 50:1841–1843.
- Kaiser MA, Barton CA, Botelho M, Buck RC, Buxton LW, Gannon J, Kao CPC, Larsen BS, Russell MH, Wang N., et al. 2006. Understanding the transport of anthropogenic fluorinated compounds in the environment. *Organohalogen Compd* 68:675–678.
- Kallenborn R, Berger U, Järnberg U. 2004. Perfluorinated alkylated substances (PFAS) in the Nordic environment. Nordic Council Publication 2004:552 (ISBN 92-893-1051-0). Copenhagen (DK): Nordic Council.
- Kannan K, Corsolini S, Falandysz J, Fillmann G, Kumar KS, Loganathan BG, Mohd MA, Olivero J, Van Wouwe N, Yang JH., et al. 2004. Perfluorooctanesulfonate and related fluorochemicals in human blood from several countries. *Environ Sci Technol* 38:4489–4495.
- Kato K, Calafat AM, Needham LL. 2009. Polyfluoroalkyl chemicals in house dust. *Environ Res* 109:518–523.
- Katsushima A, Hisamoto I, Nagai M. 1964. Preparation of fluorine-containing carboxylic acids. Washington (DC): US Patent Office. Patent 3,525,758. Daikin.
- Kausch CM, Leising JE, Medsker RE, Russell VM, Thomas RR, Malik AA. 2002. Synthesis, characterization, and unusual surface activity of a series of novel architecture, water-dispersible poly(fluorooxetane)s. *Langmuir* 18:5933–5938.
- Kausch CM, Kim Y, Russell VM, Medsker RE, Thomas RR. 2003a. Surface tension and adsorption properties of a series of bolaamphiphilic poly(fluorooxetane)s. *Langmuir* 19:7182–7187.
- Kausch CM, Kim Y, Russell VM, Medsker RE, Thomas RR. 2003b. Interfacial rheological properties of a series of bolaamphiphilic poly(fluorooxetane)s. *Langmuir* 19:7354–7361.
- Kim S-K, Kannan K. 2007. Perfluorinated acids in air, rain, snow, surface runoff, and lakes: Relative importance of pathways to contamination of urban lakes. *Environ Sci Technol* 41:8328–8334.
- Kirchgeorg T, Weinberg I, Dreyer A, Ebinghaus R. 2010. Perfluorinated compounds in marine surface waters: data from the Baltic Sea and methodological challenges for future studies. *Environ Chem* 7:429–434.
- Kirchhof B, Wong D, Van Meurs J, Hilgers RD, Macek M, Lois N, Schrage NF. 2002. Use of perfluorohexyloctane as a long-term internal tamponade agent in complicated retinal detachment surgery. *Am J Ophthalmol* 133:95–101.
- Kirchner JR. 1989. Polyfluoro nitrogen-containing organic compounds. Washington (DC): US Patent Office. Patent 5,414,111. Du Pont.
- Kissa E. 1994. Fluorinated surfactants: Synthesis—Properties—Applications (Surfactant science series 50). New York (NY): Marcel Dekker. 469 p.
- Kissa E. 2001. Fluorinated surfactants and repellents (2nd edition revised and expanded) (Surfactant science series 97). New York (NY): Marcel Dekker. 640 p.
- Koch V, Knap W, Fiebig S, Geffke T, Schulze D. 2009. Biodegradation kinetic and estimated half-life of a Clariant fluorotelomer-based acrylate polymer: Results from a test on aerobic transformation in soil. *Reprod Toxicol* 27:420–421.
- Kwok KY, Taniyasu S, Yeung LWY, Murphy MB, Lam PKS, Horii Y, Kannan K, Petrick G, Sinha RK, Yamashita N. 2010. Flux of perfluorinated chemicals through wet deposition in Japan, the United States, and several other countries. *Environ Sci Technol* 44:7043–7049.
- Lange CC. 2000. The aerobic biodegradation of *N*-EtFOSE alcohol by the microbial activity present in municipal wastewater treatment sludge. 3M Company, Project CA058. Washington (DC): USEPA, docket AR226-1030a.078. Available from: OPPT.NCIC@epamail.epa.gov
- Lange CC. 2001. The 18-day aerobic biodegradation study of perfluorooctanesulfonyl-based chemistries. 3M Company, Project CA097. Washington (DC): USEPA, docket AR226-1030a.038. Available from: OPPT.NCIC@epamail.epa.gov
- Langer V, Dreyer A, Ebinghaus R. 2010. Polyfluorinated compounds in residential and nonresidential indoor air. *Environ Sci Technol* 44:8075–8081.
- Larsen BS, Kaiser MA. 2007. Challenges in perfluorocarboxylic acid measurements. *Anal Chem* 79:3966–3973.
- Larsen BS, Kaiser MA, Botelho MA, Bachmura SF, Buxton LW. 2006. Efficient “total” extraction of perfluorooctanoate from polytetrafluoroethylene fluoropolymer. *Analyst* 131:1105–1108.
- Lau C, Anitole K, Hodes C, Lai D, Pfahles-Hutchens A, Seed J. 2007. Perfluoroalkyl acids: A review of monitoring and toxicological findings. *Toxicol Sci* 99:366–394.
- Lee H, Mabury SA. 2011. A pilot survey of legacy and current commercial fluorinated chemicals in human sera from United States donors in 2009. *Environ Sci Technol* DOI: 10.1021/es200167q
- Lee H, D’eon J, Mabury SA. 2010. Biodegradation of polyfluoroalkyl phosphates as a source of perfluorinated acids to the environment. *Environ Sci Technol* 44:3305–3310.
- Lehmle H-J. 2005. Synthesis of environmentally relevant fluorinated surfactants—a review. *Chemosphere* 58:1471–1496.
- Lehmle H-J. 2009. Production of fluorinated surfactants by electrochemical fluorination. In: Zoller U, Sosis P, editors. Handbook of detergents, Part F (Surfactant science series 142). Boca Raton (FL): CRC. p 301–321.
- Liu J, Wang N, Szostek B, Buck RC, Panciroli PK, Folsom PW, Sulecki LM, Bellin CA. 2010. 6-2 Fluorotelomer alcohol aerobic biodegradation in soil and mixed bacterial culture. *Chemosphere* 78:437–444.
- Llorca M, Farré M, Picó Y, Barceló D. 2011. Analysis of perfluorinated compounds in sewage sludge by pressurized solvent extraction followed by liquid chromatography-mass spectrometry. *J Chromatogr A* 1218:4840–4846.
- Loewen M, Halldorson T, Wang F, Tomy G. 2005. Fluorotelomer carboxylic acids and PFOS in rainwater from an urban center in Canada. *Environ Sci Technol* 39:2944–2951.
- Löfstrand K, Jörundsdóttir H, Tomy G, Svavarsson J, Weihe P, Nygård T, Bergman A. 2008. Spatial trends of polyfluorinated compounds in guillemot (*Uria aalge*) eggs from North-Western Europe. *Chemosphere* 72:1475–1480.
- Loos R, Gawlik BM, Locoro G, Rimaviciute E, Contini S, Bidoglio G. 2009. EU-wide survey of polar organic persistent pollutants in European river waters. *Environ Pollut* 157:561–568.
- Loos R, Locoro G, Comero S, Contini S, Schwesig D, Werres F, Balsaa P, Gans O, Weiss S, Blaha L., et al. 2010. Pan-European survey on the occurrence of selected polar organic persistent pollutants in ground water. *Water Res* 44:4115–4126.
- Mahmoud MAM, Kärrman A, Oono S, Harada KH, Koizumi A. 2009. Polyfluorinated telomers in precipitation and surface water in an urban area of Japan. *Chemosphere* 74:467–472.
- Martin JW, Muir DCG, Moody CA, Ellis DA, Kwan WC, Solomon KR, Mabury SA. 2002. Collection of airborne fluorinated organics and analysis by gas chromatography/chemical ionization mass spectrometry. *Anal Chem* 74:584–590.
- Martin JW, Mabury SA, Solomon KR, Muir DCG. 2003a. Dietary accumulation of perfluorinated acids in juvenile rainbow trout (*Oncorhynchus mykiss*). *Environ Toxicol Chem* 22:189–195.

- Martin JW, Mabury SA, Solomon KR, Muir DCG. 2003b. Bioconcentration and tissue distribution of perfluorinated acids in rainbow trout (*Oncorhynchus mykiss*). *Environ Toxicol Chem* 22:196–204.
- Martin JW, Mabury SA, O'Brien PJ. 2005. Metabolic products and pathways of fluorotelomer alcohols in isolated rat hepatocytes. *Chem Biol Interact* 155:165–180.
- Martin JW, Ellis DA, Mabury SA, Hurley MD, Wallington TJ. 2006. Atmospheric chemistry of perfluoroalkanesulfonamides: Kinetic and product studies of the OH radical and Cl atom initiated oxidation of *N*-ethyl perfluorobutanesulfonamide. *Environ Sci Technol* 40:864–872.
- Martin JW, Asher BJ, Beesoon S, Benskin JP, Ross MS. 2010. PFOS or PreFOS? Are perfluorooctane sulfonate precursors (PreFOS) important determinants of human and environmental perfluorooctane sulfonate (PFOS) exposure? *J Environ Monit* 12:1979–2004.
- Mason Chemical. 2011. Masurf[®] FS-710 and FS-780. [cited 2011 February 8]. Available from: http://www.masonsurfactants.com/Products/Masurf_FS_710.htm
- Mertens JJWM, Sved D, Marit GB, Myers NR, Stetson PL, Murphy SR, Schmit B, Shinohara M, Farr CH. 2010. Subchronic toxicity of S-111-S-WB in Sprague Dawley rats. *Int J Toxicol* 29:358–371.
- Miljeteig C, Strøm H, Gavrilov MV, Volkov A, Jenssen BM, Gabrielsen GW. 2009. High levels of contaminants in ivory gull *Pagophila eburnea* eggs from the Russian and Norwegian Arctic. *Environ Sci Technol* 43:5521–5528.
- Millauer H. 1971. Process for the preparation of pentafluoroethyl iodide and heptafluoroisopropyl iodide. Washington (DC): US Patent Office. Patent 3,829,512. Hoechst.
- Möller A, Ahrens L, Sturm R, Westerveld J, van der Wielen F, Ebinghaus R, de Voogt P. 2010. Distribution and sources of polyfluoroalkyl substances (PFAS) in the River Rhine watershed. *Environ Pollut* 158:3243–3250.
- Mundt DJ, Mundt KA, Luippold RS, Schmidt MD, Farr CH. 2007. Clinical epidemiological study of employees exposed to surfactant blend containing perfluorononanoic acid. *Occup Environ Med* 64:589–594.
- Murakami M, Kuroda K, Sato N, Fukushima T, Takizawa S, Takada H. 2009. Groundwater pollution by perfluorinated surfactants in Tokyo. *Environ Sci Technol* 43:3840–3846.
- Murakami M, Shinohara H, Takada H. 2009. Evaluation of wastewater and street runoff as sources of perfluorinated surfactants (PFSS). *Chemosphere* 74:487–493.
- Myers AL, Mabury SA. 2010. The fate of fluorotelomer acids in a soil-water microcosm. *Environ Toxicol Chem* 29:1689–1695.
- Nabb DL, Szostek B, Himmelstein MW, Mawn MP, Gargas ML, Sweeney LM, Stadler JC, Buck RC, Fasano WJ. 2007. *In vitro* metabolism of 8-2 fluorotelomer alcohol: Interspecies comparisons and metabolic pathway refinement. *Toxicol Sci* 100:333–344.
- Nagasaki Y, Oda Y, Takahashi M. 1988. Production of fluorine-containing carboxylic acid ammonium salts. Japanese Patent 2169553. Tosoh.
- Nakayama T, Takahashi K, Matsumi Y, Toft A, Sulbaek Andersen MP, Nielsen OJ, Waterland RL, Buck RC, Hurley MD, Wallington TJ. 2007. Atmospheric chemistry of $\text{CF}_3\text{CH}=\text{CH}_2$ and $\text{C}_4\text{F}_9\text{CH}=\text{CH}_2$: Products of the gas-phase reactions with Cl atoms and OH radicals. *J Phys Chem A* 111:909–915.
- Napoli M. 1996. Diblock and triblock semifluorinated *n*-alkanes: preparations, structural aspects and applications. *J Fluorine Chem* 79:59–69.
- Nguyen VT, Reinhard M, Karina GY-H. 2011. Occurrence and source characterization of perfluorochemicals in an urban watershed. *Chemosphere* 82:1277–1285.
- Nilsson H, Kärrman A, Rotander A, van Bavel B, Lindström G, Westberg H. 2010. Inhalation exposure to fluorotelomer alcohols yield perfluorocarboxylates in human blood? *Environ Sci Technol* 44:7717–7722.
- Norwegian Pollution Control Authority. 2008. Screening of polyfluorinated organic compounds at four fire training facilities in Norway Report 2444. Oslo (NO): Norwegian Pollution Control Authority.
- Oakes KD, Benskin JP, Martin JW, Ings JS, Heinrichs JY, Dixon DG, Servos MR. 2010. Biomonitoring of perfluorochemicals and toxicity to the downstream fish community of Etobicoke Creek following deployment of aqueous film-forming foam. *Aquat Toxicol* 98:120–129.
- [OECD] Organisation for Economic Co-operation and Development. 2002. Hazard assessment of perfluorooctane sulfonate (PFOS) and its salts. Report ENV/JM/RD(2002)17/FINAL.
- [OECD] Organisation for Economic Co-operation and Development. 2007. Lists of PFOS, PFAS, PFOA, PFCA, related compounds and chemicals that may degrade to PFCA. Report ENV/JM/MONO(2006)15.
- [OECD] Organisation for Economic Co-operation and Development. 2011. OECD portal on perfluorinated chemicals. [cited 2011 February 8]. Available from: http://www.oecd.org/site/0,3407,en_21571361_44787844_1_1_1_1_1,00.html
- Olsen GW, Burris JM, Lundberg JK, Hansen KJ, Mandel JH, Zobel LR. 2002. Identification of fluorochemicals in human sera: III. Pediatric participants in a group A streptococci clinical trial investigation (3M Company, Medical Department, Epidemiology, St. Paul, MN) USEPA public docket AR-226-1085.
- Olsen GW, Huang HY, Helzlsouer KJ, Hansen KJ, Butenhoff JL, Mandel JH. 2005. Historical comparison of perfluorooctanesulfonate, perfluorooctanoate, and other fluorochemicals in human blood. *Environ Health Perspect* 113:539–545.
- Olsen GW, Mair DC, Church TR, Ellefson ME, Reagen WK, Boyd TM, Herron RM, Medhizadehkashi Z, Nobiletti JB, Rios JA, et al. 2008. Decline in perfluorooctanesulfonate and other polyfluoroalkyl chemicals in American Red Cross adult blood donors, 2000–2006. *Environ Sci Technol* 42:4989–4995.
- Olsen GW, Chang S-C, Noker PE, Gorman GS, Ehresman DJ, Lieder PH, Butenhoff JL. 2009. A comparison of the pharmacokinetics of perfluorobutanesulfonate (PFBS) in rats, monkeys, and humans. *Toxicology* 256:65–74.
- Omnova Solutions. 2011. PolyFox—A novel fluorochemical. [cited 2011 February 8]. Available from: <http://www.omnova.com/products/chemicals/PolyFox.aspx>
- Oono S, Harada KH, Mahmoud MAM, Inoue K, Koizumi A. 2008. Current levels of airborne polyfluorinated telomers in Japan. *Chemosphere* 73:932–937.
- Oono S, Matsubara E, Harada KH, Takagi S, Hamada S, Asakawa A, Inoue K, Watanabe I, Koizumi A. 2008. Survey of airborne polyfluorinated telomers in Keihan area, Japan. *Bull Environ Contam Toxicol* 80:102–106.
- Ostertag SK, Chan HM, Moisey J, Dabeka R, Tittlemier SA. 2009. Historic dietary exposure to perfluorooctane sulfonate, perfluorinated carboxylates, and fluorotelomer unsaturated carboxylates from the consumption of store-bought and restaurant foods for the Canadian population. *J Agric Food Chem* 57:8534–8544.
- Paul AG, Jones KC, Sweetman AJ. 2009. A first global production, emission, and environmental inventory for perfluorooctane sulfonate. *Environ Sci Technol* 43:386–392.
- Peng H, Wei Q, Wan Y, Giesy JP, Li L, Hu J. 2010. Tissue distribution and maternal transfer of poly- and perfluorinated compounds in Chinese sturgeon (*Acipenser sinensis*): Implications for reproductive risk. *Environ Sci Technol* 44:1868–1874.
- Peschka M, Fichtner N, Hierse W, Kirsch P, Montenegro E, Seidel M, Wilken RD, Knepper TP. 2008. Synthesis and analytical follow-up of the mineralization of a new fluorosurfactant prototype. *Chemosphere* 72:1534–1540.
- Piekarz AM, Primbs T, Field JA, Barofsky DF, Simonich S. 2007. Semivolatile fluorinated organic compounds in Asian and Western U.S. air masses. *Environ Sci Technol* 41:8248–8255.
- Pistocchi A, Loos R. 2009. A map of European emissions and concentrations of PFOS and PFOA. *Environ Sci Technol* 43:9237–9244.
- Plassmann MM, Berger U. 2010. Trace analytical methods for semifluorinated *n*-alkanes in snow, soil, and air. *Anal Chem* 82:4551–4557.
- Plumlee MH, McNeill K, Reinhard M. 2009. Indirect photolysis of perfluorochemicals: Hydroxyl radical-initiated oxidation of *N*-ethyl perfluorooctane sulfonamido acetate (*N*-EtFOSAA) and other perfluoroalkanesulfonamides. *Environ Sci Technol* 43:3662–3668.
- Powley CR, George SW, Russell MH, Hoke RA, Buck RC. 2008. Polyfluorinated chemicals in a spatially and temporally integrated food web in the Western Arctic. *Chemosphere* 70:664–672.
- Prevedouros K, Cousins IT, Buck RC, Korzeniowski SH. 2006. Sources, fate and transport of perfluorocarboxylates. *Environ Sci Technol* 40:32–44.
- Rao NS, Baker BE. 1994. Textile finishes and fluorosurfactants. In: Banks RE, Smart BE, Tatlow JC, editors. Organofluorine chemistry: Principles and commercial applications. New York (NY): Plenum. p 321–338.

- Rayne S, Forest K. 2009a. A new class of perfluorinated acid contaminants: Primary and secondary substituted perfluoroalkyl sulfonamides are acidic at environmentally and toxicologically relevant pH values. *J Environ Sci Health A* 44:1388–1399.
- Rayne S, Forest K. 2009b. Perfluoroalkyl sulfonic and carboxylic acids: A critical review of physicochemical properties, levels and patterns in waters and wastewaters, and treatment methods. *J Environ Sci Health A* 44:1145–1149.
- Rayne S, Forest K. 2010a. Theoretical studies on the pK_a values of perfluoroalkyl carboxylic acids. *J Mol Struct THEOCHEM* 949:60–69.
- Rayne S, Forest K. 2010b. Aqueous phase hydration and hydrate acidity of perfluoroalkyl and $n:2$ fluorotelomer aldehydes. *Nature Precedings* 8 June 2010.
- Rayne S, Forest K. 2010c. Modeling the hydrolysis of perfluorinated compounds containing carboxylic and phosphoric acid ester functions and sulfonamide groups. *J Environ Sci Health A* 45:432–446.
- Rayne S, Forest K, Friesen KJ. 2008. Congener-specific numbering systems for the environmentally relevant C_4 through C_8 perfluorinated homologue groups of alkyl sulfonates, carboxylates, telomer alcohols, olefins, and acids, and their derivatives. *J Environ Sci Health A* 43:1391–1401.
- Reagen WK, Lindstrom KR, Jacoby CB, Purcell RG, Kestner TA, Payfer RM, Miller JW. 2007. Environmental characterization of 3M electrochemical fluorination derived perfluorooctanoate and perfluorooctane sulfonate. Abstracts of the Society of Environmental Toxicology and Chemistry North America 28th Annual Meeting; 2007 November 11–15; Milwaukee, Wisconsin, USA.
- Renner R. 2006. The long and the short of perfluorinated replacements. *Environ Sci Technol* 40:12–13.
- Rhoads KR, Janssen EM-L, Luthy RG, Criddle CS. 2008. Aerobic biotransformation and fate of N -ethyl perfluorooctane sulfonamidoethanol (N -EtFOSE) in activated sludge. *Environ Sci Technol* 42:2873–2878.
- Riddell N, Arsenault G, Benskin JP, Chittim B, Martin JW, McAlees A, McCrindle R. 2009. Branched perfluorooctane sulfonate isomer quantification and characterization in blood serum samples by HPLC/ESI-MS/MS. *Environ Sci Technol* 43:7902–7908.
- Ritter SK. 2010. Fluorochemicals go short. *Chem Eng News* 88:12–17.
- Rogowski I, Leonard D, Gauthier J-Y, Lanteri P. 2007. Influence of fluorine-based additive content on the physical and physicochemical properties of ski gliding wax. *Cold Reg Sci Technol* 49:145–150.
- Ruan T, Wang Y, Wang T, Zhang Q, Ding L, Liu J, Wang C, Qu G, Jiang G. 2010. Presence and partitioning behavior of polyfluorinated iodine alkanes in environmental matrices around a fluorochemical manufacturing plant: Another possible source for perfluorinated carboxylic acids? *Environ Sci Technol* 44:5755–5761.
- Russell MH, Berti WR, Szostek B, Buck RC. 2008. Investigation of the biodegradation potential of a fluoroacrylate polymer product in aerobic soils. *Environ Sci Technol* 42:800–807.
- Russell MH, Wang N, Berti WR, Szostek B, Buck RC. 2009. Comment on “Degradability of an acrylate-linked, fluorotelomer polymer in soil.” *Environ Sci Technol* 44:848–848.
- Russell MH, Berti WR, Szostek B, Wang N, Buck RC. 2010. Evaluation of PFO formation from the biodegradation of a fluorotelomer-based urethane polymer product in aerobic soils. *Polym Degrad Stab* 95:79–85.
- Schenker U, Scheringer M, MacLeod M, Martin JW, Cousins IT, Hungerbühler K. 2008. Contribution of volatile precursor substances to the flux of perfluorooctanoate to the Arctic. *Environ Sci Technol* 42:3710–3716.
- Schultz MM, Barofsky DF, Field JA. 2004. Quantitative determination of fluorotelomer sulfonates in groundwater by LC MS/MS. *Environ Sci Technol* 38:1828–1835.
- Schulze P-E, Norin H. 2006. Fluorinated pollutants in all-weather clothing Report 2/ 2006. Oslo (NO): Friends of the Earth Norway.
- Scott BF, Spencer C, Mabury SA, Muir DCG. 2006. Poly and perfluorinated carboxylates in North American precipitation. *Environ Sci Technol* 40:7167–7174.
- Scott BF, De Silva AO, Spencer C, Lopez E, Backus SM, Muir DCG. 2010. Perfluoroalkyl acids in Lake Superior water: Trends and sources. *J Great Lakes Res* 36:277–284.
- Seacat AM, Thomford PJ, Hansen KJ, Clemen LA, Eldridge SR, Elcombe CR, Butenhoff JL. 2003. Sub-chronic dietary toxicity of potassium perfluorooctanesulfonate in rats. *Toxicology* 183:117–131.
- Sepulveda JG, Blaine AC, Hundal LS, Higgins CP. 2011. Occurrence and fate of perfluorochemicals in soil following the land application of municipal biosolids. *Environ Sci Technol* DOI: 10.1021/es103903d
- Shoeib M, Harner T, Wilford BH, Jones KC, Zhu J. 2005. Perfluorinated sulfonamides in indoor and outdoor air and indoor dust: Occurrence, partitioning, and human exposure. *Environ Sci Technol* 39:6599–6606.
- Shoeib M, Vlahos P, Harner T, Peters A, Graustein M, Narayan J. 2010. Survey of polyfluorinated chemicals (PFCs) in the atmosphere over the northeast Atlantic Ocean. *Atmos Environ* 44:2887–2893.
- Shoeib M, Harner T, Webster GM, Lee SC. 2011. Indoor sources of poly- and perfluorinated compounds (PFCs) in Vancouver, Canada: Implications for human exposure. *Environ Sci Technol* DOI: 10.1021/es103562v
- Sianesi D, Marchionni G, De Pasquale RJ. 1994. Perfluoropolyethers (PFPEs) from perfluoroolefin photooxidation: Fomblin[®] and Galden[®] fluids. In: Banks RE, Smart BE, Tatlow JC, editors. Organofluorine chemistry: Principles and commercial applications. New York (NY): Plenum. p 431–462.
- Sinclair E, Kannan K. 2006. Mass loading and fate of perfluoroalkyl surfactants in wastewater treatment plants. *Environ Sci Technol* 40:1408–1414.
- Smart BE. 1994. Characteristics of C-F systems. In: Banks RE, Smart BE, Tatlow JC, editors. Organofluorine chemistry: Principles and commercial applications. New York (NY): Plenum. p 57–88.
- So MK, Yamashita N, Taniyasu S, Jiang Q, Giesy JP, Chen K, Lam PKS. 2006. Health risks in infants associated with exposure to perfluorinated compounds in human breast milk from Zhoushan, China. *Environ Sci Technol* 40:2924–2929.
- So MK, Miyake Y, Yeung WY, Ho YM, Taniyasu S, Rostkowski P, Yamashita N, Zhou BS, Shi XJ, Wang JX., et al. 2007. Perfluorinated compounds in the Pearl River and Yangtze River of China. *Chemosphere* 68:2085–2095.
- Solomon S, Burkholder JB, Ravishankara AR, Garcia RR. 1994. Ozone depletion and global warming potentials of CF_3I . *J Geophys Res D* 99:20929–20935.
- Solvay Solexis. 2011. Fluorolink[®] and Solvera[®] products. [cited 2011 February 8]. Available from: http://www.solvaysites.com/sites/solvayplastics/EN/specialty_polymers/Fluorinated_Fluids/Pages/Fluorinated_Fluids.aspx
- Stemmler I, Lammel G. 2010. Pathways of PFOA to the Arctic: variabilities and contributions of oceanic currents and atmospheric transport and chemistry sources. *Atmos Chem Phys* 10:9965–9980.
- Stock NL, Furdul VI, Muir DCG, Mabury SA. 2007. Perfluoroalkyl contaminants in the Canadian Arctic: Evidence of atmospheric transport and local contamination. *Environ Sci Technol* 41:3529–3536.
- Strynar MJ, Lindstrom AB. 2008. Perfluorinated compounds in house dust from Ohio and North Carolina, USA. *Environ Sci Technol* 42:3751–3756.
- Stump DG, Holson JF, Murphy SR, Farr CH, Schmit B, Shinohara M. 2008. An oral two-generation reproductive toxicity study of S-111-S-WB in rats. *Reprod Toxicol* 25:7–20.
- Sturm R, Ahrens L. 2010. Trends of polyfluoroalkyl compounds in marine biota and in humans. *Environ Chem* 7:457–484.
- Taniyasu S, Kannan K, So MK, Gulkowska A, Sinclair E, Okazawa T, Yamashita N. 2005. Analysis of fluorotelomer alcohols, fluorotelomer acids, and short- and long-chain perfluorinated acids in water and biota. *J Chromatogr A* 1093:89–97.
- Taniyasu S, Kannan K, Yeung LWY, Kwok KY, Lam PKS, Yamashita N. 2008. Analysis of trifluoroacetic acid and other short-chain perfluorinated acids (C_2 – C_4) in precipitation by liquid chromatography–tandem mass spectrometry: Comparison to patterns of long-chain perfluorinated acids (C_5 – C_{18}). *Anal Chim Acta* 619:221–230.
- Taves DR. 1968. Evidence that there are two forms of fluoride in human serum. *Nature* 217:1050–1051.
- Taylor CK. 1999. Fluorinated surfactants in practice. In: Karsa D, editor. Design and selection of performance surfactants: Annual surfactants review. New York (NY): John Wiley & Sons. p 271–316.
- Thomas RR. 2006. Fluorinated surfactants. In: Farn RJ, editor. Chemistry and technology of surfactants. Oxford (UK): Blackwell. p 227–235.

- Tittlemier SA, Pepper K, Edwards L, Tomy G. 2005. Development and characterization of a solvent extraction-gas chromatographic/mass spectrometric method for the analysis of perfluorooctanesulfonamide compounds in solid matrices. *J Chromatogr A* 1066:189–195.
- Tittlemier SA, Pepper K, Edwards L. 2006. Concentrations of perfluorooctanesulfonamides in Canadian Total Diet Study composite food samples collected between 1992 and 2004. *J Agric Food Chem* 54:8385–8389.
- Toms L-ML, Calafat AM, Kato K, Thompson J, Harden F, Hobson P, Sjödin A, Mueller JF. 2009. Polyfluoroalkyl chemicals in pooled blood serum from infants, children, and adults in Australia. *Environ Sci Technol* 43:4194–4199.
- Tomy GT, Budakowski W, Halldorson T, Helm PA, Stern GA, Friesen K, Pepper K, Tittlemier SA, Fisk AT. 2004. Fluorinated organic compounds in an Eastern Arctic marine food web. *Environ Sci Technol* 38:6475–6481.
- Tomy GT, Tittlemier SA, Palace VP, Budakowski WR, Braekevelt E, Brinkworth L, Friesen K. 2004. Biotransformation of *N*-ethyl perfluorooctanesulfonamide by rainbow trout (*Oncorhynchus mykiss*) liver microsomes. *Environ Sci Technol* 38:758–762.
- Trier X, Granby K, Christensen J. 2011. Polyfluorinated surfactants (PFS) in paper and board coatings for food packaging. *Environ Sci Pollut Res* 18:1108–1120.
- Trier X, Nielsen NJ, Christensen JH. 2011. Structural isomers of polyfluorinated di- and tri-alkylated phosphate ester surfactants present in industrial blends and in microwave popcorn bags. *Environ Sci Pollut Res* 18:1422–1432.
- Tsuda N, Sawachi C, Sawada Y, Shimizu T. 2003. Process for producing aqueous fluoropolymer dispersion and aqueous fluoropolymer dispersion. Washington (DC): US Patent Office. Patent 7,714,047. Daikin Industries.
- Ubel FA, Sorenson SD, Roach DE. 1980. Health status of plant workers exposed to fluorochemicals: A preliminary report. *Am Ind Hyg Assoc J* 41:584–589.
- Ukhashi H, Hayashi T, Takasaki Y. 1977. Process for producing perfluorocarboxylic acid. British Patent 1473807. Asahi Glass.
- [UNEP] UN Environment Programme. 2009. Governments unite to step-up reduction on global DDT reliance and add nine new chemicals under international treaty. [cited 2010 December 11]. Available from: <http://chm.pops.int/Convention/Pressrelease/COP4Geneva8May2009/tabid/542/language/en-US/Default.aspx>
- United Nations. 1998. Kyoto Protocol to the United Nations Framework Convention on Climate Change. [cited 2010 September 28]. Available from: <http://unfccc.int/resource/docs/convkp/kpeng.pdf>
- [USEPA] US Environmental Protection Agency. 2000. EPA and 3M announce phase-out of PFOS. [cited 2011 January 9]. Available from: <http://yosemite.epa.gov/opa/admpress.nsf/0/33aa946e6cb11f35852568e1005246b4>
- [USEPA] US Environmental Protection Agency. 2006a. Meeting with Registrant on perfluorinated chemical (Masurf-FS-780, Fluowet PL 80-B and Fluowet PL-80) use as inerts. USEPA Administrative Record AR226-3804. Available from: OPPT.NCIC@epamail.epa.gov
- [USEPA] US Environmental Protection Agency. 2006b. 2010/2015 PFOA Stewardship Program. [cited 2011 9 January]. Available from: <http://www.epa.gov/opptintr/pfoa/pubs/stewardship/>
- [USEPA] US Environmental Protection Agency. 2009. Long-chain perfluorinated chemicals (PFCs) action plan. [cited 2010 December 11]. Available from: http://www.epa.gov/opptintr/existingchemicals/pubs/pfcs_action_plan1230_09.pdf
- [USEPA] US Environmental Protection Agency. 2010. Perfluorooctanoic Acid (PFOA) and Fluorinated Telomers - Related EPA Actions - PFOA-related research. [cited 2010 December 11]. Available from: <http://www.epa.gov/opptintr/pfoa/pubs/activities.html#ord>
- van der Putte I, Murin M, van Velthoven M, Affourtit F. 2010. Analysis of the risks arising from the industrial use of perfluorooctanoic acid (PFOA) and ammonium perfluorooctanoate (APFO) and from their use in consumer articles. Delft (NL): RPS Advies.
- van Leeuwen SPJ, de Boer J. 2007. Extraction and clean-up strategies for the analysis of poly- and perfluoroalkyl substances in environmental and human matrices. *J Chromatogr A* 1153:172–185.
- Vésine E, Bossoutrot V, Mellouki A, Le Bras G, Wenger J, Sidebottom H. 2000. Kinetic and mechanistic study of OH- and Cl-initiated oxidation of two unsaturated HFCs: C₄F₉CH=CH₂ and C₆F₁₃CH=CH₂. *J Phys Chem A* 104: 8512–8520.
- Visca M, Chittofrati A, Kapeliouchko V, Malvasi M. 2003. Aqueous dispersions of fluoropolymers. European Patent 1334996. Solvay Solexis.
- Wallington TJ, Hurley MD, Xia J, Wuebbles DJ, Sillman S, Ito A, Penner JE, Ellis DA, Martin J, Mabury SA., et al. 2006. Formation of C₇F₁₅COOH (PFOA) and other perfluorocarboxylic acids during the atmospheric oxidation of 8:2 fluorotelomer alcohol. *Environ Sci Technol* 40:924–930.
- Wang N, Szostek B, Folsom PW, Sulecki LM, Capka V, Buck RC, Berti WR, Gannon JT. 2005. Aerobic biotransformation of ¹⁴C-labeled 8-2 telomer B alcohol by activated sludge from a domestic sewage treatment plant. *Environ Sci Technol* 39:531–538.
- Wang N, Szostek B, Buck RC, Folsom PW, Sulecki LM, Gannon JT. 2009. 8-2 Fluorotelomer alcohol aerobic soil biodegradation: Pathways, metabolites, and metabolite yields. *Chemosphere* 75:1089–1096.
- Wang N, Liu J, Buck RC, Korzeniowski SH, Wolstenholme BW, Folsom PW, Sulecki LM. 2011. 6:2 Fluorotelomer sulfonate aerobic biotransformation in activated sludge of waste water treatment plants. *Chemosphere* 82:853–858.
- Wania F. 2007. A global mass balance analysis of the source of perfluorocarboxylic acids in the Arctic Ocean. *Environ Sci Technol* 41:4529–4535.
- Washburn ST, Bingman TS, Braithwaite SK, Buck RC, Buxton LW, Clewell HJ, Haroun LA, Kester JE, Rickard RW, Shipp AM. 2005. Exposure assessment and risk characterization for perfluorooctanoate in selected consumer articles. *Environ Sci Technol* 39:3904–3910.
- Washington JW, Ellington JJ, Jenkins TM, Evans JJ, Yoo H, Hafner SC. 2009a. Degradability of an acrylate-linked, fluorotelomer polymer in soil. *Environ Sci Technol* 43:6617–6623.
- Washington JW, Ellington JJ, Jenkins TM, Yoo H. 2009b. Response to comments on “Degradability of an acrylate-linked, fluorotelomer polymer in soil.” *Environ Sci Technol* 44:849–850.
- Webster E, Ellis DA. 2010. The potential role of sea spray generation in the atmospheric transport of perfluorocarboxylic acids. *Environ Toxicol Chem* 29:1703–1708.
- Webster E, Ellis DA, Reid LK. 2010. Modeling the environmental fate of perfluorooctanoic acid and perfluorooctanoate: An investigation of the role of individual species partitioning. *Environ Toxicol Chem* 29:1466–1475.
- Weihe P, Kato K, Calafat AM, Nielsen F, Wanigatunga AA, Needham LL, Grandjean P. 2008. Serum concentrations of polyfluoroalkyl compounds in Faroese whale meat consumers. *Environ Sci Technol* 42:6291–6295.
- Weinberg I, Dreyer A, Ebinghaus R. 2011a. Landfills as sources of polyfluorinated compounds, polybrominated diphenyl ethers and musk fragrances to ambient air. *Atmos Environ* 45:935–941.
- Weinberg I, Dreyer A, Ebinghaus R. 2011b. Waste water treatment plants as sources of polyfluorinated compounds, polybrominated diphenyl ethers and musk fragrances to ambient air. *Environ Pollut* 159:125–132.
- Xie W, Wu Q, Kania-Korwel I, Tharappel J, Telu S, Coleman M, Glauert H, Kannan K, Mariappan S, Spitz D., et al. 2009. Subacute exposure to *N*-ethyl perfluorooctanesulfonamidoethanol results in the formation of perfluorooctanesulfonate and alters superoxide dismutase activity in female rats. *Arch Toxicol* 83:909–924.
- Xu L, Krenitsky DM, Seacat AM, Butenhoff JL, Anders MW. 2004. Biotransformation of *N*-ethyl-*N*-(2-hydroxyethyl)perfluorooctanesulfonamide by rat liver microsomes, cytosol, and slices and by expressed rat and human cytochromes P450. *Chem Res Toxicol* 17:767–775.
- Xu L, Krenitsky DM, Seacat AM, Butenhoff JL, Tephly TR, Anders MW. 2006. *N*-Glucuronidation of perfluorooctanesulfonamide by human, rat, dog, and monkey liver microsomes and by expressed rat and human UDP-glucuronosyl transferases. *Drug Metab Dispos* 34:1406–1410.
- Yarwood G, Kemball-Cook S, Keinath M, Waterland RL, Korzeniowski SH, Buck RC, Russell MH, Washburn ST. 2007. High-resolution atmospheric modeling of fluorotelomer alcohols and perfluorocarboxylic acids in the North American troposphere. *Environ Sci Technol* 41:5756–5762.
- Yeung LWY, Miyake Y, Wang Y, Taniyasu S, Yamashita N, Lam PKS. 2009. Total fluorine, extractable organic fluorine, perfluorooctane sulfonate and other related fluorochemicals in liver of Indo-Pacific humpback dolphins (*Sousa chinensis*) and finless porpoises (*Neophocaena phocaenoides*) from South China. *Environ Pollut* 157:17–23.

- Yoo H, Yamashita N, Taniyasu S, Lee KT, Jones PD, Newsted JL, Khim JS, Giesy JP. 2009. Perfluoroalkyl acids in marine organisms from Lake Shihwa, Korea. *Arch Environ Contam Toxicol* 57:552–560.
- Yoo H, Washington JW, Ellington JJ, Jenkins TM, Neill MP. 2010. Concentrations, distribution, and persistence of fluorotelomer alcohols in sludge-applied soils near Decatur, Alabama, USA. *Environ Sci Technol* 44:8397–8402.
- Yoo H, Washington JW, Jenkins TM, Ellington JJ. 2011. Quantitative determination of perfluorochemicals and fluorotelomer alcohols in plants from biosolid-amended fields using LC/MS/MS and GC/MS. *Environ Sci Technol* DOI: 10.1021/es102972m
- Young CJ, Mabury SA. 2010. Atmospheric perfluorinated acid precursors: chemistry, occurrence, and impacts. *Rev Environ Contam Toxicol* 208:1–109.
- Young CJ, Hurley MD, Wallington TJ, Mabury SA. 2008. Atmospheric chemistry of 4:2 fluorotelomer iodide ($n\text{-C}_4\text{F}_9\text{CH}_2\text{CH}_2\text{I}$): Kinetics and products of photolysis and reaction with OH radicals and Cl atoms. *J Phys Chem A* 112:13542–13548.
- Yue R. 2008. Additional information on production and use of PFOS (30 April 2008). [cited 2011 May 29]. Available from: http://chm.pops.int/Portals/0/Repository/addinfo_2008/UNEP-POPS-POPRC-SUB-F08-PFOS-ADIN-CHI.English.pdf
- Zhang T, Sun H, Gerecke AC, Kannan K, Müller CE, Alder AC. 2010. Comparison of two extraction methods for the analysis of per- and polyfluorinated chemicals in digested sewage sludge. *J Chromatogr A* 1217: 5026–5034.
- Zushi Y, Tamada M, Kanai Y, Masunaga S. 2010. Time trends of perfluorinated compounds from the sediment core of Tokyo Bay, Japan (1950s–2004). *Environ Pollut* 158:756–763.
- Zushi Y, Ye F, Motegi M, Nojiri K, Hosono S, Suzuki T, Kosugi Y, Yaguchi K, Masunaga S. 2011. Spatially detailed survey on pollution by multiple perfluorinated compounds in the Tokyo Bay basin of Japan. *Environ Sci Technol* 45:2887–2893.