



LUND UNIVERSITY

Preoperative Anxiety and Depression Correlate With Dissatisfaction After Total Knee Arthroplasty

A Prospective Longitudinal Cohort Study of 186 Patients, With 4-Year Follow-Up

Ali, Abdulemir; Lindstrand, Anders; Sundberg, Martin; Flivik, Gunnar

Published in:
Journal of Arthroplasty

DOI:
[10.1016/j.arth.2016.08.033](https://doi.org/10.1016/j.arth.2016.08.033)

2017

Document Version:
Peer reviewed version (aka post-print)

[Link to publication](#)

Citation for published version (APA):

Ali, A., Lindstrand, A., Sundberg, M., & Flivik, G. (2017). Preoperative Anxiety and Depression Correlate With Dissatisfaction After Total Knee Arthroplasty: A Prospective Longitudinal Cohort Study of 186 Patients, With 4-Year Follow-Up. *Journal of Arthroplasty*, 32(3), 767-770. <https://doi.org/10.1016/j.arth.2016.08.033>

Total number of authors:
4

Creative Commons License:
CC BY-NC-ND

General rights

Unless other specific re-use rights are stated the following general rights apply:
Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Read more about Creative commons licenses: <https://creativecommons.org/licenses/>

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LUND UNIVERSITY

PO Box 117
221 00 Lund
+46 46-222 00 00

1

2

3

4 **Preoperative Anxiety and Depression Correlate With Dissatisfaction After Total Knee**
5 **Arthroplasty: A Prospective Longitudinal Cohort Study of 186 patients, With 4-year**
6 **Follow-Up.**

7

8

9

10

11

12 Abdulemir Ali, MD , Anders Lindstrand, MD , PhD, Martin Sundberg, MD , PhD, Gunnar
13 Flivik, MD , PhD

14 Department of Orthopaedics, Skåne University Hospital, Clinical Sciences, Lund University,
15 Lund, Sweden

16 Keywords: TKA, dissatisfaction, HAD, KOOS, VAS pain.

17

18

19

20 Correspondence: abdulemir.ali@med.lu.se

21 **Abstract**

22 **Background:**

23 After more than 4 decades experience of total knee arthroplasty (TKA) there is still a group of
24 patients who are not satisfied with the outcome. In spite of the improvement of many aspects
25 around the procedure, for unexplainable reasons patient dissatisfaction is still approximately
26 the same. We conducted this study to analyse correlations between preoperative psychological
27 aspects and dissatisfaction after TKA.

28 **Methods:**

29 A total of 186 patients were operated with a primary TKA. Patients filled out the Hospital
30 Anxiety and Depression Scale, Visual Analogue Pain Scale (0-100) and Knee injury and
31 Osteoarthritis Outcome Score preoperatively and 4 years postoperatively. Four years
32 postoperatively the patients also scored their satisfaction degree with the outcome of the
33 surgery.

34 **Results:**

35 Of 186 patients, 27 (15%) reported that they were dissatisfied or uncertain with the result of
36 their TKA 4 years postoperatively. Sixteen of those 27 patients had reported anxiety /
37 depression preoperatively compared to 11 of 159 (7%) in the satisfied or very satisfied
38 groups. Patients with preoperative anxiety or depression had more than 6 times higher risk to
39 be dissatisfied compared to patients with no anxiety or depression (p-value < 0.001). Patients
40 with deep prosthetic infection had 3 times higher risk to be dissatisfied with the operation
41 outcome (p-value = 0.03). Dissatisfied patients had 1-day longer hospital stay compared to the
42 satisfied group (p-value < 0.001).

43 **Conclusion:**

44 Preoperative anxiety and/or depression is an import predictor for dissatisfaction after TKA.
45 Psychological assessment and treatment preoperatively might improve degree of satisfaction.

46

47 **Introduction:**

48 Dissatisfaction after unrevised TKA has historically been between 6% and 14% [1-12] and if
49 one adds the group uncertain to the dissatisfied group, it can be up to 28% [13]. There are
50 several explainable reasons of poor outcome after TKA like patella related problems,
51 infection, stiffness, instability, periprosthetic fracture, tendon rapture, loosening and nerve
52 injury. Other well-known extra-articular reasons of poor outcome like hip, spine, vascular
53 disease, or chronic regional pain syndrome might be contributing [14]. There are studies of
54 joint arthroplasties that suggest that factors not primarily related to structural tissue changes
55 but rather of psychological nature are involved [13, 15-17]. In our previous study from the
56 Swedish Knee Arthroplasty Register we found that patients, who were dissatisfied, had
57 similar performance tests, clinical and radiographic findings as compared with those who
58 were very satisfied. The patients who reported poor response after TKA were unhappy, as
59 demonstrated by Visual Analog Pain Scale (VAS) pain and Hospital Anxiety and Depression
60 scale (HAD), despite the absence of a discernible objective reason for revision [18]. A
61 limitation of the study was the absence of preoperative data regarding psychological
62 assessment of patients. Based on this we designed the present study to better analyze
63 correlations between preoperative psychological factors and dissatisfaction.

64

65

66

67 **Patients and methods:**

68 A total of 186 consecutive TKA patients having primary osteoarthritis were included. We
69 excluded patients having bilateral TKA, dementia, or not being able to speak Swedish. The
70 patients in this study were included in another study regarding the effect of continuous
71 intraarticular analgesia on pain and rehabilitation after total knee arthroplasty [19]. Total
72 number of patients included in that original article were 200, and 14 of these patients were
73 lost during the follow-up period (12 deceased, 2 refused to participate). The patients were
74 operated on between January 2010 and April 2011. All patients had a standard straight central
75 skin incision, medial parapatellar arthrotomy, and preparation of femur and tibia according to
76 the instructions of the prosthesis manufacturer. Patients received either the cruciate retaining
77 Triathlon knee (Stryker, United Kingdom) in 151 patients or the cruciate retaining PFC knee
78 (DePuy, United Kingdom) in 35 patients, depending on the surgeon's preference. Five
79 orthopedic surgeons who were subspecialized in arthroplasty performed the surgeries. Spinal
80 anaesthesia was used as a standard method (87%), while the remaining patients received
81 general anaesthesia. Premedication and postoperative analgesia were standardized. Patients
82 filled out the HAD [20], VAS pain (0–100 mm, where 0 = no pain and 100 = intolerable
83 pain), Knee injury and Osteoarthritis Outcome Score (KOOS) questionnaire (0-100, where 0
84 = major problem and 100 = no problem) [21], and pain drawing with predefined body region
85 to identify patients with chronic widespread pain [22] preoperatively, and 4 years
86 postoperatively. Four years postoperatively patients also filled out their satisfaction degree
87 regarding the operated knee as very satisfied, satisfied, uncertain or dissatisfied [10]. For
88 statistical analysis reasons, we merged the satisfaction degree data to 2 groups: satisfied (very
89 satisfied and satisfied groups) and not satisfied (uncertain and dissatisfied groups). Active
90 range of motion (ROM) of the knee (goniometry) was measured preoperatively; VAS pain,
91 analgesic consumption, and wound-healing complications were also recorded. Preoperative

92 radiographic assessment was done according to Kellgren and Lawrence system for
93 classification of osteoarthritis. The patient's files regarding complications and reoperation
94 were checked 4 years postoperatively in a complications registry and patient files.

95

96 **Statistics:**

97 A Cox multiple regression analysis with constant follow up and robust variance estimation
98 [23] was used to study relative risks for categorical variables among the dissatisfied group.
99 Regarding continuous variables, like the mean difference between 2 groups, they were
100 analyzed by the analysis of covariance method. In both methods, patients' gender, age and
101 body mass index preoperatively and at 4 years postoperatively were included. A p-value of <
102 0.05 was considered to be statistically significant. A power analysis had been performed for
103 the original article [19], which estimated that 200 patients were sufficient to find differences
104 between the 2 groups for that study, and our statistician considered it to be valid also for the
105 outcome in this study. Statistical analyses were performed using the Stata 12.0 program.

106 **Ethics:**

107 The study and study registration was performed in compliance with the Helsinki Declaration,
108 and all patients had given their informed written consent. The ethics committee of the Faculty
109 of Medicine, Lund University, approved the study (Dnr 2009/368). This is the same
110 approvement number as the original study about continuous intraarticular analgesia [19].

111

112

113 **Results:**

114 Patient characteristics and the overall result are shown in table 1-3. A total of 27 of the 186
115 patients (15%) reported that they were not satisfied (uncertain or dissatisfied) with the result
116 of their TKA 4 years after surgery. A total of 16 of those 27 patients reported anxiety and/or
117 depression according to the HAD score compared with 11 of 159 (7%) in the satisfied group
118 (satisfied and very satisfied) at the 4-year follow-up. As shown in table 2, we found that the
119 patients who preoperatively had anxiety/depression had more than 6 times higher risk to be
120 dissatisfied after TKA as compared with patients without preoperative anxiety/depression.
121 Mean length of stay in hospital at the time of surgery for the group that was dissatisfied at 4-
122 year follow-up was 1 day more as compared with the satisfied group. Patients who had a
123 postoperative deep infection had 3 times higher risk to be dissatisfied 4 years after TKA while
124 superficial infection or stiffness as well as preoperative radiographic mild osteoarthritis and
125 chronic widespread pain did not have any higher risk of dissatisfaction 4 years
126 postoperatively. All KOOS 5 subscales were significantly improved in both groups.

127 **Discussion:**

128 We found in this study that preoperative psychological distress had a significant correlation
129 with patient dissatisfaction 4 years after TKA. In fact, psychological distress, defined as
130 anxiety and/or depression, had the strongest statistical correlation with dissatisfaction 4 years
131 after TKA of all variables studied. The presence of radiographic grade 1 or 2 osteoarthritis
132 according to the Kellgren and Lawrence classification did not correlate with dissatisfaction.
133 Longer postoperative hospital stay correlated as well with dissatisfaction, which could be
134 explained as patients with anxiety/depression might feel safer in the hospital and are maybe
135 more afraid of going home early. In our hospital, length of stay is standardized to large extent,
136 and patients with preoperative higher risks or major postoperative complications are
137 frequently performed or moved to our larger university emergency hospital. Another reason

138 for longer hospital stay is postoperative complications, but it is unlikely that the anxiety
139 and/or depression group is more prone to general postoperative complications.

140 A limitation of this study is lack of clinical examination, performance tests, and radiographic
141 examination at the 4-year follow-up. However, in our previous study, we could not find
142 differences between dissatisfied and very satisfied patients regarding these subjects [18].

143 Another limitation in this study could be the use of 2 different types of prosthesis, and that 5
144 different surgeons were involved. On the other hand, this might be advantageous to generalize
145 the result. In addition, all surgeons in our study have nearly equal TKA outcome according to
146 our local registry.

147 An important reason for dissatisfaction after TKA is pain and poor knee function, and there
148 are several well-known explanations for these symptoms such as patella-related problems,
149 infection, stiffness, loosening, instability, fracture, tendon rupture, neurovascular injuries. In
150 Sweden, the 4 most common (more than 85%) reasons for primary TKA revisions of patients
151 with osteoarthritis during the period 2004-2013 were infection, loosening, patella-related
152 problems and instability [24].

153 Since the introduction of TKA there has been a continuous refinement in many of the aspects
154 of the surgery, including navigation, prosthesis design, preoperative templating, pain
155 management and infection prophylaxis. Despite these improvements the proportion of
156 dissatisfied patient is still relatively high (8%-28%) [13]. Khatib et al [13] reported in a
157 systematic review article of 19 studies that preoperative psychological status may affect the
158 outcome of TKA. However, the follow-up time was 1 year or less in 16 of the 19 studies
159 which is a relatively short follow-up time. Bonnin et al [15] reported in a review article of 10
160 studies that factors associated with a painful knee after TKA were preoperative anxiety /
161 depression, female gender and age less than 60 years. We presented our results as both crude

162 and adjusted to age, gender and body mass index. In a review article of 10 TKA cohort studies
163 Paulsen et al. found a correlation between preoperative distress and functional outcome in 6
164 studies while 4 did not [16]. Several studies have shown correlation between preoperative
165 psychological distress and poor surgical outcome after TKA [1, 25-32]. Most of these studies
166 had either fewer patients, or shorter follow-up times compared with our study. On the other
167 hand, there are studies showing no correlation between preoperative psychological distress
168 and surgical outcomes [33-36]. Valdes et al [37] found that patients with lower preoperative
169 radiographic scores and depression reported higher postoperative pain score. We found,
170 however, no correlation between the radiographic degree of OA and dissatisfaction. Dápuzzo
171 et al. [38] found that fibromyalgia patients after TKA had more complications and higher
172 incidence of remaining pain, but despite this, most patients felt satisfied with the procedure.
173 We did not find that patients with chronic widespread pain had higher risk of being
174 dissatisfied after TKA. Matsuda et al [39] found a negative correlation between satisfaction
175 and limited ROM. On the other hand Devers et al. [40] found no correlation between knee
176 flexion and satisfaction. Interestingly, in our study, all KOOS 5 subscales were significantly
177 improved at 4 years postoperatively, including the dissatisfied group, even though patients
178 reported being dissatisfied with the surgical outcome (Figure 1).

179

180 In summary, our results confirm the strong correlation between preoperative anxiety and
181 depression and later dissatisfaction of the surgical outcome. Psychological assessment and
182 treatment preoperatively might improve degree of satisfaction for TKA patients.

183 **Acknowledgments**

184 The authors thank Professor Jonas Ranstam, RC Syd, Clinical Sciences, Lund University, for
185 advice on statistics. They are grateful to all the medical staff of the Orthopaedics Department,

186 in Lund and Trelleborg Hospital especially Åsa Björkqvist and Ewa Persson, for excellent
187 cooperation. Ali collected and analyzed the data and prepared the manuscript. All the authors
188 designed the study and helped in writing the manuscript. Ali, Flivik, and Sundberg recruited
189 patients and performed the operations.

190

191 **Table 1:**

192 Patient characteristic

	Not Satisfied (n = 27)	Satisfied (n = 159)
Age, y ^a	72 (8)	73 (10)
Sex, F/M	16/11	104/55
BMI ^a	30 (5)	30 (5)
ASA 1/2/3	7/17/3	34/108/17
Charnley A/B/C	7/5/15	49/48/62
Anesthesia spinal/general	23 / 4	139 / 20
LOS ^a , d	5 (1)	4 (1)
ROM ^a preop	109 (9)	113 (14)
K&L grade 1-2	9/27	37/159
Chronic widespread pain preop.	7/27	21/159
Chronic widespread pain 4y	14/27	26/159
VAS ^a pain preop.	65 (12)	60 (16)
VAS ^a pain 4 y.	56 (18)	11 (10)
Anxiety/depression preop.	14/27	12/159
Anxiety/depression 4y	16/27	11/159
Deep infection	2/27	3/159
Superficial infection	2/27	4/159
Stiffness (flexion < 90 degree)	2/27	7/159

193 BMI, body mass index;LOS, length of stay in hospital; K&L, Kellgren and Lawrence; VAS,
194 Visual Analog Pain Scale; ROM, range of motion; ASA, American Society of
195 Anaesthesiologists Physical Status Classification.

196 ^a mean value, standard deviation in brackets

197

198 **Table 2.**

199 Crude estimates, Relative Risk (RR) for dissatisfaction.

Factor	RR	95% CI	P Value
Age	0.98	0.94-1.03	.5
Gender	0.80	0.39-1.63	.5
BMI	1.04	0.97-1.11	.3
Deep infection	2.90	0.93-9.04	.07
Superficial infection	2.40	0.73-7.91	.2
Stiffness (flexion < 90 °)	1.57	0.44-5.66	.5
K&L grade 1-2	0.66	0.32-1.36	.3
ASA	0.89	0.47-1.71	.7
LOS	1.56	1.34-1.82	<.001
Chronic widespread pain preop.	1.37	0.87-2.14	.2
VAS pain preop.	1.02	1.00-1.05	.07
ROM preop.	0.98	0.96-1.00	.08
Anxiety/depression preop.	6.63	3.52-12.49	<.001
KOOS Pain preop.	0.99	0.97-1.01	.6
KOOS Symptoms preop.	0.99	0.96-1.01	.2
KOOS ADL preop.	0.99	0.97-1.01	.2
KOOS Sport/Rec preop.	0.99	0.97-1.02	.4
KOOS QOL preop.	0.99	0.97-1.02	.6

200 CI, confidence interval; BMI, body mass index; K&L, Kellgren and Lawrence; LOS, length of
 201 stay in hospital; VAS, Visual Analog Pain Scale; ROM, range of motion; ASA, American
 202 Society of Anaesthesiologists Physical Status Classification; KOOS, Knee Injury and
 203 Osteoarthritis Outcome Score; ADL, activity of daily living; QOL, quality of life.

204

205

206 **Table 3:**
 207 Relative Risk (RR) for dissatisfaction adjusted for differences in age,
 208 gender and BMI

Factor	RR	95% CI	P Value
Deep infection	3.1	1.1-8.4	.03
Superficial infection	2.3	0.78-6.6	.1
Stiffness (flexion < 90 °)	1.6	0.43-6.2	.5
K&L grade 1-2	0.70	0.33-1.5	.4
ASA	0.85	0.42-1.7	.6
LOS	1.6	1.3–2.0	<.001
Chronic widespread pain preop.	1.4	0.90-2.3	.1
VAS pain preop.	1.02	1.00-1.05	.1
ROM preop.	0.98	0.96-1.00	.08
Anxiety/depression preop.	6.5	3.5-12	<.001
KOOS Pain preop.	1.00	0.98-1.02	.8
KOOS Symptoms preop.	0.99	0.96-1.01	.2
KOOS ADL preop.	0.99	0.97-1.01	.3
KOOS Sport/Rec preop.	0.99	0.96-1.02	.5
KOOS QOL preop.	1.00	0.97-1.02	.8

209 CI, confidence interval; BMI, body mass index; K&L, Kellgren and Lawrence; LOS, length of
 210 stay in hospital; VAS, Visual Analog Pain Scale; ROM, range of motion; ASA, American
 211 Society of Anaesthesiologists Physical Status Classification; KOOS, Knee Injury and
 212 Osteoarthritis Outcome Score; ADL, activity of daily living; QOL, quality of life.

213

214

215


216

217

218

219
220
221
222
223
224
225

Figure: 1
KOOS 5 subscales preop. and 4 years postoperative. KOOS, Knee Injury and Osteoarthritis Outcome Score; ADL activity of daily living; QOL, quality of life.



226
227

228 1. Scott CE, Howie CR, MacDonald D, Biant LC. Predicting dissatisfaction following total knee
229 replacement: a prospective study of 1217 patients. *The Journal of bone and joint surgery British*
230 *volume 92(9): 1253, 2010*

231 2. Bourne RB, Chesworth BM, Davis AM, Mahomed NN, Charron KD. Patient satisfaction after total
232 knee arthroplasty: who is satisfied and who is not? *Clinical orthopaedics and related research 468(1):*
233 *57, 2010*

234 3. Kim TK, Chang CB, Kang YG, Kim SJ, Seong SC. Causes and predictors of patient's dissatisfaction
235 after uncomplicated total knee arthroplasty. *The Journal of arthroplasty 24(2): 263, 2009*

236 4. Wylde V, Learmonth I, Potter A, Bettinson K, Lingard E. Patient-reported outcomes after fixed-
237 versus mobile-bearing total knee replacement: a multi-centre randomised controlled trial using the
238 Kinemax total knee replacement. *The Journal of bone and joint surgery British volume 90(9): 1172,*
239 *2008*

240 5. Fisher DA, Dierckman B, Watts MR, Davis K. Looks good but feels bad: factors that contribute to
241 poor results after total knee arthroplasty. *The Journal of arthroplasty 22(6 Suppl 2): 39, 2007*

242 6. Noble PC, Conditt MA, Cook KF, Mathis KB. The John Insall Award: Patient expectations affect
243 satisfaction with total knee arthroplasty. *Clinical orthopaedics and related research 452: 35, 2006*

244 7. Brander VA, Stulberg SD, Adams AD, Harden RN, Bruehl S, Stanos SP, Houle T. Predicting total knee
245 replacement pain: a prospective, observational study. *Clinical orthopaedics and related research*
246 *(416): 27, 2003*

247 8. Robertsson O, Dunbar MJ. Patient satisfaction compared with general health and disease-specific
248 questionnaires in knee arthroplasty patients. *The Journal of arthroplasty 16(4): 476, 2001*

249 9. Heck DA, Robinson RL, Partridge CM, Lubitz RM, Freund DA. Patient outcomes after knee
250 replacement. *Clinical orthopaedics and related research (356): 93, 1998*

251 10. Robertsson O, Dunbar M, Pehrsson T, Knutson K, Lidgren L. Patient satisfaction after knee
252 arthroplasty: a report on 27,372 knees operated on between 1981 and 1995 in Sweden. *Acta*
253 *orthopaedica Scandinavica 71(3): 262, 2000*

254 11. Hawker G, Wright J, Coyte P, Paul J, Dittus R, Croxford R, Katz B, Bombardier C, Heck D, Freund D.
255 Health-related quality of life after knee replacement. *The Journal of bone and joint surgery American*
256 *volume 80(2): 163, 1998*

257 12. Anderson JG, Wixson RL, Tsai D, Stulberg SD, Chang RW. Functional outcome and patient
258 satisfaction in total knee patients over the age of 75. *The Journal of arthroplasty 11(7): 831, 1996*

259 13. Khatib Y, Madan A, Naylor JM, Harris IA. Do Psychological Factors Predict Poor Outcome in
260 Patients Undergoing TKA? A Systematic Review. *Clinical orthopaedics and related research 473(8):*
261 *2630, 2015*

262 14. Djahani O, Rainer S, Pietsch M, Hofmann S. Systematic analysis of painful total knee prosthesis, a
263 diagnostic algorithm. *Archives of bone and joint surgery 1(2): 48, 2013*

264 15. Bonnin MP, Basiglini L, Archbold HA. What are the factors of residual pain after uncomplicated
265 TKA? *Knee surgery, sports traumatology, arthroscopy : official journal of the ESSKA 19(9): 1411, 2011*

266 16. Paulsen MG, Dowsey MM, Castle D, Choong PF. Preoperative psychological distress and
267 functional outcome after knee replacement. *ANZ J Surg 81(10): 681, 2011*

268 17. Rolfson O, Dahlberg LE, Nilsson JA, Malchau H, Garellick G. Variables determining outcome in
269 total hip replacement surgery. *The Journal of bone and joint surgery British volume 91(2): 157, 2009*

270 18. Ali A, Sundberg M, Robertsson O, Dahlberg LE, Thorstensson CA, Redlund-Johnell I, Kristiansson I,
271 Lindstrand A. Dissatisfied patients after total knee arthroplasty: a registry study involving 114
272 patients with 8-13 years of followup. *Acta orthopaedica 85(3): 229, 2014*

273 19. Ali A, Sundberg M, Hansson U, Malmvik J, Flivik G. Doubtful effect of continuous intraarticular
274 analgesia after total knee arthroplasty: a randomized double-blind study of 200 patients. *Acta*
275 *orthopaedica 86(3): 373, 2015*

276 20. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand 67(6):*
277 *361, 1983*

278 21. Roos EM, Roos HP, Ekdahl C, Lohmander LS. Knee injury and Osteoarthritis Outcome Score
279 (KOOS)--validation of a Swedish version. *Scandinavian journal of medicine & science in sports* 8(6):
280 439, 1998

281 22. Bergman S, Herrstrom P, Jacobsson LT, Petersson IF. Chronic widespread pain: a three year
282 followup of pain distribution and risk factors. *J Rheumatol* 29(4): 818, 2002

283 23. Barros AJ, Hiraakata VN. Alternatives for logistic regression in cross-sectional studies: an empirical
284 comparison of models that directly estimate the prevalence ratio. *BMC Med Res Methodol* 3: 21,
285 2003

286 24. SKAR. Swedish Knee Arthroplasty Register. Annual report 2015, Lunds University Hospital, Lund,
287 <http://www.knee.se>, 2015

288 25. Giesinger JM, Kuster MS, Behrend H, Giesinger K. Association of psychological status and patient-
289 reported physical outcome measures in joint arthroplasty: a lack of divergent validity. *Health and*
290 *quality of life outcomes* 11: 64, 2013

291 26. Hirschmann MT, Testa E, Amsler F, Friederich NF. The unhappy total knee arthroplasty (TKA)
292 patient: higher WOMAC and lower KSS in depressed patients prior and after TKA. *Knee surgery,*
293 *sports traumatology, arthroscopy : official journal of the ESSKA* 21(10): 2405, 2013

294 27. Duivenvoorden T, Vissers MM, Verhaar JA, Busschbach JJ, Gosens T, Bloem RM, Bierma-Zeinstra
295 SM, Reijman M. Anxiety and depressive symptoms before and after total hip and knee arthroplasty: a
296 prospective multicentre study. *Osteoarthritis and cartilage / OARS, Osteoarthritis Research Society*
297 *21(12): 1834, 2013*

298 28. Lingard EA, Katz JN, Wright EA, Sledge CB, Kinemax Outcomes G. Predicting the outcome of total
299 knee arthroplasty. *The Journal of bone and joint surgery American volume* 86-A(10): 2179, 2004

300 29. Brander V, Gondek S, Martin E, Stulberg SD. Pain and depression influence outcome 5 years after
301 knee replacement surgery. *Clinical orthopaedics and related research* 464: 21, 2007

302 30. Desmeules F, Dionne CE, Belzile EL, Bourbonnais R, Champagne F, Fremont P. Determinants of
303 pain, functional limitations and health-related quality of life six months after total knee arthroplasty:
304 results from a prospective cohort study. *BMC sports science, medicine and rehabilitation* 5: 2, 2013

305 31. Caracciolo B, Giaquinto S. Self-perceived distress and self-perceived functional recovery after
306 recent total hip and knee arthroplasty. *Arch Gerontol Geriatr* 41(2): 177, 2005

307 32. Bistolfi A, Bettoni E, Aprato A, Milani P, Berchiolla P, Graziano E, Massazza G, Lee GC. The
308 presence and influence of mild depressive symptoms on post-operative pain perception following
309 primary total knee arthroplasty. *Knee surgery, sports traumatology, arthroscopy : official journal of*
310 *the ESSKA, 2015*

311 33. Perez-Prieto D, Gil-Gonzalez S, Pelfort X, Leal-Blanquet J, Puig-Verdie L, Hinarejos P. Influence of
312 depression on total knee arthroplasty outcomes. *The Journal of arthroplasty* 29(1): 44, 2014

313 34. Blackburn J, Qureshi A, Amirfeyz R, Bannister G. Does preoperative anxiety and depression
314 predict satisfaction after total knee replacement? *The Knee* 19(5): 522, 2012

315 35. Riddle DL, Wade JB, Jiranek WA, Kong X. Preoperative pain catastrophizing predicts pain outcome
316 after knee arthroplasty. *Clinical orthopaedics and related research* 468(3): 798, 2010

317 36. Lingard EA, Riddle DL. Impact of psychological distress on pain and function following knee
318 arthroplasty. *The Journal of bone and joint surgery American volume* 89(6): 1161, 2007

319 37. Valdes AM, Doherty SA, Zhang W, Muir KR, Maciewicz RA, Doherty M. Inverse relationship
320 between preoperative radiographic severity and postoperative pain in patients with osteoarthritis
321 who have undergone total joint arthroplasty. *Semin Arthritis Rheum* 41(4): 568, 2012

322 38. D'Apuzzo MR, Cabanela ME, Trousdale RT, Sierra RJ. Primary total knee arthroplasty in patients
323 with fibromyalgia. *Orthopedics* 35(2): e175, 2012

324 39. Matsuda S, Kawahara S, Okazaki K, Tashiro Y, Iwamoto Y. Postoperative alignment and ROM
325 affect patient satisfaction after TKA. *Clinical orthopaedics and related research* 471(1): 127, 2013

326 40. Devers BN, Conditt MA, Jamieson ML, Driscoll MD, Noble PC, Parsley BS. Does greater knee
327 flexion increase patient function and satisfaction after total knee arthroplasty? *The Journal of*
328 *arthroplasty* 26(2): 178, 2011

