

# The Multifaceted Roles Neutrophils Play in the Tumor Microenvironment

Ronit Vogt Sionov · Zvi G. Fridlender · Zvi Granot

Received: 1 January 2014 / Accepted: 19 May 2014 / Published online: 4 June 2014  
© Springer Science+Business Media Dordrecht 2014

**Abstract** Neutrophils are myeloid cells that constitute 50–70 % of all white blood cells in the human circulation. Traditionally, neutrophils are viewed as the first line of defense against infections and as a major component of the inflammatory process. In addition, accumulating evidence suggest that neutrophils may also play a key role in multiple aspects of cancer biology. The possible involvement of neutrophils in cancer prevention and promotion was already suggested more than half a century ago, however, despite being the major component of the immune system, their contribution has often been overshadowed by other immune components such as lymphocytes and macrophages. Neutrophils seem to have conflicting functions in cancer and can be classified into anti-tumor (N1) and pro-tumor (N2) sub-populations. The aim of this review is to discuss the varying nature of neutrophil function in the cancer microenvironment with a specific emphasis on the mechanisms that regulate neutrophil mobilization, recruitment and activation.

**Keywords** Neutrophil function · Tumor Microenvironment · Pre-metastatic niche · Chemokines

## Introduction

Within the tumor microenvironment, there is a continuous crosstalk between tumor cells, stromal cells and cells of the

immune system [1–3]. Tumor cells secrete a wide range of cytokines and chemokines that not only attract and affect macrophages, neutrophils, dendritic cells, NK-, B- and T-cells, but also regulate other stromal cells, endothelial cells and the tumor cells themselves. Cells of the innate and adaptive immune system in turn mutually affect the activation status of other immune cells and attract more leukocytes into the tumor cell mass. Surprisingly, non-malignant immune cells can make up 90 % of the total tumor mass [2–5]. However, it is the integration of all signals encountered within the tumor microenvironment that affects the composition and activation of the various immune cells within the tumor cell mass and ultimately leads to either anti-tumor or pro-tumor activities.

Signals emanating from the tumor may also affect immune cell activities in the circulation and in distal pre-metastatic organs such as the liver and the lung to either restrain or encourage metastasis. Immune cell inhibition of tumor growth is part of the process termed immune surveillance, where circulating or tissue-associated immune cells eliminate malignancies. Conversely, immune cells secrete factors that can stimulate angiogenesis, remodel the extracellular matrix and promote cell growth thereby facilitating tumor growth and dissemination. The balance between these opposite actions is dictated by multiple factors secreted by the tumor cells, the tumor-associated stromal cells and the immune cells themselves. A future goal is to develop strategies that can modulate the tumor microenvironment in such a way that the anti-tumorigenic activities of the immune cells will prevail.

The goal of this review is to provide insight into the complexity of the tumor-stroma interaction, with a specific emphasis on the role played by neutrophils, which have recently emerged as a central player in the above-described interplay.

---

R. V. Sionov · Z. Granot (✉)  
Department of Developmental Biology and Cancer Research,  
Institute for Medical Research Israel Canada, Hebrew University  
Medical School, 91120 Jerusalem, Israel  
e-mail: zvikag@ekmd.huji.ac.il

Z. G. Fridlender  
Institute of Pulmonary Medicine, Hadassah-Hebrew University  
Medical Center, 96920 Jerusalem, Israel

## The Cancer-Neutrophil Crosstalk

In this review we will explore the influence of the microenvironment or niche on the phenotype and function of neutrophils in the context of tumor initiation, growth and metastatic progression. Significant controversy surrounds the function of neutrophils in the context of cancer. Neutrophils are often recruited to the tumor microenvironment where they have been shown to either promote or inhibit tumor growth [6–11]. There are several lines of evidence suggesting that neutrophils may exert direct cytotoxic activities towards the tumor, or indirectly lead to tumor regression through recruitment and induction of tumor-specific T cell responses [12–18]. For instance, tumor infiltrating neutrophils in tumor-bearing rats secrete chemotactic factors for T lymphocytes and recruit T lymphocytes to the tumor bed [19]. When neutrophils were selectively depleted at the time of *in vivo* priming with  $\gamma$ -irradiated tumor cells, the growth of subsequent transplanted syngeneic tumors was not inhibited, indicating that neutrophils are important for mounting specific anti-tumor immune responses [19]. However, even in the presence of activated neutrophils, cancer cells may generate an immunosuppressed microenvironment, preventing cytotoxic immune responses, while enhancing the tumor-promoting activities of the immune cells.

Below we discuss the interplay between cancer cells and neutrophils, and their ultimate role in tumor growth and metastatic progression. We propose that in the context of tumor growth and metastatic progression, neutrophils possess both pro- and anti-tumor properties and that their function is determined in a niche-dependent fashion.


## Niche-dependent Neutrophil Function

While chronic inflammation may promote tumorigenesis [5, 20], the tumor cells themselves also attract immune cells through the secretion of a wide range of chemokines such as IL-8 (CXCL8 in human/CXCL2 in mouse), CCL2 (MCP-1), CCL3 (MIP-1 $\alpha$ ), CCL5 (RANTES), CXCL6 (huGCP-2), and KC (CXCL1), and cytokines such as IL-1 $\beta$ , IL-6, TNF $\alpha$ , GM-CSF, and G-CSF, thereby inducing inflammation [21–32]. Cytokines from tumors may regulate tumor growth or modify the anti-tumor immune responses [33–36]. Cytokines and chemokines from tumor cells also modify their normal surrounding non-malignant stroma by modulating the function of epithelial cells, endothelial cells, fibroblasts and inflammatory cells to generate a supportive microenvironment [37–39]. For instance, TNF $\alpha$ , IL-6 and IL-17 were shown to promote tumor growth by modifying the function of stromal cells surrounding the tumor [40–42]. TNF $\alpha$  produced by tumor cells or inflammatory cells in the tumor microenvironment can promote tumor cell survival through the

induction of NF $\kappa$ B-dependent anti-apoptotic molecules [43]. TNF $\alpha$  was also shown to promote angiogenesis [44], and induce the expression of VEGF and HIF-1 $\alpha$  in tumor cells [45]. IL-6 promotes angiogenesis and the expression of VEGF [46] through JAK2/STAT3 signaling [42] and the tumor-promoting effects of IL-17 are in part mediated through up-regulation of IL-6 [42].

Traditionally, immune cells are viewed as protectors of the host where they take part in immune surveillance, eliminating both microbial infections and potentially cancerous cells. However, in the context of a tumor, the function of these cells is modified and they are “alternatively activated” to act against the host and promote tumor growth and metastasis. It is thought that tumors secrete factors that elicit a wound-repair response by tumor-associated macrophages (TAMs) and tumor-associated neutrophils (TANs) and that this response inadvertently stimulates tumor progression [47].

The phenomenon of alternative activation and the shift from anti-tumor to pro-tumor function of immune cells has been extensively described for macrophages. Macrophages have both pro- and anti-tumor actions depending on the activation signals, and have accordingly been classified into anti-tumor “M1” and pro-tumor “M2” macrophages (Fig. 1) [48, 6]. M1 macrophages show increased production of inflammatory cytokines (e.g., IL-1 $\beta$ , TNF $\alpha$ , IL-6, IL-12, IL-23) and Th1, Th17 and NK cell-attracting chemokines such as CXCL9/Mig and CXCL10/IP-10 [6]. Alternatively activated M2 macrophages have a strikingly different gene expression profile compared with M1 macrophages and express the immunosuppressive cytokine IL-10, tumor growth factors (e.g., EGF, FGF1, TGF $\beta$ 1), pro-angiogenic factors (e.g., VEGF), matrix remodeling factors (e.g., fibrin and matrix metalloproteinases), and chemokines such as CCL17/TARC, CCL22/MDC and CCL24/Eotaxin-2, that are involved in regulatory T (Treg) cell, Th2 cell, eosinophil and basophil recruitment [6, 49, 50]. M2 macrophages contribute to the formation of an immunosuppressed microenvironment. TAMs often show a M2 phenotype [51] and are characterized by the secretion of VEGF, HIF, TGF $\beta$ , IL-10, Arginase I and reactive oxygen species (ROS) [52], as well as various chemokines such as CCL2, CCL5 (RANTES), CXCL9, CXCL10, and CXCL16 [53]. TAMs contribute to tumor growth and progression through extracellular matrix remodeling, promotion of tumor cell invasion and metastasis, angiogenesis, lymphangiogenesis and immune suppression [6]. TAMs may also secrete cytokines IL-1 $\beta$ , IL-6 and IL-23 that trigger the proliferation of IL-17-producing CD8<sup>+</sup> Tc17 cells [54], a T cell subset that is often found to be present in various tumors [55]. IL-17 increases the synthesis of C-X-C chemokines from epithelial cells [56] leading to increased neutrophil infiltration into the tumor [57]. IL-17 also induces the production of G-CSF [58], thus creating a G-CSF-IL-17-IL-23 feedback axis that regulate neutrophil homeostasis [59].



**Fig. 1** Polarization of neutrophils and macrophages. Two major activation pathways have been described for macrophages and neutrophils. These immune cells can be activated through the classical pathway involving microbial components and T-cell derived factors such as IL-12 and IFN $\gamma$ , into Type 1 inflammatory cells that combat acute infection. Recent data suggest that IFN $\beta$  is also important for the polarization of neutrophils into N1 cells. In the presence of immunosuppressive components such as IL-10, TGF $\beta$ , PGE $_2$ , or excessive Th2 responses

characterized by IL-13 and IL-4 production, macrophages and neutrophils undergo an “alternative” activation pathway into Type 2 cells, which exhibit immunosuppressive activities. Also, excessive G-CSF and TNF $\alpha$  production may further promote Type 2 activation of neutrophils. Usually, the Type 1 activation prevails under acute inflammation, while Type 2 activation dominates under chronic inflammation, including the tumor microenvironment. Type 2 activation seems to be a negative feedback mechanism that counteracts excessive immune responses

Tumor-associated neutrophils (“TANs”), like macrophages, may acquire an anti-tumorigenic “N1” or pro-tumorigenic “N2” phenotype and are classified according to their activation state, cytokine repertoire and effects on tumor growth [7, 9–11, 60]. The N1 cells are characterized by cytotoxic activity towards tumor cells and an immunostimulatory profile (i.e., TNF $\alpha$ <sup>high</sup>, CCL3<sup>high</sup>, ICAM-1<sup>high</sup>, Arginase<sup>low</sup>), whereas N2 neutrophils are characterized by upregulation of the chemokines CCL2, 3, 4, 8, 12, and 17, and CXCL1, 2, 8 and 16 [10]. N1 cells produce more superoxide and hydrogen peroxide and express higher levels of Fas, TNF $\alpha$ , CCL3 and ICAM-1, but lower levels of Arginase, CCL2, CCL5, VEGF, CXCR4 and MMP-9 than N2 cells [10]. Proinflammatory N1 neutrophils have also been shown to promote CD8<sup>+</sup> T cells recruitment and activation by producing T cell-attracting chemokines (e.g., CCL3, CXCL9, and CXCL10) and proinflammatory cytokines (e.g., IL-12, TNF $\alpha$ , GM-CSF, and VEGF) [61]. There is a mutual interplay between neutrophils and CD4<sup>+</sup> T helper 17 cells (Th17) [62]. While IL-17 and CXCL8 secreted by Th17 cells induce the

recruitment of neutrophils, secretion of CCL2 and CCL20 by activated neutrophils attracts Th17 cells [61–63]. Th17 cells may further modulate neutrophil activity through secretion of TNF $\alpha$ , IFN $\gamma$  and GM-CSF [62].

The immunosuppressive transforming growth factor  $\beta$  (TGF $\beta$ ) was shown to promote the N2 neutrophil phenotype [13], while interferon  $\beta$  (IFN $\beta$ ) promotes the N1 phenotype [64] (Fig. 1). Inhibition of TGF $\beta$  signaling using the small molecule SM16 increased the mRNA levels for neutrophil chemoattractants (CXCL2 and CXCL5) in macrophages isolated from the tumor and increased the recruitment of CD11b<sup>+</sup> neutrophils into the tumor with concomitant reduced tumor growth [13]. Depletion of neutrophils prevented the SM16-induced growth inhibition, suggesting an anti-tumor activity mediated by neutrophils [13].

TANs differ from naïve neutrophils and G-MDSCs (the granulocytic sub-population of myeloid-derived suppressor cells) as they harbor low amounts of the various neutrophil granules and generate low amounts of ROS, while showing an enhanced chemokine secretion profile [10]. TANs

express higher levels of CXCL2, CXCL1 and CCL3 than naïve bone-marrow neutrophils [65], thus generating a positive feedback loop for recruiting more neutrophils as well as other immune cells to the tumor site. While TANs at the primary tumor site seem to acquire an N2 phenotype induced by high TGF $\beta$  levels, circulating neutrophils and neutrophils in metastasis-free organs tend to have an N1 anti-tumor-phenotype, due to paracrine exposure to activating tumor-secreted factors such as the chemokines IL-8, CCL2, CCL5 and CXCL5 [32, 66]. Indeed, two studies have clearly demonstrated an anti-tumor function for neutrophils at the pre-metastatic lung [32, 66]. These neutrophils have been coined tumor-entrained neutrophils (TENs), as these have been educated by the tumor to possess anti-tumor properties [32].

### Immunosuppression in the Tumor Microenvironment

Ultimately, tumors do develop in the presence of a functional immune system, suggesting that tumor cells acquire properties that help them evade immune surveillance [67]. Indeed, tumors have been shown to escape the anti-tumor immune responses by generating an immunosuppressed tumor microenvironment. Tumors often produce soluble immunosuppressive factors, such as TGF $\beta$  [13, 68, 69], VEGF [69, 70], IL-10 [69, 71], iNOS [72, 73], PGE<sub>2</sub> [74, 75], and gangliosides [76, 77], that act on neutrophils and other tumor infiltrating immune cells. Often, progressive immunosuppression is observed at advanced tumor stages, which is partially mediated by tumor-infiltrating immunosuppressive immune cells such as regulatory T (Treg) cells, Th17 cells, regulatory dendritic cells, TAMs, TANs and MDSCs. The mechanisms that mediate the immunosuppressive microenvironment deserve an in-depth exploration, which is beyond the scope of this review. Here we will only describe the effects of some of these immunosuppressive factors that are relevant to neutrophil function.

#### TGF $\beta$

Transforming growth factor  $\beta$  (TGF $\beta$ ), which exists in at least three isoforms  $\beta$ 1,  $\beta$ 2 and  $\beta$ 3, has been linked to the regulation of tumor initiation, progression and metastasis [78] and TGF $\beta$ 1 is frequently upregulated in human cancers [79, 80]. Tumor-secreted TGF $\beta$  is usually sequestered to the extracellular matrix as an inactive complex, and becomes activated through enzymes such as neutrophil-derived elastase and matrix metalloproteinase (MMP)-9, or expression of  $\alpha_v\beta_6$  integrin [78]. In addition, reactive oxygen free radicals produced by activated neutrophils can activate latent TGF $\beta$  [81]. Thus, activated neutrophils, through production of elastase, MMP-9 and ROS, may contribute to TGF $\beta$ -mediated immunosuppression, a mechanism that may drive a negative

feedback that prevents excessive immune responses. Tumor-associated MDSCs and TAMs are also significant sources of TGF $\beta$  production [82, 83]. As MDSC and TAMs phenotypes are affected by TGF $\beta$  [84], their initial appearance in the tumor would further contribute to the immunosuppressed microenvironment by enhancing the recruitment of these cells to the tumor bed.

The importance of TGF $\beta$  in immunosuppression was demonstrated by introducing the expression of a dominant-negative TGF $\beta$  type II receptor in mouse tumor-specific T cells. These *ex vivo* expanded TGF $\beta$ -insensitive CD8<sup>+</sup> T cells infiltrated the tumor and mediated apoptosis in tumor cells [68]. Fridlender et al. [13] showed that inhibition of TGF $\beta$  signaling using the small molecule inhibitor SM16 conferred anti-tumorigenic activity to neutrophils. Low concentrations of TGF $\beta$ 1 were further shown to inhibit neutrophil degranulation, as measured by lactoferrin release, in response to lipopolysaccharide (LPS) and formyl peptides [85]. TGF $\beta$  was also shown to prevent the production of ROS, reactive nitrogen intermediates and IL-1 $\beta$  by neutrophils [86]. Furthermore, TGF $\beta$  has even been shown to be a potent chemoattractant for neutrophils taking a central part in their recruitment to sites of inflammation [87–89]. However, another study showed that TGF $\beta$  reduces the expression of the adhesion molecule L-Selectin, resulting in impaired neutrophil recruitment to sites of inflammation [86]. Blockage of TGF $\beta$  signaling increased the numbers of neutrophils in tumors, which was associated with increased amount of chemokines and cytokines within the tumor, concomitant with increased ICAM-1 expression on endothelial cells [13, 90]. The increase in intra-tumor neutrophil number observed following anti-TGF $\beta$  signaling therapy can be explained by the ability of TGF $\beta$  to inhibit endothelial adhesiveness of neutrophils and neutrophil transmigration *in vivo* [91]. Abrogation of TGF $\beta$  signaling in mammary carcinomas also led to increased infiltration of Gr-1<sup>+</sup>CD11b<sup>+</sup> MDSCs into the invasive front of tumor tissues facilitating tumor cell invasion and metastasis through a process involving metalloproteinase activity [82, 92]. Interestingly, MDSCs from TGF $\beta$  signaling deficient tumor-bearing hosts produced higher levels of VEGF, MMP-2, MMP-13 and MMP-14 than those isolated from normal mice [82].

#### PGE<sub>2</sub>

Prostaglandins (PGs) are small-molecular derivatives of arachidonic acid, produced by cyclooxygenases (COXs; constitutively active cyclooxygenase COX-1 and inducible COX-2) and PG synthases [93]. COX-2 expression is induced by cytokines and growth factors at sites of inflammation and is usually not detected in normal tissues [94]. Increased COX-2 expression and PGE<sub>2</sub> production has been reported in



human colon carcinoma [95], breast cancer [96, 97], renal cell carcinoma [98], and lung carcinoma [99].

Prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) directly suppresses various immune cells such as macrophages, neutrophils, Th1, CTL and NK cells, while it promotes Th2, Th17 and regulatory T cell responses as well as the development of tumor-associated suppressive macrophages [93, 100–103]. The Th17-promoting activity of PGE<sub>2</sub> is related to its ability to suppress the production of the Th17-inhibitory IL-12, while enhancing the Th17-supporting IL-23 secretion by dendritic cells [104]. As mentioned above, IL-17 promotes neutrophil recruitment [59], such that increased Th17 activity caused by PGE<sub>2</sub> has indirect effect on neutrophil function. Indeed, PGE<sub>2</sub> has been shown to stimulate IL-23/IL-17-induced neutrophil migration in inflammation by inhibiting IL-12 and IFN $\gamma$  production, which may be one mechanism for its pro-inflammatory effect [105]. PGE<sub>2</sub> also promotes MDSC recruitment to tumor through the local induction of CXCL12/SDF-1 [100]. In addition, PGE<sub>2</sub> promotes the recruitment of CD4<sup>+</sup>CD25<sup>+</sup> Tregs to the tumor and has direct positive effects on tumor progression [106]. PGE<sub>2</sub> may also promote tissue influx of neutrophils [107], macrophages [108] and mast cells [109]. One mechanism is through PGE<sub>2</sub>-stimulated production of IL-8 by epithelial cells [107] and pulmonary microvascular endothelial cells [110], as well as MCP-1 by mast cells [108]. Furthermore, activated neutrophils have been shown to express COX-2 and secrete PGE<sub>2</sub> [111, 112], which may represent another mechanism for immune response restriction. However, PGE<sub>2</sub> may also contribute to dysregulated inflammatory responses by increasing vascular permeability that facilitates influx of pro-inflammatory polypeptides. While PGE<sub>2</sub> prolongs neutrophil half-life through upregulation of intracellular cAMP levels and inhibition of apoptosis [113], it also attenuates PMA-, fMLP- or GM-CSF-stimulated ROS and LTB<sub>4</sub> production in neutrophils [114–117]. The Th2 cytokine IL-13 was found to increase neutrophil PGE<sub>2</sub> production concomitant with increased expression of complement receptor type 1 (CR1) and type 3 (CR3), and increased neutrophil phagocytosis [118].

In line with the immunosuppressive, tumor promoting function of PGE<sub>2</sub>, inhibitors of prostaglandin synthesis, such as indomethacin and aspirin, have been shown to inhibit tumor growth [101] and restore anti-tumor activity by altering the balance between IL-10 and IL-12 [119]. Recently, this issue has reached a renaissance where specific COX2 inhibitors have been proposed to be potential drugs for tumor prevention [120]. Non-steroidal anti-inflammatory drugs (NSAIDs), such as diclofenac, increased neutrophil superoxide production, but also promoted L-Selectin down-regulation [121].

Interestingly, a mutual interplay between TGF $\beta$  and PGE<sub>2</sub> seems to exist, where TGF $\beta$  stimulation of CD4<sup>+</sup> T cells induces production of PGE<sub>2</sub>, and PGE<sub>2</sub> contributes to TGF $\beta$ -induced suppression of T cells [122].

#### Other Immunosuppressive Molecules

The expression of indoleamine 2,3-dioxygenase (IDO), a tryptophan catabolizing enzyme, in tumors may lead to dysfunctional T cell response through depletion of tryptophan from the tumor microenvironment [123, 124]. IDO is also produced by MDSCs, regulatory dendritic cells and TAMs [6, 52, 125, 126]. Silencing of IDO within the tumor using *Salmonella typhimurium* as a tumor-homing vector to deliver a short-hairpin RNA targeting IDO, allowed tumor infiltration of activated ROS-producing neutrophils and consequent tumor cell death [127].

Another mechanism leading to suppression of T-cell mediated immune responses is excessive production of adenosine by the cell surface enzyme CD73 (ecto-5'-nucleotidase) [128, 129]. CD73 is usually expressed on endothelial and epithelial cells [130], subsets of leukocytes [131] and Foxp3<sup>+</sup> Tregs [128, 129], but also on several cancer types [132, 133]. CD73 acts in concert with CD39 (ecto-apyrase) to produce adenosine in a coordinated two-step enzymatic conversion. Both CD39 and CD73 seem to attenuate neutrophil trafficking into the lungs during LPS-induced injury [134] suggesting that CD73 expression on tumor cells is likely to limit neutrophil infiltration. CD73-deficient mice have increased anti-tumor immunity and are resistant to experimental metastases [135]. Similarly, anti-CD73 antibody therapy was found to inhibit breast tumor growth and metastasis [136].

#### Regulation of Neutrophil Mobilization, Recruitment and Activation in Cancer

##### Neutrophil Mobilization in Cancer

Human cancers often induce elevated numbers of circulating neutrophils [6, 7, 9–11, 137–159]. The consequences of cancer-induced neutrophilia in human patients will be further discussed in the section discussing “Prognostic Values of Neutrophils and Other Myeloid Subtypes in Cancer Patients”. In tumor bearing mice, a phenomenon similar to what occurs upon inflammation, is observed, namely, the number of circulating neutrophils increase dramatically and are associated with the progression of the disease [32, 160–162]. For example, using the 4T1 mammary tumor model in Balb/c mice, we showed that within 1 week after orthotopic inoculation of the tumor, circulating neutrophil numbers increased from ~17 % to over 30 % [32]. This increase continues with the

progression of the disease, reaching a state of acute neutrophilia with neutrophils making ~90 % of all circulating white blood cells [32]. Similar increase in circulating neutrophil numbers was seen in other mouse models of cancer including Lewis lung carcinoma [163] as well as in spontaneous mouse models such as the MMTV-PyMT and MMTV-Wnt1 transgenic mice [32], where tumor initiation is driven by a transgene, rather than an engrafted tumor. In a rat model of 13762NF mammary adenocarcinoma cells, the number of circulating neutrophils did not increase in poorly metastatic cells, whereas the number rose 50-fold in rats bearing a highly metastatic clone [164]. An intermediate rise in neutrophil number (12–14-fold) was observed in moderately metastatic tumors [164]. The increase in neutrophil number correlated with the ability of the tumor cells to secrete CSF [165]. These tumor cells did not induce a cytotoxic neutrophil response, while *i.v.* co-injections of tumor-elicited neutrophils caused a dose-dependent increase in extrapulmonary metastases that was associated with increased production of heparanase and type IV collagenolytic enzymes by the neutrophils [164]. In contrast to tumor-elicited neutrophils, normal or protease peptone-elicited neutrophils did not alter the invasive potential [166].

*How are the Neutrophils Mobilized?* Unlike the situation in infection and inflammation where neutrophil mobilizing factors are secreted by endothelial cells and other stromal cells, in the context of cancer, neutrophil mobilizing factors are often secreted by the tumor cells themselves [22]. The most common neutrophil chemoattractants produced by tumors include IL-8 (CXCL8/CXCL2), MIP-1 $\alpha$  (CCL3), huGCP-2 (CXCL6) and KC (CXCL1) [167–171]. G-CSF is ectopically expressed in several human tumors such as leukemia [172], bladder [173], pancreatic [174], cervical [175], ovarian [176], head and neck [177], colorectal [178] and breast carcinoma [179]. Similarly, some human cancers show elevated GM-CSF expression levels [31, 180, 181]. It is therefore not surprising that elevated numbers of circulating neutrophils are seen in a wide variety of human malignancies.

GM-CSF and G-CSF are broadly used therapeutically in cancer patients for their positive effects on bone marrow mobilization and immune functions. They are especially important for overcoming neutropenia caused by various anti-neoplastic treatments. However, accumulating studies show that these factors also promote the expansion of myeloid suppressive components, with undesirable consequences on tumor antigen-specific immune responses [182]. For instance, GM-CSF-based anti-tumor vaccine to human metastatic melanoma patients induced a subset of immunosuppressive MDSCs that involved TGF $\beta$  secretion [183]. GM-CSF may increase immune responses when administered at low doses, while causing an opposite effect at high doses [182]. While physiological concentrations of GM-CSF are required for

normal myelopoiesis, chronic administration of GM-CSF resulted in the generation of immune suppressive Gr-1<sup>+</sup>CD11b<sup>+</sup> cells in mice [184]. Experimental tumors overexpressing GM-CSF induced a systemic increase of immature myeloid cells, which was associated with suppression of T cell immune responses [184–186]. However, irradiated cancer cells engineered to secrete GM-CSF elicited potent anti-tumor immune responses in various animal tumor models [187].

Similarly, tumor-secreted G-CSF that contributes to neutrophil mobilization, activation and stimulation of oxidative metabolism [188–191], is also involved in the polarization towards immunosuppressive MDSCs [192, 193]. The presence of G-MDSCs was shown to be important for promoting tumor growth [193]. Besides stimulating neutrophils, G-CSF may stimulate non-hematopoietic malignant tumor cell growth in an autocrine fashion [173, 194]. G-CSF receptor (G-CSFR) expression has been observed in bladder cancer cells [195], ovarian cancer [196], colorectal cancer [197] and Ewing sarcoma [198]. G-CSF administration increased tumor growth of Erwin sarcoma [198], and must therefore be carefully considered before use for stimulating neutrophil recruitment following chemotherapy.

#### Neutrophil Recruitment to the Tumor Microenvironment

Neutrophils make up a significant proportion of the non-malignant stroma that contributes to the tumor microenvironment [199]. The increase in circulating neutrophil numbers in the context of cancer (see “[Neutrophil Mobilization in Cancer](#)” section) may passively lead to an increase in the absolute number of neutrophils marginating at the tumor microenvironment. However, neutrophils are also actively recruited into the tumor microenvironment in both cancer patients and mouse models of cancer. In humans, intratumoral infiltration of neutrophils has been detected in gastric carcinoma [200–202], bronchioloalveolar carcinoma [170, 203], non-small cell lung carcinoma [204], pancreatic neoplasia [205], pancreatic ductal adenocarcinoma [206], bladder cancer [147], glioma [207], cervical carcinoma [139] and breast carcinoma [4, 208] (Table 1). The clinical consequences of intratumoral and circulating neutrophils in cancer patients will be discussed in “[Prognostic Values of Neutrophils and Other Myeloid Subtypes in Cancer Patients](#)” section.

Neutrophils are actively recruited to the tumor microenvironment along a chemotactic gradient of tumor-secreted factors. Production of CXCL8/IL-8 and related chemokines occurs downstream to oncogene activation [209–211]. Bellocq et al. [170] observed a direct correlation between the number of neutrophils and IL-8 levels in bronchoalveolar lavage fluids from bronchioloalveolar carcinoma patients, suggesting a role for this cytokine in neutrophil recruitment. Neutrophils were mainly located in the alveolar lumen, while seldom in the alveolar wall [170]. CXCR1 and CXCR2 are the major

**Table 1** Tumor-associated neutrophils (TANs) in human cancer

Cancer type	Major findings	Reference
Bladder cancer	<ul style="list-style-type: none"> <li>• Increase in circulating CD11b<sup>+</sup>CD15<sup>high</sup>CD33<sup>low</sup> granulocytes, while no increase in CD11b<sup>+</sup>CD15<sup>low</sup>CD33<sup>high</sup> monocytes. Both cell types are activated.</li> <li>• Presence of circulating monocyte-macrophage CD11b<sup>+</sup>HLA-DR<sup>+</sup>, and granulocytic CD11b<sup>+</sup>CD15<sup>+</sup>HLA-DR<sup>-</sup> myeloid cells.</li> <li>• Myeloid cells secrete CCL2, CCL3, CCL4, G-CSF, IL-8 and IL-6.</li> <li>• Granulocytes inhibit in vitro T cell proliferation through induction of CD4<sup>+</sup>Foxp3<sup>+</sup> T regulatory cells.</li> </ul>	[147]
Metastatic adenocarcinomas of the pancreas, colon, and breast	<ul style="list-style-type: none"> <li>• Unusually large number of circulating granulocytes co-purified with low density PBMCs on a density gradient.</li> <li>• Increased oxidative stress and production of H<sub>2</sub>O<sub>2</sub> in circulating granulocytes.</li> <li>• H<sub>2</sub>O<sub>2</sub>-dependent reduction in T-cell receptor ζ chain expression and IFNγ production in T cells.</li> </ul>	[278]
Bronchioloalveolar carcinoma	<ul style="list-style-type: none"> <li>• Neutrophils are present in the alveolar lumen. The number of neutrophils correlate with IL-8 levels.</li> </ul>	[170]
Stage III and IV lung and stomach cancer	<ul style="list-style-type: none"> <li>• Patients with lung and stomach cancer showed increased blood neutrophil count, but decreased level of leukocyte cationic proteins.</li> </ul>	[386]
Recurrent localized cervical cancer	<ul style="list-style-type: none"> <li>• High density of CD66b<sup>+</sup> neutrophils and CD163<sup>+</sup> macrophages in peritumoral compartment.</li> </ul>	[139]
Hepatocellular carcinoma (HCC)	<ul style="list-style-type: none"> <li>• Intratumoral CD66b<sup>+</sup> neutrophils correlate with CD8<sup>+</sup> T cells, TGFβ expression, BCLC stage and early recurrence.</li> <li>• Increased intratumoral neutrophil numbers were associated with decreased overall survival, while peritumoral neutrophils were not associated with the outcome.</li> </ul>	[387]
Hepatocellular carcinoma (HCC)	<ul style="list-style-type: none"> <li>• Neutrophils from HCC patients produced CCL2 and CCL3. High CCL2 production was associated with reduced overall survival.</li> </ul>	[388]
Hepatocellular carcinoma (HCC)	<ul style="list-style-type: none"> <li>• Pro-inflammatory IL-17-producing cells recruit neutrophils into the peritumoral stroma of hepatocellular carcinoma (HCC) by epithelium-derived CXC chemokines.</li> <li>• Neutrophils promote angiogenesis at the adjacent tumor-invading edge via MMP-9 signaling.</li> <li>• Selective depletion of neutrophils inhibits tumor progression and growth.</li> </ul>	[57]
Resectable non-small cell lung cancer	<ul style="list-style-type: none"> <li>• Intratumoral CD66b<sup>+</sup> neutrophils were elevated in 50 % of the patients.</li> <li>• An increase in CD66b<sup>+</sup> cells was associated with high incidence of relapse and worse overall survival.</li> </ul>	[204]
Colorectal cancer	<ul style="list-style-type: none"> <li>• Increase in MPO<sup>+</sup> and CD15<sup>+</sup> cell infiltrate in the mucosa.</li> <li>• While MPO<sup>+</sup> cells were largely CD15<sup>+</sup>CD66b<sup>+</sup>, a high percentage of CD15<sup>+</sup>CD66<sup>-</sup> cells were MPO<sup>-</sup>.</li> <li>• Only high density of MPO<sup>+</sup> cell infiltration was associated with improved survival.</li> </ul>	[362]
Colorectal carcinoma	<ul style="list-style-type: none"> <li>• High intratumoral CD66<sup>+</sup> cells were associated with a poorer prognosis.</li> </ul>	[365]
Melanoma stage I/II	<ul style="list-style-type: none"> <li>• Presence of CD66b<sup>+</sup> tumor infiltrating neutrophils was associated with poor prognosis.</li> </ul>	[389]
Renal cell carcinoma	<ul style="list-style-type: none"> <li>• The presence of intratumoral neutrophils was associated with increased tumor size and shorter recurrence-free survival.</li> </ul>	[390]
Cervical cancer	<ul style="list-style-type: none"> <li>• The highest densities of CD66b<sup>+</sup> neutrophils and CD163<sup>+</sup> macrophages were observed in the peritumoral environment.</li> <li>• High peritumoral and stromal neutrophils were associated with shorter recurrence-free survival.</li> </ul>	[139]
Various cancer samples	<ul style="list-style-type: none"> <li>• CD15<sup>+</sup> neutrophils were found in substantial amounts in untreated malignant hepatocellular, cervical, colorectal and gastric carcinoma.</li> <li>• The CD15<sup>+</sup> cells were more abundant in the peritumoral stroma than in the cancer nest.</li> <li>• The peritumoral stroma CD15<sup>+</sup> cell density was associated with intrahepatic metastasis in liver cancer and lymph node metastasis in gastric cancer.</li> </ul>	[391]

Evidence for the presence of neutrophils within human cancer specimens and its prognostic value

neutrophil receptors that mediate neutrophil chemotaxis to the tumor microenvironment in response to IL-8. The same receptors also mediate neutrophil response to other ligands such as CXCL1/2/3, ENA-78 (epithelial-cell-derived neutrophil attractant-78; CXCL5), GCP-2 (granulocyte chemotactic protein-2; CXCL6) and NAP-2 (neutrophil-activating peptide 2; CXCL7) [212–215]. The neutrophil-sensitive human HT-29 colorectal adenocarcinoma and FaDu pharyngeal

squamous-cell carcinoma cells secrete IL-8 and GROα, and induced the adhesion of neutrophils to ICAM-1 on microvascular-endothelial cell monolayers resulting in transmigration through the endothelial cell monolayer [216]. Recruited neutrophils also release cytokines and chemokines, which enhance their own recruitment and activation in addition to inducing the migration of other immune cells [217]. Neutrophil-secreted MMP-9 may process the chemokine

CXCL5 to further promote neutrophil recruitment [214]. Similarly, neutrophil-derived Cathepsin G may increase the chemotactic activity of CXCL8, CXCL5 and CCL15 through N-terminal truncation [218].

Ly6G<sup>+</sup> neutrophils are the dominant source of CXCR2 in the blood, and CXCR2 deficiency attenuates neutrophil recruitment to the tumor bed [219]. Depletion of Ly6G<sup>+</sup> cells purged CXCR2-dependent tumor-associated leukocytes, suppressed established skin tumor growth and colitis-associated tumorigenesis [219]. CXCR2 is thus a potent pro-tumorigenic chemokine receptor that directs recruitment of tumor-promoting leukocytes into tissues during tumor-inducing and tumor-driven inflammation [219]. This notion correlates well with the important role CXCR2 plays in recruiting neutrophils to inflammatory sites.

The granulocyte chemotactic protein (GCP)-2/CXCL6 was shown to be important for the recruitment of neutrophils to melanoma tumors, and is associated with angiogenesis and tumor growth [171, 220]. CXCL6 specific antibodies reduced the recruitment of neutrophils to the tumor, with concomitant reduction in tumor growth [220]. There are several lines of evidence to suggest that CD8<sup>+</sup> T cells may promote the recruitment of neutrophils to the tumor, supposedly through secretion of IFN $\gamma$  [14] and vice versa, neutrophils contribute to T cell recruitment and activation [14]. Elimination of either neutrophils or CD8<sup>+</sup> T cells, or administration of IFN $\gamma$  neutralizing antibodies prevented tumor regression, suggesting a tight co-operation between neutrophils and CD8<sup>+</sup> cells in eliminating the tumor [14]. In contrast, regulatory T cells may limit neutrophil recruitment to the tumor site, which might be related to decreased expression of neutrophil chemoattractants such as CCL3, CXCL1 and CXCL2 [221].

CXCL12 and VEGF are also suggested to act as chemoattractants for myeloid cells [222, 223] and may act in concert for recruiting neutrophil to the tumor microenvironment. Under normal conditions, a tight cooperation between myeloid cells and endothelial progenitor cells is required for proper neovascularization [223]. VEGF induces bone marrow-derived myeloid cell mobilization to the circulation and through VEGF-mediated upregulation of SDF1/CXCL12 in activated perivascular myofibroblasts, the myeloid cells are kept in close proximity to angiogenic vessels [223]. VEGF is upregulated in a wide variety of tumors [69, 224–227], as is CXCL12 [228]. CXCL12, that is also upregulated by hypoxia [229], augments CXCR4 expression on vascular endothelial cells [230] and attracts CXCR4<sup>+</sup> cells, including neutrophils, to the tumor [223].

While tumor cells are capable of secreting neutrophil chemoattractants, other cells in the tumor microenvironment may also contribute to neutrophil recruitment. For example, tumor-infiltrating T helper type 17 (Th17) cells and IL-17 induce the expression of G-CSF, leading to immature

myeloid-cell mobilization and recruitment into the tumor microenvironment [231]. IL-17 may also induce the expression of C-X-C chemokines, notably CXCL8/IL-8, in epithelial cells that in turn recruit neutrophils to the tumor [57]. The presence of Th17 cells within the tumor microenvironment could antagonize and counter the tumor-suppressive IFN $\gamma$ -producing CD4<sup>+</sup> Th1 cells, and are thereby likely involved in the promotion of tumor growth [83]. However, in a lung melanoma mouse cancer model, adoptive transfer of Th17 cells promoted tumor-specific CD8<sup>+</sup> T cell activation [232], suggesting for a dual role for Th17 cells in tumor biology. The Th17 cells promoted dendritic cell recruitment into the tumor tissues and stimulated CCL20 chemokine production by the tumor [232].

Recently, another tumor-derived cytokine, IL-35, was shown to promote tumor growth by enhancing myeloid cell accumulation and angiogenesis [233]. IL-35 does not directly inhibit tumor-associated CD8<sup>+</sup> T cell activation, differentiation, or effector functions, but IL-35-treated cancer cells showed increased expression of gp130 and reduced sensitivity to CTL destruction [233]. Altogether, these data suggest that tumor cells induce a chronic inflammatory response.

#### Neutrophil Activation in Cancer

Neutrophils are traditionally perceived as the first line of defense against microbial infections and as mediators of inflammation, which possess favorable properties that protect the host. In the context of cancer, neutrophils were shown to exert both pro- and anti-tumor activities suggesting for a dual mode of activation which is the basis for the distinction between N1 and N2 neutrophils, as described in “Niche-dependent Neutrophil Function” section. Interestingly, several tumor-derived cytokines can activate neutrophil cytotoxic activities, including CCL2, CCL5, CCL3, CXCL1, SDF1/CXCL12 and CXCL16 [32, 66]. However, these chemokines are usually associated with pro- rather than anti-tumor activities. For example, CCL2 (MCP-1), which is overexpressed in a wide range of cancers [234], is associated with poor prognosis in breast, colorectal, cervical and thyroid cancers [235–239]. Tumor-derived CCL2 plays a pro-tumor role by recruiting inflammatory monocytes to pulmonary metastases [240] and MDSCs to the tumor microenvironment [241], as well as promoting angiogenesis [242], tumor cell proliferation [243] and migration [244]. On the other hand, CCL2 prevents apoptosis of neutrophils [245] and activates neutrophils in the pre-metastatic lung towards an anti-tumor phenotype where they produce H<sub>2</sub>O<sub>2</sub> to kill disseminated tumor cells [32, 66], suggesting a dual role of this chemokine. Also, CXCL1 was shown to promote tumor cell proliferation [246], tumor angiogenesis [247], invasion and migration [248], in addition to its ability to recruit and activate neutrophils [249]. CXCL1 was shown to be involved in a paracrine



network mediating both metastatic progression and chemoresistance [250]. IL-8 is another chemokine expressed in variety of tumors and is associated with neutrophil recruitment and activation [251]. For example, IL-8 secreted by human fibrosarcoma and prostate carcinoma cells promoted the infiltration of MMP-9<sup>+</sup> neutrophils [252] and melanoma secreted IL-8 was shown to increase CD11b/CD18 expression on neutrophils, an indication for their activation [253, 254].

A dual role has also been observed for G-CSF. On the one hand, G-CSF is an essential cytokine for mobilization of neutrophils, and under certain conditions, activates them. On the other hand, G-CSF may polarize granulocytes to promote tumor growth and metastasis [26]. These seemingly conflicting roles played by both neutrophils and the tumor-derived chemokines, put forth the question of what determines the overall contribution of neutrophils to cancer? It is likely that the pro- and anti-tumor activities of neutrophils are determined by the overall cytokine and chemokine milieu provided by the tumor and the tumor infiltrating cells rather than the expression level of a specific chemokine. This notion is supported by the fact that TGF $\beta$  was shown to be a potent repressor of neutrophil cytotoxicity both in vivo and in vitro [13, 32]. This is best exemplified by the differences in the function of neutrophils at the primary tumor site versus their activity at the pre-metastatic niche. High TGF $\beta$  activity in the tumor microenvironment generates immune suppressive conditions, attenuating the potential cytotoxicity of activated neutrophils recruited to the tumor bed [13]. At the same time, the full extent of activated neutrophil anti-tumor cytotoxicity may be manifested in distant organs, i.e. the pre-metastatic niche, where TGF $\beta$  levels are low [32, 66].

### **Tumor-Associated Neutrophils (TANs): Pro- and Anti-Tumor Mechanisms**

#### Neutrophils in Tumor Initiation - Inflammation-Associated Tumorigenesis


Chronic inflammation has been associated with increased susceptibility for cancer [20]. This is well demonstrated in chronic Hepatitis B virus infection [255], where the persisting insult and chronic inflammation ultimately leads to hepatocellular carcinoma. Another well-studied example is the correlation between inflammatory bowel (IBD) disease and colorectal cancer (CRC) with up to 30 % of IBD patients developing CRC [256]. The inflammatory process involves a wide range of immune cells of which neutrophils make a significant fraction. It is therefore reasonable to assume that neutrophils also play a part in inflammation-driven tumorigenesis. Indeed,

neutrophils were shown to directly promote tumorigenesis by causing genomic instability [257]. Neutrophil-induced genotoxicity has been related to induction of oxidative DNA damage through release of ROS and myeloperoxidase-related metabolites [258–260]. In addition, neutrophils were shown to promote experimental chronic colitis-associated carcinogenesis in mice [261]. Neutrophil recruitment to the inflamed submucosa was mediated by the chemokine CXCL2 through the neutrophil receptor CXCR2, resulting in increased neutrophil secretion of MMP-9 and neutrophil elastase, and consequent excessive angiogenesis and cell proliferation [261]. Elastase degrades barrier-forming proteins on epithelial cells [262] and neutrophil-derived elastase was found to degrade the adhesion molecule E-Cadherin on pancreatic ductal adenocarcinoma, leading to tumor cell dyshesion and increased migratory capacity [206]. Neutrophil infiltration of pancreatic ductal adenocarcinoma cells was also associated with epithelial-to-mesenchymal transition (EMT). In vitro cultivation of tumor cells with neutrophils led to enhanced expression of the transcription factor TWIST, translocation of  $\beta$ -Catenin to the nucleus, appearance of ZEB1 in the nucleus and downregulation of keratins, a sign of (EMT) [263].

Under certain circumstances, administration of neutrophil-neutralizing antibodies reduced the number and size of tumors [261, 264, 265]. The tumorigenic effect of MMP-9, predominantly provided by neutrophils and mast cells, was also reported in a mouse model of skin cancer where MMP-9 knockout mice showed decreased incidence of invasive tumors [266]. While there is a large body of evidence to support the pro-tumorigenic function of neutrophils, several studies show that neutrophils also possess anti-tumorigenic functions. For example, neutrophils were shown to participate in immune surveillance and eliminate potentially malignant cells and neutrophil MMP-8 was shown to provide protection against carcinogen-induced skin tumors [267].

#### Neutrophils at the Primary Tumor Site

Neutrophils are frequently found in solid tumors [268] (see “Neutrophil Recruitment to The Tumor Microenvironment” section), and together with macrophages and NK cells make up the vast majority of tumor infiltrating cells. The fact that neutrophils are frequent residents of solid tumors is not surprising as many of these tumors express and secrete high levels of neutrophil mobilizing factors and neutrophil chemoattractants. For example, IL-8/CXCL8, a potent neutrophil chemotactic factor and activator, was found to be expressed in a wide variety of human cancers [253, 269–272]. In most cases, IL-8 is secreted by the tumor cells themselves [210], while in others, other stromal cells such as fibroblasts [273] and macrophages [274] secrete IL-8. Some human cancers secrete high levels of G-CSF and GM-CSF



**Fig. 2** The crosstalk between cancer cells, neutrophils and other immune cells in the tumor microenvironment. The tumor microenvironment is characterized by a state of chronic inflammation, where tumor cells secrete a range of cytokines and chemokines that recruit and activate neutrophils and other immune cells. Accumulating evidence suggest that neutrophils are essential for mounting an adaptive immune response, as this fails to occur upon depletion of neutrophils. Neutrophils exert both pro-tumor and anti-tumor activities. Through production of ROS, nitric oxide and TRAIL, neutrophils can directly kill the tumor cells. Neutrophils also indirectly prevent tumor growth through eliciting specific anti-

tumor CD8<sup>+</sup> cytotoxic responses. By virtue of their multifunctional tasks, neutrophils may also stimulate tumor growth through secretion of ECM remodeling enzymes and pro-angiogenic factors. As in chronic inflammation, a negative feedback mechanism involved in restricting immune responses, such as the generation of MDSCs, also occurs within the tumor, generating an immunosuppressed environment that antagonizes the anti-tumorigenic activities of neutrophils. The secretion of immunosuppressive factors by the tumor cells themselves further fortifies this immunosuppressed milieu. (*TANs* Tumor Associated Neutrophils, *TAMs* Tumor Associated Macrophages)

[275, 276], acting as both potent mediators of neutrophil mobilization, recruitment and activation (Fig. 2).

Recruited neutrophils were shown to acquire a pro-tumor phenotype similar to that of tumor-associated monocytes, which by analogy to the M2 monocyte phenotype, was referred to as an N2 neutrophil phenotype [13]. TANs were shown to support tumor growth by producing angiogenic factors and matrix-degrading enzymes [8, 57, 64, 252, 264, 277]. Neutrophils were also shown to promote the acquisition of a metastatic phenotype [265] and to suppress anti-tumor immune responses [278]. Neutrophil-derived CXCL2 enhances angiogenesis as well as neutrophil accumulation, ultimately inducing carcinogenesis [247]. Several studies have shown that G-CSF may stimulate angiogenesis and promote tumor growth [279–281]. This effect of G-CSF is mediated through recruitment of CD11b<sup>+</sup>Gr1<sup>+</sup> MDSCs and increased number of endothelial progenitor cells [279]. The pro-angiogenic effects of MDSCs is in part driven by Bv8

(prokineticin 2/PK2), which is upregulated by G-CSF [282, 283] through a STAT3 dependent mechanism [284]. G-CSF may further contribute to tumor angiogenesis by inducing VEGF-A production in neutrophils [285]. The neutrophil secreted MMP-9 has been functionally implicated in VEGF activation [286], thus further fortifying angiogenesis. A cooperation between cancer cells and neutrophils in promoting angiogenesis was demonstrated by Queen et al. [4], where GM-CSF secreted by breast cancer cells stimulated neutrophils to produce oncostatin M, which in turn stimulated the tumor cells to produce VEGF, thereby augmenting tumor-associated angiogenesis.

IL-17 produced by Th17 cells induces expression of G-CSF, leading to immature myeloid-cell mobilization and recruitment into the tumor microenvironment [231]. These CD11b<sup>+</sup>Gr1<sup>+</sup> MDSCs produce the pro-angiogenic Bv8 [283], that bypasses VEGF and renders tumors refractory to anti-VEGF therapy [231]. Anti-Bv8 treatment reduced the number of CD11b<sup>+</sup>Gr1<sup>+</sup>

cells in peripheral blood and in tumors along with suppression of angiogenesis [283]. Thus, G-CSF may induce angiogenesis through a Bv8-dependent mechanism.

The important role of neutrophils in the induction of the angiogenic switch in cancer, was further illustrated in a RIP1-Tag2 transgenic mouse model of pancreatic  $\beta$ -cell carcinogenesis [287]. MMP-9-expressing neutrophils were predominantly found in angiogenic islet dysplasias and in tumors, whereas MMP-9-expressing macrophages were localized along the periphery of these lesions [287]. Transient depletion of neutrophils reduced the frequency of initial angiogenic switching in islet dysplasias. Jablonska et al. [64] observed that IFN $\beta$ -deficient mice develop faster growing melanoma and fibrosarcoma tumors with better developed blood vessels than did wild-type mice. These tumors displayed enhanced infiltration of CD11b<sup>+</sup>Gr1<sup>+</sup> neutrophils that expressed high levels of VEGF, MMP-9 and CXCR4. The transcription factors c-Myc and STAT3 regulating the expression of these proteins, were also elevated in the neutrophils [64]. Thus, endogenous IFN $\beta$  seems to inhibit tumor angiogenesis through repression of genes encoding pro-angiogenic and homing factors in tumor-infiltrated neutrophils [64]. Neutrophils in bronchioloalveolar carcinoma produce hepatocyte growth factor (HGF) that stimulates the migration of the tumor cells [203]. Similarly, human cholangiocellular and hepatocellular carcinoma cells induce HGF secretion by neutrophils, which in turn enhances the invasiveness of the cancer cells [288]. TANs can also stimulate tumor growth by releasing growth factors such as epidermal growth factor, TGF $\beta$  and platelet-derived growth factor (PDGF) from the extracellular matrix [289]. The important role neutrophils play at the primary tumor site was further demonstrated using neutrophil depletion experiments in mice. In neutrophil depleted mice tumor growth was significantly reduced supporting the notion of pro-tumorigenic neutrophil function at the primary tumor site [287, 264]. While these observations clearly demonstrate a pro-tumor function for neutrophils in the tumor-microenvironment, there are several convincing observations showing the opposite (See “Anti-Tumor Functions of Neutrophils” section). The complexity of the mechanisms that regulate neutrophil function in the primary tumor microenvironment is exemplified in Fig. 2.

#### Promotion of Cancer Cell Dissemination by Neutrophils

There are several studies showing that neutrophils promote tumor cell motility, migration and invasion [146]. Neutrophils may facilitate invasion through secretion of enzymes degrading the extracellular matrix such as elastase, Cathepsin G, proteinase-3, MMP-8 and MMP-9 [146]. Neutrophils may also indirectly contribute to the degradation of the extracellular matrix through activation of tumor-derived pro-MMP-2 (pro-Gelatinase A), mediated by the neutrophil-derived elastase, cathepsin G and proteinase-3 [290].

IL-8-producing tumor cells have frequently been shown to be more metastatic than the corresponding non-producer cells [291]. IL-8 is a potent chemoattractant for neutrophils and normally recruits neutrophils to the site of wounds. However, in the tumor settings, IL-8 induces the release of specific proteases and heparanases by recruited neutrophils. These proteases and heparanases remodel the extracellular matrix making it easier for tumor cells to intravasate. Neutrophil elastase activates other latent proteases in the tumor microenvironment which cleave and inactivate plasminogen activator inhibitor-1, leading in turn to the release of embedded growth factors such as bFGF, a potent angiogenic factor [291]. An interesting interplay between neutrophil elastase and its inhibitor alpha 1-antitrypsin seems to have a significant impact on cancer development [292]. A deficiency in alpha 1-antitrypsin is associated with increased risk of liver cancer, bladder cancer, gall bladder cancer, malignant lymphoma and lung cancer [292]. Conversely, elevated concentrations of neutrophil elastase might promote the development, invasion, and metastasis of many cancers [292]. Neutrophils also release high levels of MMP-9/Gelatinase B, which further affects extracellular matrix remodelling. HOCl produced by myeloperoxidase of activated neutrophils oxidizes specific sulfur-containing amino acids and activates the MMP pro-enzymes. In parallel, HOCl inactivates TIMP-1, thereby further increasing the proteolytic activity of MMPs [291].

Neutrophils activated by melanoma-secreted IL-8 can facilitate melanoma cell extravasation through an interaction between ICAM-1 (CD54) on the melanoma cells and Mac-1 (CD11b/CD18) on neutrophils [253, 254, 293]. Neutrophils were also shown to be important for entrapping circulating melanoma cells and thus facilitating lung metastasis development [294]. Melanoma-secreted IL-8 not only attracts neutrophils, but also strongly upregulates  $\beta$ 2 integrins, which interact with ICAM-1 on melanoma cells, thereby promoting the anchoring of the tumor cells to vascular endothelium [294]. Other studies further show that IL-8 increases neutrophil  $\beta$ 2 integrin expression with simultaneous shedding of L-selectin (CD62L), thereby promoting their sequestration in the lung [295]. Similarly, neutrophils were shown to promote breast cancer cell transendothelial migration through an ICAM-1-CD11b/CD18 interaction dependent mechanism [296]. Neutrophils were also shown to interact with colon carcinoma cells, facilitating their dissemination [297]. The interaction was mediated by CD11b, CD11a and L-selectin on neutrophils and CD54 and sialylated, O-linked, protease-sensitive ligands on the tumor cells [297, 298].

#### *Neutrophils in Pre-Metastatic Organs - The “Seed and Soil” Hypothesis*

In 1889, Stephen Paget proposed that tumors do not metastasize to random organs and that there are preferable sites of metastasis for specific tumors. This has led to the realization

that successful colonization of a distant organ requires a degree of compatibility between the tumor cell and the future site of metastasis. This realization was the basis of the “Seed and soil” theory. This theory was expanded upon when tumors were shown to directly enhance the seeding of circulating cells in the lungs [299], suggesting that the tumor can modulate the “soil” to make it more compatible with the “seed”. Later studies have shown the direct involvement of bone marrow derived VEGFR1<sup>+</sup> progenitor cells [300, 301] in generating the pre-metastatic niche that is more receptive toward incoming tumor cells. These cells were identified as of myeloid lineage, and appeared not to differentiate, but maintained their expression of immature surface markers including c-Kit and Sca-1 within the tissue parenchyma [300]. Of note, the formation of the pre-metastatic niche was found to be organ specific. Yan et al. [302] observed that CD11b<sup>+</sup>Gr-1<sup>+</sup> myeloid cells are increased in lung of mice bearing mammary adenocarcinomas before tumor cell arrival. These immature myeloid cells decreased IFN $\gamma$  production, but increased the production of pro-inflammatory cytokines in the premetastatic lung [302]. In addition, the CD11b<sup>+</sup>Gr-1<sup>+</sup> cells produce large amounts of MMP-9 and promote vascular and ECM remodeling [302]. Deletion of MMP-9 normalized aberrant vasculature in the pre-metastatic lung and diminished lung metastasis [302]. More recently, Gao and colleagues [303] showed that it was the CD11b<sup>+</sup>Ly6C<sup>high</sup> monocytic myeloid subpopulation that was promoting metastatic seeding in the lungs rather than the neutrophilic CD11b<sup>+</sup>Ly6G<sup>high</sup> subpopulation. Interestingly, the pro-metastatic effect of CD11b<sup>+</sup>Ly6C<sup>high</sup> monocytes at the pre-metastatic lung was found to be mediated by secretion of versican which in turn leads to mesenchymal to epithelial transition (MET) and to enhanced proliferation [303]. Still, while these studies do not implicate neutrophils in the process of priming the pre-metastatic niche, the fact the neutrophil chemoattractants such as S100A8 and S100A9 [301] are available in the pre-metastatic lung, together with our own observations [32] showing neutrophils accumulating early in the pre-metastatic lung, support the notion that neutrophils may contribute to the formation of the pre-metastatic niche. Furthermore, Kowanz et al. [26] showed that depletion of Gr1<sup>+</sup> or Ly6G<sup>+</sup> cells from the pre-metastatic lung results in reduced metastasis. Similarly, Sceneay et al. [304] also observed increased numbers of granulocytic CD11b<sup>+</sup>Ly6C<sup>med</sup>Ly6G<sup>+</sup> myeloid cells in the pre-metastatic lungs in mice injected with melanoma or breast carcinoma cells with a concomitant reduction in the cytotoxic activity of NK cells [304]. Erler et al. [305] observed that lysyl oxidase (LOX) secreted by hypoxic breast tumor cells is required for the recruitment of immature myeloid CD11b<sup>+</sup>F4/80<sup>-</sup> cells to the premetastatic lung. Tumor-secreted LOX was found co-localized with fibronectin in the lung, to which CD11b<sup>+</sup> cells bound. The CD11b<sup>+</sup> cells secreted MMP-2, generating chemoattractive collagen IV peptides, a process leading to enhanced invasion and recruitment of metastasizing

tumor cells as well as bone-marrow-derived cells into the lung [305]. Knocking down LOX in the breast tumor cells prevented the recruitment of CD11b<sup>+</sup> cells to the premetastatic lung, with concomitant reduction in metastatic growth [305].

A mechanism involving microvascular deposition of neutrophil extracellular traps (NETs) was found to directly facilitate the formation of hepatic micrometastases [306]. Neutrophils were also shown to promote liver metastasis of Lewis lung carcinoma (H-59 subline) cells through Mac1 (CD11b/CD18)-mediated interaction with circulating tumor cells [307]. Neutrophil depletion prior to cancer cell inoculation reduced the number of liver metastases [307]. This effect was reversed when inflamed neutrophils were co-inoculated with tumor cells [307]. H-59 cells showed reduced adhesion to liver sinusoids of CD11b deficient mice [307]. In contrast, our own work, using mouse models of breast cancer, shows that neutrophils are stimulated by tumor-secreted factors and acquire an anti-tumor phenotype [32]. These neutrophils accumulate in the lungs in large numbers during the pre-metastatic phase and provide anti-metastatic protection by eliminating incoming disseminated tumor cells [32]. These tumor-entrained neutrophils (TENs) produced high levels of H<sub>2</sub>O<sub>2</sub> and acquire the capacity to kill tumor cells in a contact dependent fashion [32]. Similarly, Lopez-Lago and colleagues [66] showed that tumor-secreted CXCL1 stimulates neutrophils and induces an anti-metastatic response in a model of human renal cell carcinoma. Furthermore, they demonstrated that the metastatic potential of the tumors inversely correlates with the extent of neutrophil mobilization [66]. Finally, we have shown that tumoricidal neutrophils also exist in the circulation of breast cancer patients, but not in healthy individuals, suggesting that anti-tumor neutrophils are generated during the natural course of the disease in patients [32]. These conflicting observations provide the basis for the controversy that surrounds the function of neutrophils at the pre-metastatic niche; do they possess favorable or unfavorable properties? The answer to this question remains obscure. However, it seems that neutrophils may possess both pro- and anti-tumor functions and their actual function in situ is determined by the microenvironment and chemokine milieu (Figs. 1 and 2).

#### Anti-Tumor Functions of Neutrophils

The recognition that neutrophils have potential anti-tumor functions was first brought forward in the 1970s where neutrophils from patients with bladder cancer were shown to be cytotoxic toward bladder cancer cells [308]. Since then neutrophils were shown to exert their anti-tumor functions via direct cytotoxicity, antibody dependent cell mediated cytotoxicity (ADCC) and through the presentation of specific antigens (Table 2) [7]. Neutrophils need to be activated in order to exert their anti-tumor activities. Various cytokines (e.g., G-CSF, IFN $\gamma$ , TNF $\alpha$ ) and chemokines (e.g., CCL-2, CCL-5,



**Table 2** Evidence for anti-tumorigenic effects of neutrophils

Major findings	Reference
<b>Human neutrophils</b>	
• Normal human peripheral blood granulocytes destroy various human cancer cells in vitro, including osteosarcoma, melanoma and lung squamous carcinoma cells, at a higher efficiency than normal fibroblasts.	[392]
• Leukocytes from patients with urinary bladder carcinomas showed cytotoxicity toward human bladder carcinoma cells in vitro.	[308]
• Leukocytes from pregnant women, and patients with toxemia, uterine myoma, ovarian and endometrial carcinoma had cytotoxic effects toward primary ovarian carcinoma cultures.	[393]
• Neutrophils from Stage III and IV lung and stomach cancer patients were more cytotoxic to K562 cells than those from healthy controls.	[386]
• Normal human peripheral blood granulocytes showed cytostatic effect on various tumor cell lines including human K562 myeloid leukemia, human F265 lymphoblastoid, mouse TU-5 kidney cells, mouse RBL-5 lymphoma, human RAJI Burkitt's lymphoma, human W1-38 embryonal lung line, human SV40 transformed fibroblasts.	[394]
• The neutrophils were cytolytic to Chang-A hepatocarcinoma cells, an effect that was enhanced by antibodies.	
• Granulocytes formed clusters among themselves and attached to the target cells.	
• Detachment of adherent cells was observed after 6 h.	
• PMA-stimulated neutrophils mediated cytotoxicity against CEM T-lymphoblast cells.	[314]
• Neutrophil cytotoxicity was mediated by H <sub>2</sub> O <sub>2</sub> and myeloperoxidase.	
• Killing of CEM could be inhibited by catalase, the MPO inhibitor cyanide, and the HOCl scavengers tryptophan, methionine and alanine, suggesting a role for the MPO-H <sub>2</sub> O <sub>2</sub> -Cl <sup>-</sup> system.	
• The cytotoxic effect was dependent on pH and the effector cell number.	
• Concanavalin A-activated neutrophils can release cytotoxic quantities of H <sub>2</sub> O <sub>2</sub> and myeloperoxidase, which in concert with a halide (chloride or iodide) lysed murine LSTRA lymphoma cells.	[315]
• SOD did not affect killing.	
• Unstimulated neutrophils had no effect.	
• Azide, cyanide and catalase prevented the cytotoxicity.	
• Neutrophils were cytotoxic and cytostatic to human tumor cell lines T24 bladder carcinoma, LR melanoma, and SV40-transformed fibroblasts, as well as K562 CML, Raji Burkitt lymphoma and CEM T-ALL cells.	[395]
• The killing of tumor cells was selective, as the neutrophil didn't kill normal human fibroblasts.	
• Protease inhibitors did not inhibit neutrophil cytotoxicity.	
• PMA-activated granulocytes are cytotoxic to CEM T-ALL cells.	[316]
• The myeloperoxidase-hydrogen peroxide-halide system is involved as azide, cyanide and catalase inhibited the killing.	
• The killing was dependent on the presence of a halide.	
• PMA-activated neutrophils are cytotoxic to Raji Burkitt lymphoma cells when incubated at 20:1 E:T ratio.	[317]

**Table 2** (continued)

Major findings	Reference
• The myeloperoxidase-hydrogen peroxide-halide system is involved as the cytotoxicity can be inhibited by azide, cytochrome C and catalase.	
• Superoxide dismutase enhanced the cytotoxic effect.	
• Neutrophils had a stronger cytotoxic effect on malignant targets (MA-160 prostate cancer, Garr1 colon cancer) than on nonmalignant targets (human embryonic lung and intestinal cells).	[396]
• Neutrophils from colon and breast cancer patients with stage I disease possessed neutrophils which were less effective in killing tumor cells than neutrophils from normal donors.	
• In contrast, neutrophils from colon and breast carcinoma patients with stage IV disease were more effective in killing tumor cells than normal cells.	
• PMA-activated neutrophils kill K562 CML cells. Neutrophil cytotoxicity was inhibited by catalase, while augmented by SOD.	[318]
• Lidocaine enhanced, while verapamil and exogenous adenosine 5'-triphosphate inhibited neutrophil-mediated tumor cytotoxicity.	[397–400]
• Prednisolone inhibited, while chloroquine had no effect on neutrophil-mediated tumor cytotoxicity.	
• TNF-treated neutrophils acquire cytotoxic activity towards tumor cells such as Raji, K562, UCLA-SO-M14 and U937, through production of hydrogen peroxide.	[309, 310]
• Catalase, but not SOD, sodium azide or deferoxamine, prevented the cytotoxic effect.	
• IFN $\gamma$ enhances the tumor cytostatic effect of neutrophils.	[311]
• Neutrophils induce ADCC of human GD2 <sup>+</sup> melanoma and neuroblastoma cell lines in the presence of antibodies to the ganglioside GD2.	[336]
• CD11/CD18-deficient neutrophils were defective in inducing cytolysis.	
• Antibodies to CD11b, CD11c and CD18 blocked ADCC by normal neutrophils.	
• Antibodies to FcRII and FcRIII, but not those to FcRI, blocked ADCC.	
• GM-CSF enhances anti-tumor neutrophil ADCC.	
• K652 cells that have been made resistant to H <sub>2</sub> O <sub>2</sub> through repeated exposure to increasing amounts of glucose oxidase, expressed elevated levels of Catalase and were resistant to the cytotoxic effect of TNF $\alpha$ -activated neutrophils.	[401]
• Neutrophils from colon carcinoma and breast cancer patients showed elevated cytotoxicity towards Hct-116 colon carcinoma cells, while normal neutrophils or neutrophils from melanoma patients had almost no effect.	[402]
• The cytotoxicity of neutrophils from colon carcinoma could be enhanced by fMLP and prothymosin $\alpha$ 1.	
• Prothymosin $\alpha$ 1 enhanced the oxidative responses of neutrophils.	
• GM-CSF-primed neutrophils reduced melanoma cell viability.	[325]
• These neutrophils showed increased release of O <sub>2</sub> <sup>-</sup> .	
• SOD and Catalase when applied separately could not abrogate the neutrophil killing of melanoma cells, while added together partly prevented the killing.	
• Interfering with nitric oxide (NO <sup>•</sup> ) production by N <sup>G</sup> -monomethyl-L-arginine significantly protected the melanoma cells against GM-CSF-primed neutrophils.	

**Table 2** (continued)

Major findings	Reference
<ul style="list-style-type: none"> <li>Neutrophils are cytotoxic to HT-29 colorectal cancer and FaDu pharyngeal squamous-cell carcinoma cells. These tumor cells secrete IL-8 and GRO<math>\alpha</math>, leading to the activation of cytotoxic neutrophils.</li> </ul>	[216]
<ul style="list-style-type: none"> <li>Tumor-entrained neutrophils (TENs) with anti-tumor activities are present in the peripheral blood of breast cancer patients prior to surgical resection, but not in healthy individuals.</li> </ul>	[32]
<ul style="list-style-type: none"> <li>These TENs showed cytotoxic effect towards MDA-MB-231 breast carcinoma cells.</li> </ul>	
<ul style="list-style-type: none"> <li>Human neutrophils display a higher cytotoxic activity against poorly metastatic cells compared with highly metastatic renal cell carcinoma cells.</li> </ul>	[66]
<ul style="list-style-type: none"> <li>Several neutrophil chemokines including CXCL1, CXCL2, CXCL3, CXCL5 and IL-8 were down modulated in the highly metastatic cells.</li> </ul>	
<ul style="list-style-type: none"> <li>Non-metastatic renal carcinoma cells promote recruitment of neutrophils to the lung of mice. The metastatic activity inversely correlated with the ability of tumor cells to recruit and activate these immune cells.</li> </ul>	
<b>Murine neutrophils</b>	
<ul style="list-style-type: none"> <li>Neutrophils are attracted by rat Walker Carcinoma 256 cells.</li> </ul>	[403]
<ul style="list-style-type: none"> <li>Neutrophils interacts with these cells, indent their membranes, leading to a rounding up of the tumor cells. This killing process depends on the presence of other host factors, as in vitro the tumor cells did not induce cytotoxic neutrophils.</li> </ul>	
<ul style="list-style-type: none"> <li>Cytotoxic effect of rat peritoneal neutrophils against the syngeneic ascites tumor WBP1.</li> </ul>	[404]
<ul style="list-style-type: none"> <li>Establishment of WBP1 ascites tumor was retarded when the rats were stimulated to produce large amounts of peritoneal neutrophils by intraperitoneal injections of beef heart infusion broth in combination with proteose peptone.</li> </ul>	
<ul style="list-style-type: none"> <li>Granulocytes from tumor-bearing mice were cytotoxic.</li> </ul>	[405]
<ul style="list-style-type: none"> <li>PMA-stimulated BCG- or thioglycollate-elicited peritoneal neutrophils kill efficiently P388 and TLX9 lymphoma cells, which correlated with H<sub>2</sub>O<sub>2</sub> release.</li> </ul>	[406]
<ul style="list-style-type: none"> <li>Cell lysis peaked after 4.5 h following co-cultivation with activated neutrophils.</li> </ul>	
<ul style="list-style-type: none"> <li>Increase in the proportion of myeloperoxidase-positive neutrophils in mice bearing autochthonous M-MuSV (Moloney murine sarcoma virus)-induced tumors.</li> </ul>	[407]
<ul style="list-style-type: none"> <li>The cytotoxic activity correlated with myeloperoxidase expression.</li> </ul>	
<ul style="list-style-type: none"> <li>Ha-2 Harvey-MuSV-induced sarcoma and Tu5 Simian virus-40-transformed kidney cells were highly susceptible to the cytotoxic effect of neutrophils (40–60 % killing), while MBA 3-methylcholantrene-induced sarcoma, Ta<sub>3</sub>/St 3-methylcholantrene-induced adenocarcinoma, T1699 spontaneous adenocarcinoma and 3 T3 embryonic fibroblasts showed medium sensitivity (20–40 % killing).</li> </ul>	
<ul style="list-style-type: none"> <li>The cytotoxic activity of neutrophils seems to be non-specific, leading to killing of various tumor cell lines as well as allogeneic, but not syngeneic, fibroblasts. Although the extent of killing of allogeneic mice was much lower (5–20 %) than tumor cells (20–60 %).</li> </ul>	
<ul style="list-style-type: none"> <li>Total bone marrow derived cells were much more cytotoxic than spleen-derived cells. Cells from the lymph nodes were not cytotoxic.</li> </ul>	

**Table 2** (continued)

Major findings	Reference
<ul style="list-style-type: none"> <li><math>\beta</math>-1,3-glucan, BCG, Propionibacterium acnes and Zymosan A induced anti-tumor activity of neutrophils toward MM46 mammary carcinoma, MM48 mammary carcinoma, MH134 hepatoma, EL-4 and YAC-1 lymphoma cells.</li> </ul>	[408, 409]
<ul style="list-style-type: none"> <li>Catalase, but not superoxide dismutase, cyanide or azide, inhibited the killing of tumor cells.</li> </ul>	
<ul style="list-style-type: none"> <li>Antibody to TNF prevented the cytolytic effect of caseinate-induced inflammatory neutrophils toward MM46 mammary tumor cells.</li> </ul>	[332]
<ul style="list-style-type: none"> <li>Immature neutrophils obtained from ascites fluid 6 h after caseinate injection showed stronger cytotoxic activity than mature neutrophils obtained 3 h after injection.</li> </ul>	
<ul style="list-style-type: none"> <li>Myeloperoxidase is expressed 1.6 times more in immature neutrophils.</li> </ul>	
<ul style="list-style-type: none"> <li>Intraperitoneal injection of <i>Corynebacterium parvum</i> 24 h after an intraperitoneal inoculation of a lethal number of mouse ovarian teratocarcinoma cells induced an antitumor response that cured 75 to 95 % of the mice.</li> </ul>	[410]
<ul style="list-style-type: none"> <li>Isolated peritoneal neutrophils were cytolytic to the tumor cells with cell lysis obtained within 30 min after binding of the neutrophils to the ovarian teratocarcinoma target cells.</li> </ul>	
<ul style="list-style-type: none"> <li>Overexpression of chemokines such as IL-8, MCP-1 or MIP-1 in CHO tumor cells led to recruitment of neutrophils and concomitant inhibition of tumor growth in nude mice.</li> </ul>	[411]
<ul style="list-style-type: none"> <li>IFN<math>\gamma</math> activates rat neutrophils to kill tumor cells by a mechanism dependent on nitric oxide.</li> </ul>	[312, 313]
<ul style="list-style-type: none"> <li>SOD enhanced the tumor cytotoxic effect.</li> </ul>	
<ul style="list-style-type: none"> <li>The growth of disialoganglioside (GD2) positive neuroblastoma cells was inhibited by neutrophils in the presence of antibodies towards GD2, an effect that was enhanced by GM-CSF.</li> </ul>	[412]
<ul style="list-style-type: none"> <li>In the absence of antibodies, neutrophils inhibited growth of one GD2<sup>+</sup> cell line, whereas they stimulated the growth of two other GD2<sup>+</sup> cell lines as well as the GD2<sup>-</sup> cell lines tested.</li> </ul>	
<ul style="list-style-type: none"> <li>Mouse granulocytes restrict human tumor cell growth in SCID mice.</li> </ul>	[413]
<ul style="list-style-type: none"> <li>Neutrophils, together with macrophages, mediate antibody-dependent cell cytotoxicity (ADCC) towards tumor cells, which is responsible for the efficacy of monoclonal antibody (mAb)-mediated cancer therapy.</li> </ul>	[342]
<ul style="list-style-type: none"> <li>Neutrophil-induced ADCC contributes to the anti-tumor activity of the anti-CD20 antibody Rituximab in a non-Hodgkin's lymphoma SCID mouse model.</li> </ul>	[341]
<ul style="list-style-type: none"> <li>Walker carcinoma W256 activated neutrophils in vitro to produce singlet oxygen.</li> </ul>	[326]
<ul style="list-style-type: none"> <li>ROS is crucial for neutrophil-mediated tumor cell lysis.</li> </ul>	
<ul style="list-style-type: none"> <li>Massive granulocyte infiltration at the site of W256 transplants correlates with spontaneous tumor regression.</li> </ul>	[414]
<ul style="list-style-type: none"> <li>Peripheral blood granulocytes from the tumor-bearing animals are cytotoxic to W256 cells in vitro.</li> </ul>	
<ul style="list-style-type: none"> <li>Adoptive transfer of granulocytes in the vicinity of W256 carcinoma in rats or Ehrlich ascites tumor in mice reduced tumor cell mass with concomitant increased survival.</li> </ul>	[415]
<ul style="list-style-type: none"> <li>The presence of MPO in the tumor microenvironment was accompanied by the formation of lipid peroxidase</li> </ul>	[416]

**Table 2** (continued)

Major findings	Reference
(LPO)-derived aldehydes such as acrolein, 4-hydroxy-2-nonenal and malondialdehyde.	
• The presence of acrolein and neutrophil elastase were increased in animals with regressing W256 tumor.	
• Meth A tumor cells induce infiltration of cytotoxic neutrophils, which are responsible for the IFN $\gamma$ -dependent spontaneous rejection.	[417]
• Inhibition of TGF $\beta$ signaling by using SM16 increased neutrophil mediated cytotoxicity towards AB12 mesothelioma cells in a mechanism that depends on ROS.	[13]
• While antibodies to TNF $\alpha$ and N-methylarginine, an inhibitor of iNOS, did not inhibit the cytotoxicity, blockade of superoxide and H $_2$ O $_2$ by superoxide dismutase (SOD) and catalase, respectively, blocked neutrophil cytotoxicity.	
• CD11b <sup>+</sup> MMP-9 <sup>+</sup> Neutrophils accumulate in the lungs prior to metastatic seeding of breast carcinoma cells.	[32]
• Tumor-entrained neutrophils (TENs) inhibit metastatic seeding of 4T1 breast carcinoma cells in the lungs by generating H $_2$ O $_2$ .	
• Also, neutrophil depletion in MMTV-PyMT/MMTV-cMyc tumor bearing mice resulted in enhanced metastatic seeding in the lungs.	
• TENs could not prevent the growth of local tumor.	
• Circulating blood neutrophils from 4T1-tumor bearing mice were cytotoxic to the mouse 4T1 and the human MCF7 breast carcinoma cell lines. Similarly, circulating blood neutrophils from B16 melanoma-bearing tumor mice were cytotoxic to both B16 and 4T1 cells.	
• Catalase or TGF $\beta$ (at 200pM) abrogated the cytotoxic effects of TENs.	
• Neutrophil-tumor cell contact was required for tumor cytolysis.	
• Tumor-secreted CCL2 mediates the anti-metastatic entrainment of G-CSF-stimulated neutrophils.	
• In vitro CCL2 and CCL5 simulated naïve neutrophils to produce H $_2$ O $_2$ and to kill 4T1 tumor cells.	
<b>Purified neutrophil products</b>	
• Myeloperoxidase, a product of neutrophils, shows cytotoxic effects against mouse lymphoma cells.	[418, 419]
• Cationic proteins purified from polymorphonuclear leukocytes granules exert a cytotoxic effect on mouse L8TRA lymphoma cells.	[420]
• A ~100kD protein secreted from wheat germ agglutinin or actinomycin D-stimulated neutrophils kills MM46 and MM48 mammary adenocarcinoma cells, MH134 hepatoma cells and L929 fibrosarcoma cells, but not normal splenocytes.	[421]
• No cytotoxic effect of supernatant of unstimulated neutrophils.	
• Human defensins HNP-1, HNP-2 and HNP-3 lysed various human and mouse tumor cell lines.	[335]
• Human defensins in combination with hydrogen peroxide had a synergistic cytotoxic effect on tumor cells.	
• Calprotectin (S100A8/S100A9), a calcium binding protein complex abundantly expressed in neutrophils, induces apoptotic cell death in various tumor cells.	[422]

Evidence for neutrophil mediated tumor cell killing in cancer patients, in animal models of cancer and in using neutrophil derived products

CXCL5 and IL-8) may promote the generation of anti-tumorigenic neutrophils [32, 66, 309–313]. Anti-tumorigenic neutrophils can also be generated artificially by exposing them to the phorbol ester PMA, or to the lectins Concanavalin A or wheat germ agglutinin (WGA) [314–318].

In vivo, the effect of massive infiltration of neutrophils was analyzed by sustained treatment of tumor-bearing mice with GM-CSF, leading to a strong neutrophilia around the tumors. These mice showed a 16-fold lower mortality rate than untreated mice [319]. The same therapeutic strategy was applied in a patient with advanced hepatocarcinoma, who exhibited a complete remission after 4 months of G-CSF treatment. Prolonged administration of G-CSF to squamous head and neck cancer patients led to increased disease-free survival [320]. Other clinical trials using continuous GM-CSF administration in advanced prostate cancer [321] or sustained G-CSF in stage IV melanoma with brain metastases [322] reported better survival, suggesting for an anti-tumor effects of neutrophils.

#### Direct Cytotoxicity

Neutrophils are armed with a variety of toxic molecules, most of which have anti-microbial properties and are harmless toward eukaryotic cells [323, 324]. Still, several anti-bacterial molecules are also involved in neutrophil cytotoxicity toward tumor cells. Of special importance are the reactive oxygen species H $_2$ O $_2$  and HOCl, generated during an “oxidative burst” that is mediated by the NADPH oxidase complex and by myeloperoxidase, respectively [325–327]. These molecules are directly involved in the anti-tumor activity [325–327]. Inhibition of myeloperoxidase by azide or cyanide, or addition of catalase that catalyzes the conversion of hydrogen peroxide to water and oxygen, prevents neutrophil-mediated tumor cell killing [13, 32, 309, 314–318]. Neutrophils from patients with myeloperoxidase (MPO) deficiency or defective H $_2$ O $_2$  production are not cytotoxic to tumor cells [327]. However, addition of superoxide dismutase (SOD) that catalyzes the conversion of superoxide into oxygen and hydrogen peroxide, did not prevent the killing, but sometimes even enhanced it [13, 32, 309, 314–318], possibly by accelerated hydrogen peroxide production, the key effector molecule in neutrophil cytotoxicity. Dissemmond and colleagues showed that SOD in combination with catalase could partly prevent tumor cell killing with an involvement of nitric oxide in the killing process [325]. Rat neutrophils stimulated with IFN $\gamma$  produced nitric oxide that prevented tumor cell growth [312] and induced apoptosis of the tumor cells [313]. However, another study [13] could not relieve the cytotoxic effect with a nitric oxide scavenger. This apparent discrepancy could be due to different neutrophil activation modes and the presence of several neutrophil generated cytotoxic molecules where the inhibition of one is compensated for by the other.

Several studies, including our own, have shown that neutrophil cytotoxicity requires physical contact between neutrophils and tumor cells [4, 32, 328]. Caruso et al. [329] demonstrated intimate contact between neutrophils and tumor cells in samples of early gastric cancer at a stage prior to the appearance of neutrophilia. Following the interaction with neutrophils, tumor cells show varying degrees of damage including disorganization of the intermediate filaments and dilation of the rough endoplasmic reticulum [329]. However, neutrophil cytotoxicity induced by the phorbol ester PMA, a potent agonist of  $H_2O_2$  production and secretion, does not require physical contact [32]. This observation suggests that high levels of neutrophil-generated  $H_2O_2$  are sufficient for cytotoxicity, even when physical contact is restricted. These seemingly conflicting observations were resolved when we realized that rather than secreting  $H_2O_2$  spontaneously, neutrophils secrete  $H_2O_2$  in a contact-triggered fashion [32]. Similarly, Saito et al. [330] demonstrated by visualizing the oxidative process by luminol-dependent chemiluminescence, that ROS accumulate at sites where neutrophils come in contact with tumor cells. Also, the intensity of the hydrogen peroxide-sensitive tracer dichlorofluorescein diacetate preloaded in tumor cells rapidly increased after adding the neutrophils [330]. The interaction between neutrophils and tumor cells results in loss of tumor cell membrane integrity beginning within 15 min of the binding step and completion of the lytic event within 45 min [328]. It is noteworthy to point out that molecules other than  $H_2O_2$ , HOCl and  $NO\cdot$  may also mediate the tumoricidal effects of neutrophils. These include proteases [331], membrane-perforating agents [331],  $TNF\alpha$  [331, 332], TRAIL [333] and defensins [334, 335].

#### *Antibody-Dependent Cell-Mediated Cytotoxicity (ADCC)*

Another mechanism for neutrophil-mediated tumor cell killing is antibody-dependent cell-mediated cytotoxicity (ADCC) [7, 331] where specific antibodies are used to target malignant cells. Neutrophils express several subtypes of FcRs capable of inducing ADCC, including  $Fc\gamma RI$  (CD64),  $Fc\gamma RIIa$  (CD32),  $Fc\gamma RIIIa$  (CD16a), and  $Fc\gamma RIIIb$  (CD16b) [336–338], whose surface expression is increased following G-CSF stimulation [338, 339].  $Fc\gamma RI$  was shown to mediate neutrophil ADCC activity against glioma, squamous cell and ovarian carcinoma cells [338]. However, binding of IgG to the inhibitory  $Fc\gamma RIIIb$ , might lead to down-regulation of immune responses [340]. The neutrophil-mediated killing of human ganglioside  $GD2^+$  melanoma and neuroblastoma cells in the presence of antibodies to  $GD2$ , was found to depend on  $Fc\gamma RII$  and  $Fc\gamma RIII$ , but not on  $Fc\gamma RI$  [336]. Neutrophils from a child with leukocyte adhesion deficiency (LAD) devoid of CD11/CD18 adhesion molecules failed to mount any detectable ADCC [336], suggesting the involvement of these

adhesion molecules in ADCC. Also, antibodies to CD11b, CD11c and CD18 efficiently blocked ADCC by normal neutrophils, providing further support to this notion [336]. GM-CSF enhanced the anti-tumor neutrophil-dependent ADCC through enhanced expression of CD11/CD18 molecules [336]. Neutrophils were also shown to contribute to the anti-tumor ADCC in Non-Hodgkin's Lymphoma using antibodies to CD20 (Rituximab) [341], in breast cancer using a Tn antigen-specific chimeric mAb [342], and in B-cell lymphoma using a bispecific single-chain fragment variable-specific for HLA class II and  $Fc\alpha RI$  (CD89) [343]. The bispecific antibody against the myeloid receptor for IgA ( $Fc\alpha RI$ ; CD89) and the B-cell surface marker CD20 induced neutrophil-dependent ADCC toward broad range of B cell lines [344]. Lysis via  $Fc\alpha RI:CD20$  bispecific antibodies was enhanced in blood from patients during therapy with G-CSF or GM-CSF [344]. Interestingly, Otten et al. [345] observed that immature neutrophils mobilized from the bone marrow upon G-CSF treatment, efficiently triggered tumor cell lysis via  $Fc\alpha RI$  (CD89), but were unable to initiate tumor cell killing via  $Fc\gamma R$ . This may provide a rationale for the disappointing results observed in some earlier clinical trials in which patients were treated with G-CSF and anti-tumor antibody binding to  $Fc\gamma R$ . An indication for the role neutrophils play in mediating ADCC in human patients was described by Cheung and colleagues [346]. In their study they found that the outcome of treating high risk neuroblastoma patients with anti- $GD2$  monoclonal antibody strongly correlated with the extent of granulocyte activation [346].

#### *Neutrophils as Tumor-Antigen Presenting Cells*

There is accumulating evidence suggesting that neutrophils contribute to the development of an adaptive anti-tumor immune response [12–16, 347]. Elimination of neutrophils prevents the mounting of anti-tumor  $CD8^+$  T cytotoxic responses [12–14, 16]. Neutrophils were shown to be required for priming rats with tumor-associated antigens to induce anti-tumor  $CD8^+$  effector T cells [348]. This may be related to the ability of neutrophils to attract and activate dendritic cells, macrophages, NK cells and T cells. By virtue of their ability to present antigens, neutrophils may directly activate a T cell response. Neutrophils may also activate dendritic cells and T cells through release of neutrophil extracellular traps (NETs) [349, 350]. On top of secreting T cell chemoattractants, neutrophil-secreted  $TNF\alpha$ , Cathepsin G and neutrophil elastase, are able to increase T cell proliferation and cytokine production, which together enhance adaptive immune responses [350]. Jackaman et al. [351] demonstrated the cooperation between tumor-infiltrating neutrophils and  $CD8^+$  T



cells in eradicating tumors that received intratumoral injection of IL-2 and anti-CD40 antibodies.

### Prognostic Values of Neutrophils and Other Myeloid Subtypes in Cancer Patients

Alterations in circulating leukocyte composition and number are often observed in cancer patients. Cancer-related inflammation and tumor-induced immune suppression are often associated with expansion of myeloid subsets including MDSCs. In parallel, the granulocytes are often activated in cancer patients with a concomitant increase in H<sub>2</sub>O<sub>2</sub> production [278]. G-CSF is often secreted by tumor cells [26, 174, 189, 190, 194, 198, 276, 352–359] and is preferentially observed in dedifferentiated or poorly differentiated tumors [360, 361, 354, 190]. Elevated G-CSF blood concentrations in cancer patients have been associated with poor clinical outcome [174, 194, 198, 276, 352, 355, 356].

Recently, a high myeloperoxidase (MPO)<sup>+</sup> CD15<sup>+</sup> cell infiltration representing neutrophils in colorectal carcinoma was found to be an independent favorable prognostic factor [362, 363]. Also, the increased presence of CD16<sup>+</sup> (FcγRIII) myeloid cells of the monocyte/macrophage lineage that were also positive for CD45, CD33, CD11b, and CD11c, but not CD64 or HLA-DR, was found to be associated with improved survival in patients with colorectal carcinoma [364]. This is in contrast to the general concept that high amounts of intratumoral myeloid cells promote tumor progression and hence correlate with poor disease outcome. In particular, colorectal carcinoma infiltration by CD66b<sup>+</sup> granulocytes was proposed as a marker of adverse prognosis [365].

Already in 1970, Riesco [366] observed a positive correlation between cancer curability and the total number of lymphocytes, while a negative correlation was found with the total number of peripheral neutrophils (segmented and nonsegmented) when 589 cases of different cancer types were investigated. From then on, a large number of clinical studies have been performed to understand the neutrophil-cancer relationship. Clinical data have often related elevated circulating neutrophil counts or elevated neutrophil-to-lymphocyte ratios (NLRs) as a predictive parameter for poor outcome and formation of distant metastasis in patients with epithelial malignancies [367], including lung [368, 170], gastric [159, 163, 369], renal cell carcinoma [370], ovarian [141], hepatic [143, 148, 371], pancreatic [138], colon cancer [145, 372], and colorectal carcinoma [152] (Table 3). However, Caruso et al. [200] observed that when analyzing the amount of tumor-infiltrated neutrophils in advanced gastric carcinoma, in female but not male, patients with higher TANs had a favorable prognosis.

Colorectal carcinoma sections were characterized by tumor-infiltrating granulocytes (TIGs) and tumor-associated macrophages (TAMs) and abnormal levels of the cytokines IL-1β, IL-6, IL-8, TNFα, G-CSF and M-CSF [178]. G-CSF was associated with a deeper tumor invasion and a more advanced tumor stage [178]. The granulocyte/lymphocyte ratio was associated with abnormal levels of G-CSF (more than 50 % of the cancer patients) and TIGs were a risk factor for lymph node metastasis in colorectal carcinoma [178]. Similarly, in gastric carcinoma, TIGs were associated with tumor stage and shorter survival time [373].

Studies on neutrophils from patients with head and neck squamous cell carcinoma (HNSCC) showed that these neutrophils differ from their counterparts in healthy donors [374]. The neutrophils from HNSCC showed lower inducible production of ROS, reduced spontaneous apoptosis and increased number of immature neutrophils [374]. The serum concentration of neutrophil related cytokines was higher in HNSCC patients [374]. HNSCC tissue exhibited considerable infiltration by neutrophils, and strong infiltration was associated with poorer survival in advanced diseases [157]. Neutrophil count, neutrophil-to-lymphocyte ratio and serum concentrations of CXCL8 (IL-8), CCL4 (MIP-1β) and CCL5 (RANTES) were significantly higher in the peripheral blood of HNSCC patients than in controls [157]. In vitro, HNSCC-conditioned medium inhibited apoptosis of neutrophils, increased neutrophil chemokinesis and chemotaxis and induced the release of lactoferrin, MMP-9 and CCL4 [157]. Further studies showed that HNSCC activates the p38-MAPK pathway in neutrophils, and stimulates the release of CCL4, CXCL8, and MMP-9 by neutrophils in a CREB-dependent manner [375]. The secretion of CCL4 and MMP-9 by neutrophils was stimulated by macrophage migration inhibitory factor (MIF) produced by the HNSCC cells [376]. MIF was also shown to be produced by other cancer cell types, including breast cancer, esophageal squamous cell carcinoma and hepatocellular carcinoma, where it contributes to angiogenesis [377–380]. Breast cancer patients with positive MIF expression in tumor tissues showed a significantly worse disease-free survival compared with MIF negative patients [377]. Thus, there seem to be a crosstalk between the tumor cells and neutrophils that may affect tumor growth and modify anti-tumor immune responses.

In human hepatocellular carcinoma, TANs accumulate in the peritumoral stroma due to IL-17-dependent release of epithelial cell derived chemokines such as CXCL1/GROα, CXCL2/MIP-2, CXCL3/GROγ and CXCL8/IL-8 [57]. High infiltration of peritumoral neutrophils correlated with tumor progression and predicted poor survival [57]. Neutrophils are the major source of MMP-9 within the hepatocellular carcinoma tissue, which was shown to stimulate the pro-angiogenic activity in hepatoma cells [57]. Selective depletion of neutrophils inhibited tumor angiogenesis and growth in vivo [57]. Also, CXCL5 has been shown to be

**Table 3** The correlation between neutrophil blood count and cancer prognosis

Cancer cell type	Effect on prognosis	Reference
Advanced pancreatic adenocarcinoma	Elevated pretreatment NLR>5 was a predictor of shorter survival.	[138]
Recurrent localized cervical cancer	Tumor-associated neutrophil count is an independent factor for short recurrence-free survival	[139]
Epithelial ovarian cancer	Pre-operative NLR in ovarian cancer subjects (mean 6.02) was significantly higher than in benign ovarian tumor subjects (mean 2.57) and healthy controls (mean 1.98). Elevated NLR may predict an adverse outcome.	[141]
Breast cancer	Patients with an NLR>2.5 showed lower disease-specific survival rate than those with an NLR<2.5. This correlation was especially seen in the luminal A subtype.	[423]
Breast cancer	Patients with an NLR>3.3 had a higher mortality rate than those with an NLR<1.8. The high NLR was associated with more advanced stages of cancer.	[424]
Hepatocellular carcinoma (HCC)	The presence of intratumoral CD66 <sup>+</sup> neutrophils was a poor prognostic factor for HCC after resection	[387]
Hepatocellular carcinoma (HCC)	Pre-operative NLR>5 was an adverse predictor of disease-free and overall survival.	[148]
Hepatocellular carcinoma (HCC)	Patients with an NLR>2.5 showed higher risk for tumor recurrence in HCC patients undergoing liver transplantation.	[371]
Hepatocellular carcinoma (HCC)	High baseline NLR was associated with higher tumor recurrence and worse overall survival in patients after radiofrequency ablation.	[140]
Hepatocellular carcinoma (HCC)	Elevated C-reactive protein together with an NLR>2.3 predicted shorter overall survival.	[425]
Small hepatocellular carcinoma	Elevated post-operative NLR change was a worse prognostic factor.	[143]
Resectable non-small cell lung cancer	An increase in CD66b <sup>+</sup> cells was associated with a high cumulative incidence of relapse (CIR) and worse overall survival.	[204]
	An increase in intratumoral neutrophil to CD8 <sup>+</sup> lymphocyte ratio (iNTR) was associated with high CIR and poor overall survival.	
Advanced non-small-cell lung cancer	A neutrophil count above 4,500/mm <sup>3</sup> was associated with shorter overall and progression-free survival.	[368]
Advanced non-small-cell lung cancer	Patients with normal leukocyte count had a longer median overall survival than those with elevated WBC count (>10×10 <sup>3</sup> /μl).	[426]
Advanced non-small-cell lung cancer	Increased preoperative NLR had a poorer prognosis.	[154]
Bronchioloalveolar carcinoma	The risk of death was increased in patients with a neutrophil percentage of >39 % in the bronchoalveolar lavage fluid.	[170]
Gastric carcinoma	Higher NLR was associated with lymph node metastasis, higher tumor stage, tumor progression and reduced 5-year survival.	[163]
Gastric cancer	Patients with an NLR>2.5 had a poorer prognosis.	[159]
Gastric cancer	NLR was influenced by tumor size. High NLR seems to be a marker for tumor recurrence.	[369]
Advanced gastric carcinoma	Female, but not male, patients with a moderate to extensive amount of tumor-infiltrating neutrophils had a 39 % reduction in their risk of mortality.	[200]
Gastric cancer stage III-IV	Progression-free survival and overall survival was worse for patients with high NLR.	[151]
Colon cancer, stage IIA	Patients with preoperative NLR>4 had a shorter recurrent-free survival.	[145]
Colon carcinoma, stage II	Patients with elevated NLR had a worse overall survival and worse disease-free survival.	[372]
Colorectal cancer	Preoperative NLR>5 correlated with poor pre-operative prognosis.	[158]
Colorectal cancer	The NLR was higher in advanced stages of cancer. The ability to produce ROS at the terminal stage was 33 % lower than in the control group.	[427]
Colorectal cancer	Patients with an NLR>2.5 had a poorer prognosis than those with an NLR<2.5.	[428]
Colorectal cancer	Patients with an NLR>5 showed a worse overall survival.	[429]
Metastatic colorectal cancer	A high NLR was associated with poor overall survival.	[430]
Colorectal liver metastases	The 5-year survival for patients undergoing resection with an NLR>5 was worse than those with normal NLR.	[150]
Non-metastatic renal cell carcinoma	Patients with a pre-operative and post-operative NLR>2.7 had a shorter recurrence-free survival rate.	[153, 431]
Localized renal cell carcinoma	The presence of intratumoral neutrophils was associated with short recurrence-free survival.	[390]
Metastatic renal cell carcinoma	Patients with elevated neutrophil count (>6,500 cells/μl) showed a shorter overall survival.	[432]
Metastatic renal cell carcinoma	Patients with a high blood neutrophil count (>6,000), intratumoral neutrophils and low intratumoral CD57 <sup>+</sup> NK cells are independent poor prognostic immunological factors.	[370]
Bladder cancer	There was no significant difference between patients with an NLR below or above 2.5 in terms of overall survival.	[144]
Bladder cancer	NLR was an independent prognostic factor in patients treated with radical cystectomy.	[149]
Urothelial carcinoma	An NLR>2.5 was found to be a predictor of invasiveness.	[433]

**Table 3** (continued)

Cancer cell type	Effect on prognosis	Reference
Melanoma, stage IV	A high pretreatment neutrophil count (>7,500 cells/ $\mu$ l) predicted shorter overall survival in patients receiving IL-2-based immunotherapy.	[156]
Oral squamous cell carcinoma	Patients with a pretreatment NLR>1.9 showed poorer disease-specific survival.	[434]
Esophageal cancer	Elevated NLR>5 is associated with poor outcomes.	[435]
Esophageal cancer	Patients with a pre-therapeutic NLR>2.2 showed poorer response to chemotherapy.	[436]
Esophageal cancer	No correlation between NLR and overall survival.	[437]
Esophageal cancer	High NLR>5 was associated with poorer overall survival.	[438]
Stomach cancer	Patients with a NLR>7.7 at post-operative day 3 showed a 4.2 times higher cancer recurrence after gastrectomy.	[439]
Metastatic Mesothelioma	An NLR>5 showed poorer overall survival in patients undergoing systemic therapy.	[440]
Metastatic Mesothelioma	An NLR>3 showed poorer prognosis after extrapleural pneumonectomy.	[441]
Metastatic Mesothelioma	A high NLR correlated with sustained neoangiogenesis and increased proliferative index.	[442]
Bone metastasis	A high preoperative N/L ratio was associated with poor prognosis after bone metastasis in the surgery group.	[443]

Evidence from clinical studies correlating neutrophil counts and Neutrophil to Lymphocyte Ratio (NLR) with cancer prognosis

overexpressed in hepatocellular carcinoma from patients with recurrent disease as well as in highly metastatic hepatocellular carcinoma cell lines [381]. The upregulation of CXCL5 in hepatocellular carcinoma cells correlated with the promotion of tumor growth, lung metastasis and intratumoral neutrophil infiltration [381].

Is it possible that the elevated neutrophil counts in cancer patients are actually the consequence of a more advanced stage, and therefore predict an apparent poorer prognosis? This was acknowledged by Liu et al. [178], who showed that the NLR was significantly associated with a more advanced colorectal tumor stage. Also Ietomi [382] observed that the NLR was higher in Stage IV stomach cancer patients than those having stage I-III stomach cancer. The NLR dropped after surgery, but upon relapse it rose again [382]. Similarly, Sarraf et al. [154] observed higher NLR in more advanced non-small cell lung cancer patients and Fossatti et al. [207] observed a marked and significant correlation between tumor grade of glioma patients and the extent of neutrophil infiltration as determined by the number of CD15<sup>+</sup>MPO<sup>+</sup> cells [207]. In low grade tumors 40–50 % show neutrophil infiltration, while in glioblastoma multiforme over 85 % of the samples show neutrophil infiltration [207]. These observations suggest that while neutrophil infiltration is associated with higher-grade tumors this does not mean that poor prognosis is due to increased neutrophil infiltration, rather the tumor is responsible for the elevated neutrophil number.

### Concluding Remarks - The Dialogue Between Cancer Cells and Neutrophils

There is a continuous interaction between cancer cells and neutrophils in the tumor microenvironment. Neutrophils are

attracted to the primary tumor by tumor-secreted chemokines, which can also induce anti-microbial as well as anti-tumor neutrophil activities. The activation of neutrophils under certain circumstances has been shown to yield prolonged life span in in vivo cancer models. These so-called “Tumor-Associated Neutrophils” (TANs) show enhanced NADPH oxidase activity which leads to the production of reactive oxygen species, especially hydrogen peroxide, that are cytotoxic to tumor cells [32]. After causing tumor cell apoptosis, these cells are then engulfed by neutrophils [383], and processed for antigen-presentation to mount an adaptive CD8<sup>+</sup> T cytotoxic anti-tumor response [15]. Neutrophils may work together with monocytes to transport tumor cell antigens to secondary lymphoid tissues, where naïve T cells are stimulated [383]. T cells further increase the activity of neutrophils through secretion of IFN $\gamma$ . The increase in oxygen radicals have additional effects, such as suppression of T cell responses [278, 384], activation of TGF $\beta$  [81], and induction of GMP-140 (P-Selectin) expression on the surface of endothelial cells, leading to enhanced neutrophil adherence and activity [385]. The activation of TGF $\beta$  suppresses excessive neutrophil function, and may polarize them into N2 neutrophils [13].

Conversely, certain tumor-stimulated neutrophils secrete ECM remodeling enzymes and pro-angiogenic factors that promote, directly or indirectly, the growth of the tumor as well as their detachment and dissemination. When the immunosuppressed environment is induced, there is a positive feedback mechanism to maintain the immunosuppressed condition. Unlike the tumor microenvironment, where immunosuppression prevails, neutrophils in the immunopermissive pre-metastatic organs may show characteristics of anti-tumorigenic cells, providing anti-metastatic protection by eliminating metastatic cells [32]. In light of this, we propose that while neutrophils are largely viewed as a homogenous

cell population, their function in cancer is dictated in a context-dependent fashion, which may seem conflicting. In depth exploration of how neutrophils interpret the orchestra of signals in the different cancer niches will enhance our understanding of how different neutrophil subsets function in the context of cancer.

**Acknowledgments** We thank Dr. Seth Salpeter for critical reading of this review. Research in the authors' laboratory is supported by the I-CORE Program of The Israel Science Foundation (Grant No. 41/11), The Abisch-Frenkel Foundation and The Rosetrees Trust (R.V.S and Z.G).

**Conflict of interest** None declared.

## References

- Pietras K, Ostman A (2010) Hallmarks of cancer: interactions with the tumor stroma. *Exp Cell Res* 316(8):1324–1331. doi:10.1016/j.yexcr.2010.02.045
- Fridman WH, Pages F, Sautes-Fridman C, Galon J (2012) The immune contexture in human tumours: impact on clinical outcome. *Nat Rev Cancer* 12(4):298–306. doi:10.1038/nrc3245
- Wels J, Kaplan RN, Rafii S, Lyden D (2008) Migratory neighbors and distant invaders: tumor-associated niche cells. *Genes Dev* 22(5):559–574. doi:10.1101/gad.1636908
- Queen MM, Ryan RE, Holzer RG, Keller-Peck CR, Jorcyk CL (2005) Breast cancer cells stimulate neutrophils to produce oncostatin M: potential implications for tumor progression. *Cancer Res* 65(19):8896–8904. doi:10.1158/0008-5472.CAN-05-1734
- Coussens LM, Werb Z (2002) Inflammation and cancer. *Nature* 420(6917):860–867. doi:10.1038/nature01322
- Galdiero MR, Bonavita E, Barajon I, Garlanda C, Mantovani A, Jaillon S (2013) Tumor associated macrophages and neutrophils in cancer. *Immunobiology*. doi:10.1016/j.imbio.2013.06.003
- Souto JC, Vila L, Bru A (2011) Polymorphonuclear neutrophils and cancer: intense and sustained neutrophilia as a treatment against solid tumors. *Med Res Rev* 31(3):311–363. doi:10.1002/med.20185
- Tazzyman S, Lewis CE, Murdoch C (2009) Neutrophils: key mediators of tumour angiogenesis. *Int J Exp Pathol* 90(3):222–231. doi:10.1111/j.1365-2613.2009.00641.x
- Gregory AD, Houghton AM (2011) Tumor-associated neutrophils: new targets for cancer therapy. *Cancer Res* 71(7):2411–2416. doi:10.1158/0008-5472.CAN-10-2583
- Fridlender ZG, Albelda SM (2012) Tumor-associated neutrophils: friend or foe? *Carcinogenesis* 33(5):949–955. doi:10.1093/carcin/bgs123
- Brandau S, Dumitru CA, Lang S (2013) Protumor and antitumor functions of neutrophil granulocytes. *Semin Immunopathol* 35(2):163–176. doi:10.1007/s00281-012-0344-6
- Fioretti F, Fradelizi D, Stoppacciaro A, Ramponi S, Ruco L, Minty A, Sozzani S, Garlanda C, Vecchi A, Mantovani A (1998) Reduced tumorigenicity and augmented leukocyte infiltration after monocyte chemotactic protein-3 (MCP-3) gene transfer: perivascular accumulation of dendritic cells in peritumoral tissue and neutrophil recruitment within the tumor. *J Immunol* 161(1):342–346
- Fridlender ZG, Sun J, Kim S, Kapoor V, Cheng G, Ling L, Worthen GS, Albelda SM (2009) Polarization of tumor-associated neutrophil phenotype by TGF-beta: "N1" versus "N2" TAN. *Cancer Cell* 16(3):183–194. doi:10.1016/j.ccr.2009.06.017
- Stoppacciaro A, Melani C, Parenza M, Mastracchio A, Bassi C, Baroni C, Parmiani G, Colombo MP (1993) Regression of an established tumor genetically modified to release granulocyte colony-stimulating factor requires granulocyte-T cell cooperation and T cell-produced interferon gamma. *J Exp Med* 178(1):151–161
- Mantovani A, Cassatella MA, Costantini C, Jaillon S (2011) Neutrophils in the activation and regulation of innate and adaptive immunity. *Nat Rev Immunol* 11(8):519–531. doi:10.1038/nri3024
- Cavallo F, Giovarelli M, Gulino A, Vacca A, Stoppacciaro A, Modesti A, Forni G (1992) Role of neutrophils and CD4+ T lymphocytes in the primary and memory response to nonimmunogenic murine mammary adenocarcinoma made immunogenic by IL-2 gene. *J Immunol* 149(11):3627–3635
- Suttman H, Riemensberger J, Bentien G, Schmaltz D, Stockle M, Jocham D, Bohle A, Brandau S (2006) Neutrophil granulocytes are required for effective Bacillus Calmette-Guerin immunotherapy of bladder cancer and orchestrate local immune responses. *Cancer Res* 66(16):8250–8257. doi:10.1158/0008-5472.CAN-06-1416
- Kousis PC, Henderson BW, Maier PG, Gollnick SO (2007) Photodynamic therapy enhancement of antitumor immunity is regulated by neutrophils. *Cancer Res* 67(21):10501–10510. doi:10.1158/0008-5472.CAN-07-1778
- Midorikawa Y, Yamashita T, Sendo F (1990) Modulation of the immune response to transplanted tumors in rats by selective depletion of neutrophils in vivo using a monoclonal antibody: abrogation of specific transplantation resistance to chemical carcinogen-induced syngeneic tumors by selective depletion of neutrophils in vivo. *Cancer Res* 50(19):6243–6247
- Ben-Neriah Y, Karin M (2011) Inflammation meets cancer, with NF-kappaB as the matchmaker. *Nat Immunol* 12(8):715–723. doi:10.1038/ni.2060
- Schreiber RD, Old LJ, Smyth MJ (2011) Cancer immunoeediting: integrating immunity's roles in cancer suppression and promotion. *Science* 331(6024):1565–1570. doi:10.1126/science.1203486
- Piccard H, Muschel RJ, Opendakker G (2012) On the dual roles and polarized phenotypes of neutrophils in tumor development and progression. *Crit Rev Oncol Hematol* 82(3):296–309. doi:10.1016/j.critrevonc.2011.06.004
- Chen Z, Malhotra PS, Thomas GR, Ondrey FG, Duffey DC, Smith CW, Enamorado I, Yeh NT, Kroog GS, Rudy S, McCullagh L, Mousa S, Quezado M, Herscher LL, Van Waes C (1999) Expression of proinflammatory and proangiogenic cytokines in patients with head and neck cancer. *Clin Cancer Res* 5(6):1369–1379
- Chen Z, Colon I, Ortiz N, Callister M, Dong G, Pegram MY, Arosarena O, Strome S, Nicholson JC, Van Waes C (1998) Effects of interleukin-1alpha, interleukin-1 receptor antagonist, and neutralizing antibody on proinflammatory cytokine expression by human squamous cell carcinoma lines. *Cancer Res* 58(16):3668–3676
- Arii K, Tanimura H, Iwahashi M, Tsunoda T, Tani M, Noguchi K, Mizobata S, Hotta T, Nakamori M, Yamaue H (2000) Neutrophil functions and cytokine production in patients with gastric cancer. *Hepatogastroenterology* 47(31):291–297
- Kowanetz M, Wu X, Lee J, Tan M, Hagenbeek T, Qu X, Yu L, Ross J, Korsisaari N, Cao T, Bou-Reslan H, Kallop D, Weimer R, Ludlam MJ, Kaminker JS, Modrusan Z, van Bruggen N, Peale FV, Carano R, Meng YG, Ferrara N (2010) Granulocyte-colony stimulating factor promotes lung metastasis through mobilization of Ly6G+ Ly6C+ granulocytes. *Proc Natl Acad Sci U S A* 107(50):21248–21255. doi:10.1073/pnas.1015855107
- Sawanobori Y, Ueha S, Kurachi M, Shimaoka T, Talmadge JE, Abe J, Shono Y, Kitabatake M, Kakimi K, Mukaida N, Matsushima K (2008) Chemokine-mediated rapid turnover of myeloid-derived suppressor cells in tumor-bearing mice. *Blood* 111(12):5457–5466. doi:10.1182/blood-2008-01-136895



28. Fridlender ZG, Buchlis G, Kapoor V, Cheng G, Sun J, Singhal S, Crisanti MC, Wang LC, Heitjan D, Snyder LA, Albelda SM (2010) CCL2 blockade augments cancer immunotherapy. *Cancer Res* 70(1):109–118. doi:10.1158/0008-5472.CAN-09-2326
29. Ninck S, Reisser C, Dyckhoff G, Helmke B, Bauer H, Herold-Mende C (2003) Expression profiles of angiogenic growth factors in squamous cell carcinomas of the head and neck. *Int J Cancer* 106(1):34–44. doi:10.1002/ijc.11188
30. Lin WW, Karin M (2007) A cytokine-mediated link between innate immunity, inflammation, and cancer. *J Clin Invest* 117(5):1175–1183. doi:10.1172/JCI15337
31. Revoltella RP, Menicagli M, Campani D (2012) Granulocyte-macrophage colony-stimulating factor as an autocrine survival-growth factor in human gliomas. *Cytokine* 57(3):347–359. doi:10.1016/j.cyto.2011.11.016
32. Granot Z, Henke E, Comen EA, King TA, Norton L, Benezra R (2011) Tumor entrained neutrophils inhibit seeding in the premetastatic lung. *Cancer Cell* 20(3):300–314. doi:10.1016/j.ccr.2011.08.012
33. Forte G, Sorrentino R, Montinaro A, Luciano A, Adcock IM, Maiolino P, Arra C, Cicala C, Pinto A, Morello S (2012) Inhibition of CD73 improves B cell-mediated anti-tumor immunity in a mouse model of melanoma. *J Immunol* 189(5):2226–2233. doi:10.4049/jimmunol.1200744
34. Kandasamy M, Bay BH, Lee YK, Mahendran R (2011) Lactobacilli secreting a tumor antigen and IL15 activates neutrophils and dendritic cells and generates cytotoxic T lymphocytes against cancer cells. *Cell Immunol* 271(1):89–96. doi:10.1016/j.cellimm.2011.06.004
35. Schwandt A, Garcia JA, Elson P, Wyckhouse J, Finke JH, Ireland J, Triozzi P, Zhou M, Dreicer R, Rini BI (2011) Clinical and immunomodulatory effects of celecoxib plus interferon-alpha in metastatic renal cell carcinoma patients with COX-2 tumor immunostaining. *J Clin Immunol* 31(4):690–698. doi:10.1007/s10875-011-9530-x
36. Yamamoto M, Kamigaki T, Yamashita K, Hori Y, Hasegawa H, Kuroda D, Moriyama H, Nagata M, Ku Y, Kuroda Y (2009) Enhancement of anti-tumor immunity by high levels of Th1 and Th17 with a combination of dendritic cell fusion hybrids and regulatory T cell depletion in pancreatic cancer. *Oncol Rep* 22(2):337–343
37. de Visser KE, Coussens LM (2006) The inflammatory tumor microenvironment and its impact on cancer development. *Contrib Microbiol* 13:118–137. doi:10.1159/000092969
38. Kenny PA, Bissell MJ (2003) Tumor reversion: correction of malignant behavior by microenvironmental cues. *Int J Cancer* 107(5):688–695. doi:10.1002/ijc.11491
39. Noguera R, Nieto OA, Tadeo I, Farinas F, Alvaro T (2012) Extracellular matrix, biotensegrity and tumor microenvironment. An update and overview. *Histol Histopathol* 27(6):693–705
40. Mocellin S, Rossi CR, Pilati P, Nitti D (2005) Tumor necrosis factor, cancer and anticancer therapy. *Cytokine Growth Factor Rev* 16(1):35–53. doi:10.1016/j.cytogfr.2004.11.001
41. Moore RJ, Owens DM, Stamp G, Arnott C, Burke F, East N, Holdsworth H, Turner L, Rollins B, Pasparakis M, Kollias G, Balkwill F (1999) Mice deficient in tumor necrosis factor-alpha are resistant to skin carcinogenesis. *Nat Med* 5(7):828–831. doi:10.1038/10552
42. Gu FM, Li QL, Gao Q, Jiang JH, Zhu K, Huang XY, Pan JF, Yan J, Hu JH, Wang Z, Dai Z, Fan J, Zhou J (2011) IL-17 induces AKT-dependent IL-6/JAK2/STAT3 activation and tumor progression in hepatocellular carcinoma. *Mol Cancer* 10:150. doi:10.1186/1476-4598-10-150
43. Luo JL, Maeda S, Hsu LC, Yagita H, Karin M (2004) Inhibition of NF-kappaB in cancer cells converts inflammation-induced tumor growth mediated by TNFalpha to TRAIL-mediated tumor regression. *Cancer Cell* 6(3):297–305. doi:10.1016/j.ccr.2004.08.012
44. Sainson RC, Johnston DA, Chu HC, Holderfield MT, Nakatsu MN, Crampton SP, Davis J, Conn E, Hughes CC (2008) TNF primes endothelial cells for angiogenic sprouting by inducing a tip cell phenotype. *Blood* 111(10):4997–5007. doi:10.1182/blood-2007-08-108597
45. Jing Y, Ma N, Fan T, Wang C, Bu X, Jiang G, Li R, Gao L, Li D, Wu M, Wei L (2011) Tumor necrosis factor-alpha promotes tumor growth by inducing vascular endothelial growth factor. *Cancer Invest* 29(7):485–493. doi:10.3109/07357907.2011.597812
46. Tzeng HE, Tsai CH, Chang ZL, Su CM, Wang SW, Hwang WL, Tang CH (2013) Interleukin-6 induces vascular endothelial growth factor expression and promotes angiogenesis through apoptosis signal-regulating kinase 1 in human osteosarcoma. *Biochem Pharmacol* 85(4):531–540. doi:10.1016/j.bcp.2012.11.021
47. Whalen GF (1990) Solid tumours and wounds: transformed cells misunderstood as injured tissue? *Lancet* 336(8729):1489–1492
48. Sica A, Mantovani A (2012) Macrophage plasticity and polarization: in vivo veritas. *J Clin Invest* 122(3):787–795. doi:10.1172/JCI59643
49. Mantovani A, Sozzani S, Locati M, Allavena P, Sica A (2002) Macrophage polarization: tumor-associated macrophages as a paradigm for polarized M2 mononuclear phagocytes. *Trends Immunol* 23(11):549–555
50. Badylak SF, Valentin JE, Ravindra AK, McCabe GP, Stewart-Akers AM (2008) Macrophage phenotype as a determinant of biologic scaffold remodeling. *Tissue Eng Part A* 14(11):1835–1842. doi:10.1089/ten.tea.2007.0264
51. Mantovani A, Locati M (2013) Tumor-associated macrophages as a paradigm of macrophage plasticity, diversity, and polarization: lessons and open questions. *Arterioscler Thromb Vasc Biol* 33(7):1478–1483. doi:10.1161/ATVBAHA.113.300168
52. Hargadon KM (2013) Tumor-altered dendritic cell function: implications for anti-tumor immunity. *Front Immunol* 4:192. doi:10.3389/fimmu.2013.00192
53. Biswas SK, Gangi L, Paul S, Schioppa T, Saccani A, Sironi M, Bottazzi B, Doni A, Vincenzo B, Pasqualini F, Vago L, Nebuloni M, Mantovani A, Sica A (2006) A distinct and unique transcriptional program expressed by tumor-associated macrophages (defective NF-kappaB and enhanced IRF-3/STAT1 activation). *Blood* 107(5):2112–2122. doi:10.1182/blood-2005-01-0428
54. Kuang DM, Peng C, Zhao Q, Wu Y, Zhu LY, Wang J, Yin XY, Li L, Zheng L (2010) Tumor-activated monocytes promote expansion of IL-17-producing CD8+ T cells in hepatocellular carcinoma patients. *J Immunol* 185(3):1544–1549. doi:10.4049/jimmunol.0904094
55. Wilke CM, Kryczek I, Wei S, Zhao E, Wu K, Wang G, Zou W (2011) Th17 cells in cancer: help or hindrance? *Carcinogenesis* 32(5):643–649. doi:10.1093/carcin/bgr019
56. Laan M, Cui ZH, Hoshino H, Lotvall J, Sjostrand M, Gruenert DC, Skoogh BE, Linden A (1999) Neutrophil recruitment by human IL-17 via C-X-C chemokine release in the airways. *J Immunol* 162(4):2347–2352
57. Kuang DM, Zhao Q, Wu Y, Peng C, Wang J, Xu Z, Yin XY, Zheng L (2011) Peritumoral neutrophils link inflammatory response to disease progression by fostering angiogenesis in hepatocellular carcinoma. *J Hepatol* 54(5):948–955. doi:10.1016/j.jhep.2010.08.041
58. Ley K, Smith E, Stark MA (2006) IL-17A-producing neutrophil-regulatory Tn lymphocytes. *Immunol Res* 34(3):229–242. doi:10.1385/IR:34:3:229
59. Stark MA, Huo Y, Burcin TL, Morris MA, Olson TS, Ley K (2005) Phagocytosis of apoptotic neutrophils regulates granulopoiesis via IL-23 and IL-17. *Immunity* 22(3):285–294. doi:10.1016/j.immuni.2005.01.011
60. Houghton AM (2010) The paradox of tumor-associated neutrophils: fueling tumor growth with cytotoxic substances. *Cell Cycle* 9(9):1732–1737

61. Scapini P, Lapinet-Vera JA, Gasperini S, Calzetti F, Bazzoni F, Cassatella MA (2000) The neutrophil as a cellular source of chemokines. *Immunol Rev* 177:195–203
62. Pelletier M, Maggi L, Micheletti A, Lazzeri E, Tamassia N, Costantini C, Cosmi L, Lunardi C, Annunziato F, Romagnani S, Cassatella MA (2010) Evidence for a cross-talk between human neutrophils and Th17 cells. *Blood* 115(2):335–343. doi:10.1182/blood-2009-04-216085
63. Liang SC, Long AJ, Bennett F, Whitters MJ, Karim R, Collins M, Goldman SJ, Dunussi-Joannopoulos K, Williams CM, Wright JF, Fouser LA (2007) An IL-17F/A heterodimer protein is produced by mouse Th17 cells and induces airway neutrophil recruitment. *J Immunol* 179(11):7791–7799
64. Jablonska J, Leschner S, Westphal K, Lienenklaus S, Weiss S (2010) Neutrophils responsive to endogenous IFN-beta regulate tumor angiogenesis and growth in a mouse tumor model. *J Clin Invest* 120(4):1151–1164. doi:10.1172/JCI37223
65. Fridlender ZG, Sun J, Mishalian I, Singhal S, Cheng G, Kapoor V, Horng W, Fridlender G, Bayuh R, Worthen GS, Albelda SM (2012) Transcriptomic analysis comparing tumor-associated neutrophils with granulocytic myeloid-derived suppressor cells and normal neutrophils. *PLoS One* 7(2):e31524. doi:10.1371/journal.pone.0031524
66. Lopez-Lago MA, Posner S, Thodima VJ, Molina AM, Motzer RJ, Chaganti RS (2013) Neutrophil chemokines secreted by tumor cells mount a lung antimetastatic response during renal cell carcinoma progression. *Oncogene* 32(14):1752–1760. doi:10.1038/onc.2012.201
67. Swann JB, Smyth MJ (2007) Immune surveillance of tumors. *J Clin Invest* 117(5):1137–1146. doi:10.1172/JCI31405
68. Zhang Q, Yang X, Pins M, Javonovic B, Kuzel T, Kim SJ, Parijs LV, Greenberg NM, Liu V, Guo Y, Lee C (2005) Adoptive transfer of tumor-reactive transforming growth factor-beta-insensitive CD8+ T cells: eradication of autologous mouse prostate cancer. *Cancer Res* 65(5):1761–1769. doi:10.1158/0008-5472.CAN-04-3169
69. Gholamin M, Moaven O, Memar B, Farshchian M, Naseh H, Malekzadeh R, Sotoudeh M, Rajabi-Mashhadi MT, Forghani MN, Farrokhi F, Abbaszadegan MR (2009) Overexpression and interactions of interleukin-10, transforming growth factor beta, and vascular endothelial growth factor in esophageal squamous cell carcinoma. *World J Surg* 33(7):1439–1445. doi:10.1007/s00268-009-0070-y
70. Gabrilovich DI, Ishida T, Nadaf S, Ohm JE, Carbone DP (1999) Antibodies to vascular endothelial growth factor enhance the efficacy of cancer immunotherapy by improving endogenous dendritic cell function. *Clin Cancer Res* 5(10):2963–2970
71. Gerlini G, Tun-Kyi A, Dudli C, Burg G, Pimpinelli N, Nestle FO (2004) Metastatic melanoma secreted IL-10 down-regulates CD1 molecules on dendritic cells in metastatic tumor lesions. *Am J Pathol* 165(6):1853–1863. doi:10.1016/S0002-9440(10)63238-5
72. Jayaraman P, Parikh F, Lopez-Rivera E, Hailemichael Y, Clark A, Ma G, Cannan D, Ramacher M, Kato M, Overwijk WW, Chen SH, Umansky VY, Sikora AG (2012) Tumor-expressed inducible nitric oxide synthase controls induction of functional myeloid-derived suppressor cells through modulation of vascular endothelial growth factor release. *J Immunol* 188(11):5365–5376. doi:10.4049/jimmunol.1103553
73. Choudhari SK, Chaudhary M, Bagde S, Gadbail AR, Joshi V (2013) Nitric oxide and cancer: a review. *World J Surg Oncol* 11: 118. doi:10.1186/1477-7819-11-118
74. Greenhough A, Smartt HJ, Moore AE, Roberts HR, Williams AC, Paraskeva C, Kaidi A (2009) The COX-2/PGE2 pathway: key roles in the hallmarks of cancer and adaptation to the tumour microenvironment. *Carcinogenesis* 30(3):377–386. doi:10.1093/carcin/bgp014
75. Brown JR, DuBois RN (2005) COX-2: a molecular target for colorectal cancer prevention. *J Clin Oncol* 23(12):2840–2855. doi:10.1200/JCO.2005.09.051
76. McKallip R, Li R, Ladisch S (1999) Tumor gangliosides inhibit the tumor-specific immune response. *J Immunol* 163(7):3718–3726
77. Hossain DM, Mohanty S, Ray P, Das T, Sa G (2012) Tumor gangliosides and T cells: a deadly encounter. *Front Biosci (Schol Ed)* 4:502–519
78. Bierie B, Moses HL (2010) Transforming growth factor beta (TGF-beta) and inflammation in cancer. *Cytokine Growth Factor Rev* 21(1):49–59. doi:10.1016/j.cytogfr.2009.11.008
79. Wojtowicz-Praga S (2003) Reversal of tumor-induced immunosuppression by TGF-beta inhibitors. *Invest New Drugs* 21(1):21–32
80. Levy L, Hill CS (2006) Alterations in components of the TGF-beta superfamily signaling pathways in human cancer. *Cytokine Growth Factor Rev* 17(1–2):41–58. doi:10.1016/j.cytogfr.2005.09.009
81. Barcellos-Hoff MH, Dix TA (1996) Redox-mediated activation of latent transforming growth factor-beta 1. *Mol Endocrinol* 10(9): 1077–1083
82. Yang L, Huang J, Ren X, Gorska AE, Chytil A, Aakre M, Carbone DP, Matrisian LM, Richmond A, Lin PC, Moses HL (2008) Abrogation of TGF beta signaling in mammary carcinomas recruits Gr-1+CD11b+ myeloid cells that promote metastasis. *Cancer Cell* 13(1):23–35. doi:10.1016/j.ccr.2007.12.004
83. Zamarron BF, Chen W (2011) Dual roles of immune cells and their factors in cancer development and progression. *Int J Biol Sci* 7(5): 651–658
84. Lechner MG, Megiel C, Russell SM, Bingham B, Arger N, Woo T, Epstein AL (2011) Functional characterization of human CD33+ and Cd11b+ myeloid-derived suppressor cell subsets induced from peripheral blood mononuclear cells co-cultured with a diverse set of human tumor cell lines. *J Transl Med* 9:90. doi:10.1186/1479-5876-9-90
85. Shen L, Smith JM, Shen Z, Eriksson M, Sentman C, Wira CR (2007) Inhibition of human neutrophil degranulation by transforming growth factor-beta1. *Clin Exp Immunol* 149(1):155–161. doi:10.1111/j.1365-2249.2007.03376.x
86. Malipiero U, Koedel U, Pfister HW, Leveen P, Burki K, Reith W, Fontana A (2006) TGFbeta receptor II gene deletion in leucocytes prevents cerebral vasculitis in bacterial meningitis. *Brain* 129(Pt 9): 2404–2415. doi:10.1093/brain/awl192
87. Parekh T, Saxena B, Reibman J, Cronstein BN, Gold LI (1994) Neutrophil chemotaxis in response to TGF-beta isoforms (TGF-beta 1, TGF-beta 2, TGF-beta 3) is mediated by fibronectin. *J Immunol* 152(5):2456–2466
88. Fava RA, Olsen NJ, Postlethwaite AE, Broadley KN, Davidson JM, Nanney LB, Lucas C, Townes AS (1991) Transforming growth factor beta 1 (TGF-beta 1) induced neutrophil recruitment to synovial tissues: implications for TGF-beta-driven synovial inflammation and hyperplasia. *J Exp Med* 173(5):1121–1132
89. Reibman J, Meixler S, Lee TC, Gold LI, Cronstein BN, Haines KA, Kolasinski SL, Weissmann G (1991) Transforming growth factor beta 1, a potent chemoattractant for human neutrophils, bypasses classic signal-transduction pathways. *Proc Natl Acad Sci U S A* 88(15):6805–6809
90. Kim S, Buchlis G, Fridlender ZG, Sun J, Kapoor V, Cheng G, Haas A, Cheung HK, Zhang X, Corbely M, Kaiser LR, Ling L, Albelda SM (2008) Systemic blockade of transforming growth factor-beta signaling augments the efficacy of immunogene therapy. *Cancer Res* 68(24):10247–10256. doi:10.1158/0008-5472.CAN-08-1494
91. Smith WB, Noack L, Khew-Goodall Y, Isenmann S, Vadas MA, Gamble JR (1996) Transforming growth factor-beta 1 inhibits the production of IL-8 and the transmigration of neutrophils through activated endothelium. *J Immunol* 157(1):360–368
92. Bierie B, Stover DG, Abel TW, Chytil A, Gorska AE, Aakre M, Forrester E, Yang L, Wagner KU, Moses HL (2008) Transforming growth factor-beta regulates mammary carcinoma cell survival and interaction with the adjacent microenvironment. *Cancer Res* 68(6): 1809–1819. doi:10.1158/0008-5472.CAN-07-5597

93. Kalinski P (2012) Regulation of immune responses by prostaglandin E<sub>2</sub>. *J Immunol* 188(1):21–28. doi:10.4049/jimmunol.1101029
94. Wang MT, Honn KV, Nie D (2007) Cyclooxygenases, prostanoids, and tumor progression. *Cancer Metastasis Rev* 26(3–4):525–534. doi:10.1007/s10555-007-9096-5
95. Eisinger AL, Prescott SM, Jones DA, Stafforini DM (2007) The role of cyclooxygenase-2 and prostaglandins in colon cancer. *Prostaglandins Other Lipid Mediat* 82(1–4):147–154. doi:10.1016/j.prostaglandins.2006.05.026
96. Schrey MP, Patel KV (1995) Prostaglandin E<sub>2</sub> production and metabolism in human breast cancer cells and breast fibroblasts. Regulation by inflammatory mediators. *Br J Cancer* 72(6):1412–1419
97. Ristimaki A, Sivula A, Lundin J, Lundin M, Salminen T, Haglund C, Joensuu H, Isola J (2002) Prognostic significance of elevated cyclooxygenase-2 expression in breast cancer. *Cancer Res* 62(3):632–635
98. Zea AH, Rodriguez PC, Atkins MB, Hernandez C, Signoretti S, Zabaleta J, McDermott D, Quiceno D, Youmans A, O'Neill A, Mier J, Ochoa AC (2005) Arginase-producing myeloid suppressor cells in renal cell carcinoma patients: a mechanism of tumor evasion. *Cancer Res* 65(8):3044–3048. doi:10.1158/0008-5472.CAN-04-4505
99. Riedl K, Krysan K, Pold M, Dalwadi H, Heuze-Vourc'h N, Dohadwala M, Liu M, Cui X, Figlin R, Mao JT, Strieter R, Sharma S, Dubinett SM (2004) Multifaceted roles of cyclooxygenase-2 in lung cancer. *Drug Resist Updat* 7(3):169–184. doi:10.1016/j.drug.2004.04.003
100. Obermajer N, Wong JL, Edwards RP, Odunsi K, Moysich K, Kalinski P (2012) PGE<sub>2</sub>-driven induction and maintenance of cancer-associated myeloid-derived suppressor cells. *Immunol Investig* 41(6–7):635–657. doi:10.3109/08820139.2012.695417
101. Plescia OJ, Smith AH, Grinwich K (1975) Subversion of immune system by tumor cells and role of prostaglandins. *Proc Natl Acad Sci U S A* 72(5):1848–1851
102. Sionov RV, Gallily R (1990) Engulfment and intracellular killing of F9 teratocarcinoma cells by non-activated murine macrophages. *Int Immunol* 2(4):291–301
103. Mulligan JK, Rosenzweig SA, Young MR (2010) Tumor secretion of VEGF induces endothelial cells to suppress T cell functions through the production of PGE<sub>2</sub>. *J Immunother* 33(2):126–135. doi:10.1097/CJI.0b013e3181b91c9c
104. Khayrullina T, Yen JH, Jing H, Ganea D (2008) In vitro differentiation of dendritic cells in the presence of prostaglandin E<sub>2</sub> alters the IL-12/IL-23 balance and promotes differentiation of Th17 cells. *J Immunol* 181(1):721–735
105. Lemos HP, Grespan R, Vieira SM, Cunha TM, Verri WA Jr, Fernandes KS, Souto FO, McInnes IB, Ferreira SH, Liew FY, Cunha FQ (2009) Prostaglandin mediates IL-23/IL-17-induced neutrophil migration in inflammation by inhibiting IL-12 and IFN $\gamma$  production. *Proc Natl Acad Sci U S A* 106(14):5954–5959. doi:10.1073/pnas.0812782106
106. Karavitis J, Zhang M (2013) COX2 regulation of breast cancer bone metastasis. *Oncoimmunology* 2(3):e23129. doi:10.4161/onci.23129
107. Yu Y, Chadee K (1998) Prostaglandin E<sub>2</sub> stimulates IL-8 gene expression in human colonic epithelial cells by a posttranscriptional mechanism. *J Immunol* 161(7):3746–3752
108. Nakayama T, Mutsuga N, Yao L, Tosato G (2006) Prostaglandin E<sub>2</sub> promotes degranulation-independent release of MCP-1 from mast cells. *J Leukoc Biol* 79(1):95–104. doi:10.1189/jlb.0405226
109. Weller CL, Collington SJ, Hartnell A, Conroy DM, Kaise T, Barker JE, Wilson MS, Taylor GW, Jose PJ, Williams TJ (2007) Chemotactic action of prostaglandin E<sub>2</sub> on mouse mast cells acting via the PGE<sub>2</sub> receptor 3. *Proc Natl Acad Sci U S A* 104(28):11712–11717. doi:10.1073/pnas.0701700104
110. Aso H, Ito S, Mori A, Morioka M, Suganuma N, Kondo M, Imaizumi K, Hasegawa Y (2012) Prostaglandin E<sub>2</sub> enhances interleukin-8 production via EP<sub>4</sub> receptor in human pulmonary microvascular endothelial cells. *Am J Physiol Lung Cell Mol Physiol* 302(2):L266–L273. doi:10.1152/ajplung.00248.2011
111. He LK, Liu LH, Hahn E, Gamelli RL (2001) The expression of cyclooxygenase and the production of prostaglandin E<sub>2</sub> in neutrophils after burn injury and infection. *J Burn Care Rehabil* 22(1):58–64
112. Cockeran R, Steel HC, Mitchell TJ, Feldman C, Anderson R (2001) Pneumolysin potentiates production of prostaglandin E<sub>2</sub> and leukotriene B<sub>4</sub> by human neutrophils. *Infect Immun* 69(5):3494–3496. doi:10.1128/IAI.69.5.3494-3496.2001
113. Ottonello L, Gonella R, Dapino P, Sacchetti C, Dallegri F (1998) Prostaglandin E<sub>2</sub> inhibits apoptosis in human neutrophilic polymorphonuclear leukocytes: role of intracellular cyclic AMP levels. *Exp Hematol* 26(9):895–902
114. Sottile A, Venza M, Venza I, Teti D (1995) Prostaglandins affect the respiratory burst of human neutrophils. *Immunopharmacol Immunotoxicol* 17(2):311–321. doi:10.3109/08923979509019753
115. Ham EA, Soderman DD, Zanetti ME, Dougherty HW, McCauley E, Kuehl FA Jr (1983) Inhibition by prostaglandins of leukotriene B<sub>4</sub> release from activated neutrophils. *Proc Natl Acad Sci U S A* 80(14):4349–4353
116. Talpain E, Armstrong RA, Coleman RA, Vardey CJ (1995) Characterization of the PGE receptor subtype mediating inhibition of superoxide production in human neutrophils. *Br J Pharmacol* 114(7):1459–1465
117. Ottonello L, Morone MP, Dapino P, Dallegri F (1995) Cyclic AMP-elevating agents down-regulate the oxidative burst induced by granulocyte-macrophage colony-stimulating factor (GM-CSF) in adherent neutrophils. *Clin Exp Immunol* 101(3):502–506
118. Yu CL, Huang MH, Kung YY, Tsai CY, Tsai YY, Tsai ST, Huang DF, Sun KH, Han SH, Yu HS (1998) Interleukin-13 increases prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) production by normal human polymorphonuclear neutrophils by enhancing cyclooxygenase 2 (COX-2) gene expression. *Inflamm Res* 47(4):167–173
119. Stolina M, Sharma S, Lin Y, Dohadwala M, Gardner B, Luo J, Zhu L, Kronenberg M, Miller PW, Portanova J, Lee JC, Dubinett SM (2000) Specific inhibition of cyclooxygenase 2 restores antitumor reactivity by altering the balance of IL-10 and IL-12 synthesis. *J Immunol* 164(1):361–370
120. Chan AT, Cook NR (2012) Are we ready to recommend aspirin for cancer prevention? *Lancet* 379(9826):1569–1571. doi:10.1016/S0140-6736(11)61654-1
121. Dominguez-Luis M, Herrera-Garcia A, Arce-Franco M, Armas-Gonzalez E, Rodriguez-Pardo M, Lorenzo-Diaz F, Feria M, Cadenas S, Sanchez-Madrid F, Diaz-Gonzalez F (2013) Superoxide anion mediates the L-selectin down-regulation induced by non-steroidal anti-inflammatory drugs in human neutrophils. *Biochem Pharmacol* 85(2):245–256. doi:10.1016/j.bcp.2012.10.024
122. Baratelli F, Lee JM, Hazra S, Lin Y, Walser TC, Schaub D, Pak PS, Elashoff D, Reckamp K, Zhang L, Fishbein MC, Sharma S, Dubinett SM (2010) PGE<sub>2</sub> contributes to TGF- $\beta$  induced T regulatory cell function in human non-small cell lung cancer. *Am J Transl Res* 2(4):356–367
123. Godin-Ethier J, Hanafi LA, Piccirillo CA, Lapointe R (2011) Indoleamine 2,3-dioxygenase expression in human cancers: clinical and immunologic perspectives. *Clin Cancer Res* 17(22):6985–6991. doi:10.1158/1078-0432.CCR-11-1331
124. Katz JB, Muller AJ, Prendergast GC (2008) Indoleamine 2,3-dioxygenase in T-cell tolerance and tumoral immune escape. *Immunol Rev* 222:206–221. doi:10.1111/j.1600-065X.2008.00610.x
125. Movahedi K, Laoui D, Gysemans C, Baeten M, Stange G, Van den Bossche J, Mack M, Pipeleers D, In't Veld P, De Baetselier P, Van Ginderachter JA (2010) Different tumor microenvironments contain



- functionally distinct subsets of macrophages derived from Ly6C(high) monocytes. *Cancer Res* 70(14):5728–5739. doi:10.1158/0008-5472.CAN-09-4672
126. Yu J, Du W, Yan F, Wang Y, Li H, Cao S, Yu W, Shen C, Liu J, Ren X (2013) Myeloid-derived suppressor cells suppress antitumor immune responses through IDO expression and correlate with lymph node metastasis in patients with breast cancer. *J Immunol* 190(7):3783–3797. doi:10.4049/jimmunol.1201449
  127. Blache CA, Manuel ER, Kaltcheva TI, Wong AN, Ellenhorn JD, Blazar BR, Diamond DJ (2012) Systemic delivery of Salmonella typhimurium transformed with IDO shRNA enhances intratumoral vector colonization and suppresses tumor growth. *Cancer Res* 72(24):6447–6456. doi:10.1158/0008-5472.CAN-12-0193
  128. Whiteside TL, Jackson EK (2013) Adenosine and prostaglandin E2 production by human inducible regulatory T cells in health and disease. *Front Immunol* 4:212. doi:10.3389/fimmu.2013.00212
  129. Deaglio S, Dwyer KM, Gao W, Friedman D, Usheva A, Erat A, Chen JF, Enjoji K, Linden J, Oukka M, Kuchroo VK, Strom TB, Robson SC (2007) Adenosine generation catalyzed by CD39 and CD73 expressed on regulatory T cells mediates immune suppression. *J Exp Med* 204(6):1257–1265. doi:10.1084/jem.20062512
  130. Colgan SP, Eltzschig HK, Eckle T, Thompson LF (2006) Physiological roles for ecto-5'-nucleotidase (CD73). *Purinergic Signal* 2(2):351–360. doi:10.1007/s11302-005-5302-5
  131. Yamashita Y, Hooker SW, Jiang H, Laurent AB, Resta R, Khare K, Coe A, Kincade PW, Thompson LF (1998) CD73 expression and fyn-dependent signaling on murine lymphocytes. *Eur J Immunol* 28(10):2981–2990. doi:10.1002/(SICI)1521-4141(199810)28:10<2981::AID-IMMU2981>3.0.CO;2-D
  132. Zhang B (2010) CD73: a novel target for cancer immunotherapy. *Cancer Res* 70(16):6407–6411. doi:10.1158/0008-5472.CAN-10-1544
  133. Jin D, Fan J, Wang L, Thompson LF, Liu A, Daniel BJ, Shin T, Curiel TJ, Zhang B (2010) CD73 on tumor cells impairs antitumor T-cell responses: a novel mechanism of tumor-induced immune suppression. *Cancer Res* 70(6):2245–2255. doi:10.1158/0008-5472.CAN-09-3109
  134. Reutershan J, Vollmer I, Stark S, Wagner R, Ngamsri KC, Eltzschig HK (2009) Adenosine and inflammation: CD39 and CD73 are critical mediators in LPS-induced PMN trafficking into the lungs. *FASEB J* 23(2):473–482. doi:10.1096/fj.08-119701
  135. Stagg J, Divisekera U, Duret H, Sparwasser T, Teng MW, Darcy PK, Smyth MJ (2011) CD73-deficient mice have increased antitumor immunity and are resistant to experimental metastasis. *Cancer Res* 71(8):2892–2900. doi:10.1158/0008-5472.CAN-10-4246
  136. Stagg J, Divisekera U, McLaughlin N, Sharkey J, Pommey S, Denoyer D, Dwyer KM, Smyth MJ (2010) Anti-CD73 antibody therapy inhibits breast tumor growth and metastasis. *Proc Natl Acad Sci U S A* 107(4):1547–1552. doi:10.1073/pnas.0908801107
  137. Almand B, Clark JI, Nikitina E, van Beynen J, English NR, Knight SC, Carbone DP, Gabrilovich DI (2001) Increased production of immature myeloid cells in cancer patients: a mechanism of immunosuppression in cancer. *J Immunol* 166(1):678–689
  138. An X, Ding PR, Li YH, Wang FH, Shi YX, Wang ZQ, He YJ, Xu RH, Jiang WQ (2010) Elevated neutrophil to lymphocyte ratio predicts survival in advanced pancreatic cancer. *Biomarkers* 15(6):516–522. doi:10.3109/1354750X.2010.491557
  139. Carus A, Ladekarl M, Hager H, Nedergaard BS, Donskov F (2013) Tumour-associated CD66b+ neutrophil count is an independent prognostic factor for recurrence in localised cervical cancer. *Br J Cancer* 108(10):2116–2122. doi:10.1038/bjc.2013.167
  140. Chen TM, Lin CC, Huang PT, Wen CF (2012) Neutrophil-to-lymphocyte ratio associated with mortality in early hepatocellular carcinoma patients after radiofrequency ablation. *J Gastroenterol Hepatol* 27(3):553–561. doi:10.1111/j.1440-1746.2011.06910.x
  141. Cho H, Hur HW, Kim SW, Kim SH, Kim JH, Kim YT, Lee K (2009) Pre-treatment neutrophil to lymphocyte ratio is elevated in epithelial ovarian cancer and predicts survival after treatment. *Cancer Immunol Immunother* 58(1):15–23. doi:10.1007/s00262-008-0516-3
  142. Choi J, Suh B, Ahn YO, Kim TM, Lee JO, Lee SH, Heo DS (2012) CD15+/CD16low human granulocytes from terminal cancer patients: granulocytic myeloid-derived suppressor cells that have suppressive function. *Tumour Biol* 33(1):121–129. doi:10.1007/s13277-011-0254-6
  143. Dan J, Zhang Y, Peng Z, Huang J, Gao H, Xu L, Chen M (2013) Postoperative neutrophil-to-lymphocyte ratio change predicts survival of patients with small hepatocellular carcinoma undergoing radiofrequency ablation. *PLoS One* 8(3):e58184. doi:10.1371/journal.pone.0058184
  144. Demirtas A, Sabur V, Akinsal EC, Demirci D, Ekmekcioglu O, Gulmez I, Tatlisn A (2013) Can neutrophil-lymphocyte ratio and lymph node density be used as prognostic factors in patients undergoing radical cystectomy? *ScientificWorldJournal* 2013:703579. doi:10.1155/2013/703579
  145. Ding PR, An X, Zhang RX, Fang YJ, Li LR, Chen G, Wu XJ, Lu ZH, Lin JZ, Kong LH, Wan DS, Pan ZZ (2010) Elevated preoperative neutrophil to lymphocyte ratio predicts risk of recurrence following curative resection for stage IIA colon cancer. *Int J Colorectal Dis* 25(12):1427–1433. doi:10.1007/s00384-010-1052-0
  146. Dumitru CA, Lang S, Brandau S (2013) Modulation of neutrophil granulocytes in the tumor microenvironment: mechanisms and consequences for tumor progression. *Semin Cancer Biol* 23(3):141–148. doi:10.1016/j.semcancer.2013.02.005
  147. Eruslanov E, Neuberger M, Daurkin I, Perrin GQ, Algood C, Dahm P, Rosser C, Vieweg J, Gilbert SM, Kusmartsev S (2012) Circulating and tumor-infiltrating myeloid cell subsets in patients with bladder cancer. *Int J Cancer* 130(5):1109–1119. doi:10.1002/ijc.26123
  148. Gomez D, Farid S, Malik HZ, Young AL, Toogood GJ, Lodge JP, Prasad KR (2008) Preoperative neutrophil-to-lymphocyte ratio as a prognostic predictor after curative resection for hepatocellular carcinoma. *World J Surg* 32(8):1757–1762. doi:10.1007/s00268-008-9552-6
  149. Gondo T, Nakashima J, Ohno Y, Choichiro O, Horiguchi Y, Namiki K, Yoshioka K, Ohori M, Hatano T, Tachibana M (2012) Prognostic value of neutrophil-to-lymphocyte ratio and establishment of novel preoperative risk stratification model in bladder cancer patients treated with radical cystectomy. *Urology* 79(5):1085–1091. doi:10.1016/j.urology.2011.11.070
  150. Halazun KJ, Aldoori A, Malik HZ, Al-Mukhtar A, Prasad KR, Toogood GJ, Lodge JP (2008) Elevated preoperative neutrophil to lymphocyte ratio predicts survival following hepatic resection for colorectal liver metastases. *Eur J Surg Oncol* 34(1):55–60. doi:10.1016/j.ejso.2007.02.014
  151. Jin H, Zhang G, Liu X, Chen C, Yu H, Huang X, Zhang Q, Yu J (2013) Blood neutrophil-lymphocyte ratio predicts survival for stages III-IV gastric cancer treated with neoadjuvant chemotherapy. *World J Surg Oncol* 11:112. doi:10.1186/1477-7819-11-112
  152. Liu H, Liu G, Bao Q, Sun W, Bao H, Bi L, Wen W, Liu Y, Wang Z, Yin X, Bai Y, Hu X (2010) The baseline ratio of neutrophils to lymphocytes is associated with patient prognosis in rectal carcinoma. *J Gastrointest Cancer* 41(2):116–120. doi:10.1007/s12029-009-9125-4
  153. Ohno Y, Nakashima J, Ohori M, Gondo T, Hatano T, Tachibana M (2012) Followup of neutrophil-to-lymphocyte ratio and recurrence of clear cell renal cell carcinoma. *J Urol* 187(2):411–417. doi:10.1016/j.juro.2011.10.026
  154. Sarraf KM, Belcher E, Raevsky E, Nicholson AG, Goldstraw P, Lim E (2009) Neutrophil/lymphocyte ratio and its association with survival after complete resection in non-small cell lung cancer. *J*



- Thorac Cardiovasc Surg 137(2):425–428. doi:10.1016/j.jtcvs.2008.05.046
155. Sato T, Omura M, Saito J, Hirasawa A, Kakuta Y, Wakabayashi Y, Nishikawa T (2000) Neutrophilia associated with anaplastic carcinoma of the thyroid: production of macrophage colony-stimulating factor (M-CSF) and interleukin-6. *Thyroid* 10(12):1113–1118
  156. Schmidt H, Suci S, Punt CJ, Gore M, Kruit W, Patel P, Lienard D, von der Maase H, Eggermont AM, Keilholz U (2007) Pretreatment levels of peripheral neutrophils and leukocytes as independent predictors of overall survival in patients with American Joint Committee on Cancer Stage IV Melanoma: results of the EORTC 18951 Biochemotherapy Trial. *J Clin Oncol* 25(12):1562–1569. doi:10.1200/JCO.2006.09.0274
  157. Trellakis S, Bruderek K, Dumitru CA, Gholaman H, Gu X, Bankfalvi A, Scherag A, Hutte J, Dominas N, Lehnerdt GF, Hoffmann TK, Lang S, Brandau S (2011) Polymorphonuclear granulocytes in human head and neck cancer: enhanced inflammatory activity, modulation by cancer cells and expansion in advanced disease. *Int J Cancer* 129(9):2183–2193. doi:10.1002/ijc.25892
  158. Walsh SR, Cook EJ, Goulder F, Justin TA, Keeling NJ (2005) Neutrophil-lymphocyte ratio as a prognostic factor in colorectal cancer. *J Surg Oncol* 91(3):181–184. doi:10.1002/jso.20329
  159. Yamanaka T, Matsumoto S, Teramukai S, Ishiwata R, Nagai Y, Fukushima M (2007) The baseline ratio of neutrophils to lymphocytes is associated with patient prognosis in advanced gastric cancer. *Oncology* 73(3–4):215–220. doi:10.1159/000127412
  160. Melani C, Chiodoni C, Forni G, Colombo MP (2003) Myeloid cell expansion elicited by the progression of spontaneous mammary carcinomas in c-erbB-2 transgenic BALB/c mice suppresses immune reactivity. *Blood* 102(6):2138–2145. doi:10.1182/blood-2003-01-0190
  161. Kusmartsev S, Gabrilovich DI (2006) Role of immature myeloid cells in mechanisms of immune evasion in cancer. *Cancer Immunol Immunother* 55(3):237–245. doi:10.1007/s00262-005-0048-z
  162. Kusmartsev S, Cheng F, Yu B, Nefedova Y, Sotomayor E, Lush R, Gabrilovich D (2003) All-trans-retinoic acid eliminates immature myeloid cells from tumor-bearing mice and improves the effect of vaccination. *Cancer Res* 63(15):4441–4449
  163. Liu H, Tabuchi T, Takemura A, Kasuga T, Motohashi G, Hiraishi K, Katano M, Nakada I, Ubukata H (2008) The granulocyte/lymphocyte ratio as an independent predictor of tumour growth, metastasis and progression: its clinical applications. *Mol Med Rep* 1(5):699–704. doi:10.3892/mmr\_00000016
  164. Aeed PA, Nakajima M, Welch DR (1988) The role of polymorphonuclear leukocytes (PMN) on the growth and metastatic potential of 13762NF mammary adenocarcinoma cells. *Int J Cancer* 42(5):748–759
  165. McGary CT, Miele ME, Welch DR (1995) Highly metastatic 13762NF rat mammary adenocarcinoma cell clones stimulate bone marrow by secretion of granulocyte-macrophage colony-stimulating factor/interleukin-3 activity. *Am J Pathol* 147(6):1668–1681
  166. Welch DR, Schissel DJ, Howrey RP, Aeed PA (1989) Tumor-elicited polymorphonuclear cells, in contrast to “normal” circulating polymorphonuclear cells, stimulate invasive and metastatic potentials of rat mammary adenocarcinoma cells. *Proc Natl Acad Sci U S A* 86(15):5859–5863
  167. Opendakker G, Van Damme J (2004) The countercurrent principle in invasion and metastasis of cancer cells. Recent insights on the roles of chemokines. *Int J Dev Biol* 48(5–6):519–527. doi:10.1387/ijdb.041796go
  168. Schaidt H, Oka M, Bogenrieder T, Nesbit M, Satyamoorthy K, Berking C, Matsushima K, Herlyn M (2003) Differential response of primary and metastatic melanomas to neutrophils attracted by IL-8. *Int J Cancer* 103(3):335–343. doi:10.1002/ijc.10775
  169. Lee LF, Hellendall RP, Wang Y, Haskill JS, Mukaida N, Matsushima K, Ting JP (2000) IL-8 reduced tumorigenicity of human ovarian cancer in vivo due to neutrophil infiltration. *J Immunol* 164(5):2769–2775
  170. Bellocq A, Antoine M, Flahault A, Philippe C, Crestani B, Bernaudin JF, Mayaud C, Milleron B, Baud L, Cadranel J (1998) Neutrophil alveolitis in bronchioloalveolar carcinoma: induction by tumor-derived interleukin-8 and relation to clinical outcome. *Am J Pathol* 152(1):83–92
  171. Gijssbers K, Gouwy M, Struyf S, Wuyts A, Proost P, Opendakker G, Penninckx F, Ectors N, Geboes K, Van Damme J (2005) GCP-2/CXCL6 synergizes with other endothelial cell-derived chemokines in neutrophil mobilization and is associated with angiogenesis in gastrointestinal tumors. *Exp Cell Res* 303(2):331–342. doi:10.1016/j.yexcr.2004.09.027
  172. Asano Y, Yokoyama T, Shibata S, Kobayashi S, Shimoda K, Nakashima H, Okamura S, Niho Y (1997) Effect of the chimeric soluble granulocyte colony-stimulating factor receptor on the proliferation of leukemic blast cells from patients with acute myeloblastic leukemia. *Cancer Res* 57(16):3395–3397
  173. Chakraborty A, Guha S (2007) Granulocyte colony-stimulating factor/granulocyte colony-stimulating factor receptor biological axis promotes survival and growth of bladder cancer cells. *Urology* 69(6):1210–1215. doi:10.1016/j.urology.2007.02.035
  174. Joshita S, Nakazawa K, Sugiyama Y, Kamijo A, Matsubayashi K, Miyabayashi H, Furuta K, Kitano K, Kawa S (2009) Granulocyte-colony stimulating factor-producing pancreatic adenocarcinoma showing aggressive clinical course. *Intern Med* 48(9):687–691
  175. Kyo S, Kanaya T, Takakura M, Inoue M (2000) A case of cervical cancer with aggressive tumor growth: possible autocrine growth stimulation by G-CSF and IL-6. *Gynecol Oncol* 78(3 Pt 1):383–387. doi:10.1006/gyno.2000.5904
  176. Savarese TM, Mitchell K, McQuain C, Campbell CL, Guardiani R, Wu J, Ollari C, Reale F, Nelson BE, Chen A, Quesenberry PJ (2001) Coexpression of granulocyte colony stimulating factor and its receptor in primary ovarian carcinomas. *Cancer Lett* 162(1):105–115
  177. Tsukuda M, Nagahara T, Yago T, Matsuda H, Yanoma S (1993) Production of granulocyte colony-stimulating factor by head and neck carcinomas. *Biotherapy* 6(3):183–187
  178. Liu H, Zhang Z, Tabuchi T, Wang S, Wang J (2013) The role of pro-inflammatory cytokines and immune cells in colorectal carcinoma progression. *Oncol Lett* 5(4):1177–1182. doi:10.3892/ol.2013.1176
  179. Chavey C, Bibeau F, Gourgu-Bourgade S, Burlinon S, Boissiere F, Laune D, Roques S, Lazennec G (2007) Oestrogen receptor negative breast cancers exhibit high cytokine content. *Breast Cancer Res* 9(1):R15. doi:10.1186/bcr1648
  180. Urdinguio RG, Fernandez AF, Moncada-Pazos A, Huidobro C, Rodriguez RM, Ferrero C, Martinez-Cambor P, Obaya AJ, Bernal T, Parra-Blanco A, Rodrigo L, Santacana M, Matias-Guiu X, Soldevilla B, Dominguez G, Bonilla F, Cal S, Lopez-Otin C, Fraga MF (2013) Immune-dependent and independent antitumor activity of GM-CSF aberrantly expressed by mouse and human colorectal tumors. *Cancer Res* 73(1):395–405. doi:10.1158/0008-5472.CAN-12-0806
  181. Braun B, Lange M, Oeckler R, Mueller MM (2004) Expression of G-CSF and GM-CSF in human meningiomas correlates with increased tumor proliferation and vascularization. *J Neurooncol* 68(2):131–140
  182. Parmiani G, Castelli C, Pilla L, Santinami M, Colombo MP, Rivoltini L (2007) Opposite immune functions of GM-CSF administered as vaccine adjuvant in cancer patients. *Ann Oncol* 18(2):226–232. doi:10.1093/annonc/mdl158
  183. Filipazzi P, Valenti R, Huber V, Pilla L, Canese P, Iero M, Castelli C, Mariani L, Parmiani G, Rivoltini L (2007) Identification of a new subset of myeloid suppressor cells in peripheral blood of melanoma

- patients with modulation by a granulocyte-macrophage colony-stimulating factor-based antitumor vaccine. *J Clin Oncol* 25(18):2546–2553. doi:10.1200/JCO.2006.08.5829
184. Bronte V, Chappell DB, Apolloni E, Cabrelle A, Wang M, Hwu P, Restifo NP (1999) Unopposed production of granulocyte-macrophage colony-stimulating factor by tumors inhibits CD8+ T cell responses by dysregulating antigen-presenting cell maturation. *J Immunol* 162(10):5728–5737
  185. Serafini P, Carbley R, Noonan KA, Tan G, Bronte V, Borrello I (2004) High-dose granulocyte-macrophage colony-stimulating factor-producing vaccines impair the immune response through the recruitment of myeloid suppressor cells. *Cancer Res* 64(17):6337–6343. doi:10.1158/0008-5472.CAN-04-0757
  186. Fu YX, Watson G, Jimenez JJ, Wang Y, Lopez DM (1990) Expansion of immunoregulatory macrophages by granulocyte-macrophage colony-stimulating factor derived from a murine mammary tumor. *Cancer Res* 50(2):227–234
  187. Dranoff G, Jaffee E, Lazenby A, Golumbek P, Levitsky H, Brose K, Jackson V, Hamada H, Pardoll D, Mulligan RC (1993) Vaccination with irradiated tumor cells engineered to secrete murine granulocyte-macrophage colony-stimulating factor stimulates potent, specific, and long-lasting anti-tumor immunity. *Proc Natl Acad Sci U S A* 90(8):3539–3543
  188. Avalos BR, Gasson JC, Hedvat C, Quan SG, Baldwin GC, Weisbart RH, Williams RE, Golde DW, DiPersio JF (1990) Human granulocyte colony-stimulating factor: biologic activities and receptor characterization on hematopoietic cells and small cell lung cancer cell lines. *Blood* 75(4):851–857
  189. Kobashi Y, Okimoto N, Sakamoto K (2004) Squamous cell carcinoma of the lung producing granulocyte colony-stimulating factor and resembling a malignant pleural mesothelioma. *Intern Med* 43(2):111–116
  190. Hsu DM, Agarwal S, Benham A, Coarfa C, Trahan DN, Chen Z, Stowers PN, Courtney AN, Lakoma A, Barbieri E, Metelitsa LS, Gunaratne P, Kim ES, Shohet JM (2013) G-CSF receptor positive neuroblastoma subpopulations are enriched in chemotherapy-resistant or relapsed tumors and are highly tumorigenic. *Cancer Res* 73(13):4134–4146. doi:10.1158/0008-5472.CAN-12-4056
  191. Panopoulos AD, Watowich SS (2008) Granulocyte colony-stimulating factor: molecular mechanisms of action during steady state and ‘emergency’ hematopoiesis. *Cytokine* 42(3):277–288. doi:10.1016/j.cyto.2008.03.002
  192. Abrams SI, Waight JD (2012) Identification of a G-CSF-Granulocytic MDSC axis that promotes tumor progression. *Oncoimmunology* 1(4):550–551
  193. Waight JD, Hu Q, Miller A, Liu S, Abrams SI (2011) Tumor-derived G-CSF facilitates neoplastic growth through a granulocytic myeloid-derived suppressor cell-dependent mechanism. *PLoS One* 6(11):e27690. doi:10.1371/journal.pone.0027690
  194. Tachibana M, Murai M (1998) G-CSF production in human bladder cancer and its ability to promote autocrine growth: a review. *Cytokines Cell Mol Ther* 4(2):113–120
  195. Tachibana M, Miyakawa A, Uchida A, Murai M, Eguchi K, Nakamura K, Kubo A, Hata JI (1997) Granulocyte colony-stimulating factor receptor expression on human transitional cell carcinoma of the bladder. *Br J Cancer* 75(10):1489–1496
  196. Ninci EB, Brandstetter T, Meinhold-Heerlein I, Bettendorf H, Sellin D, Bauknecht T (2000) G-CSF receptor expression in ovarian cancer. *Int J Gynecol Cancer* 10(1):19–26
  197. Yang X, Liu F, Xu Z, Chen C, Wu X, Li G, Li J (2005) Expression of granulocyte colony stimulating factor receptor in human colorectal cancer. *Postgrad Med J* 81(955):333–337. doi:10.1136/pgmj.2004.024646
  198. Morales-Arias J, Meyers PA, Bolontrade MF, Rodriguez N, Zhou Z, Reddy K, Chou AJ, Koshkina NV, Kleinerman ES (2007) Expression of granulocyte-colony-stimulating factor and its receptor in human Ewing sarcoma cells and patient tumor specimens: potential consequences of granulocyte-colony-stimulating factor administration. *Cancer* 110(7):1568–1577. doi:10.1002/cncr.22964
  199. Joyce JA, Pollard JW (2009) Microenvironmental regulation of metastasis. *Nat Rev Cancer* 9(4):239–252
  200. Caruso RA, Bellocco R, Pagano M, Bertoli G, Rigoli L, Inferrea C (2002) Prognostic value of intratumoral neutrophils in advanced gastric carcinoma in a high-risk area in northern Italy. *Mod Pathol* 15(8):831–837. doi:10.1097/01.MP.0000020391.98998.6B
  201. Griffiths AP, Rice A, Dixon MF (1998) Anaplastic gastric adenocarcinoma with extensive neutrophilic infiltration. *Histopathology* 33(4):392–393
  202. Rice AJ, Griffiths AP, Martin IG, Dixon MF (2000) Gastric carcinoma with prominent neutrophil infiltration. *Histopathology* 37(3):289–290
  203. Wislez M, Rabbe N, Marchal J, Milleron B, Crestani B, Mayaud C, Antoine M, Soler P, Cadranet J (2003) Hepatocyte growth factor production by neutrophils infiltrating bronchioloalveolar subtype pulmonary adenocarcinoma: role in tumor progression and death. *Cancer Res* 63(6):1405–1412
  204. Ilie M, Hofman V, Ortholan C, Bonnetaud C, Coelle C, Mouroux J, Hofman P (2012) Predictive clinical outcome of the intratumoral CD66b-positive neutrophil-to-CD8-positive T-cell ratio in patients with resectable nonsmall cell lung cancer. *Cancer* 118(6):1726–1737. doi:10.1002/cncr.26456
  205. Reid MD, Basturk O, Thirabanjasak D, Hruban RH, Klimstra DS, Bagci P, Altinel D, Adsay V (2011) Tumor-infiltrating neutrophils in pancreatic neoplasia. *Mod Pathol* 24(12):1612–1619. doi:10.1038/modpathol.2011.113
  206. Gaida MM, Steffen TG, Gunther F, Tschaharganeh DF, Felix K, Bergmann F, Schirmacher P, Hansch GM (2012) Polymorphonuclear neutrophils promote dyshesion of tumor cells and elastase-mediated degradation of E-cadherin in pancreatic tumors. *Eur J Immunol* 42(12):3369–3380. doi:10.1002/eji.201242628
  207. Fossati G, Ricevuti G, Edwards SW, Walker C, Dalton A, Rossi ML (1999) Neutrophil infiltration into human gliomas. *Acta Neuropathol* 98(4):349–354
  208. O’Sullivan C, Lewis CE (1994) Tumour-associated leucocytes: friends or foes in breast carcinoma. *J Pathol* 172(3):229–235. doi:10.1002/path.1711720302
  209. Ancrile BB, O’Hayer KM, Counter CM (2008) Oncogenic ras-induced expression of cytokines: a new target of anti-cancer therapeutics. *Mol Interv* 8(1):22–27. doi:10.1124/mi.8.1.6
  210. Spammann A, Bar-Sagi D (2004) Ras-induced interleukin-8 expression plays a critical role in tumor growth and angiogenesis. *Cancer Cell* 6(5):447–458. doi:10.1016/j.ccr.2004.09.028
  211. Lazennec G, Richmond A (2010) Chemokines and chemokine receptors: new insights into cancer-related inflammation. *Trends Mol Med* 16(3):133–144. doi:10.1016/j.molmed.2010.01.003
  212. Wuyts A, Proost P, Lenaerts JP, Ben-Baruch A, Van Damme J, Wang JM (1998) Differential usage of the CXC chemokine receptors 1 and 2 by interleukin-8, granulocyte chemotactic protein-2 and epithelial-cell-derived neutrophil attractant-78. *Eur J Biochem* 255(1):67–73
  213. Ahuja SK, Murphy PM (1996) The CXC chemokines growth-regulated oncogene (GRO) alpha, GRObeta, GROgamma, neutrophil-activating peptide-2, and epithelial cell-derived neutrophil-activating peptide-78 are potent agonists for the type B, but not the type A, human interleukin-8 receptor. *J Biol Chem* 271(34):20545–20550
  214. Song J, Wu C, Zhang X, Sorokin LM (2013) In vivo processing of CXCL5 (LIX) by matrix metalloproteinase (MMP)-2 and MMP-9 promotes early neutrophil recruitment in IL-1beta-induced peritonitis. *J Immunol* 190(1):401–410. doi:10.4049/jimmunol.1202286
  215. Schenk BI, Petersen F, Flad HD, Brandt E (2002) Platelet-derived chemokines CXC chemokine ligand (CXCL)7, connective tissue-

- activating peptide III, and CXCL4 differentially affect and cross-regulate neutrophil adhesion and transendothelial migration. *J Immunol* 169(5):2602–2610
216. Bru A, Souto JC, Alcolea S, Anton R, Remacha A, Camacho M, Soler M, Bru I, Porres A, Vila L (2009) Tumour cell lines HT-29 and FaDu produce proinflammatory cytokines and activate neutrophils in vitro: possible applications for neutrophil-based antitumour treatment. *Mediators Inflamm* 2009:817498. doi:10.1155/2009/817498
  217. Yamashiro S, Kamohara H, Wang JM, Yang D, Gong WH, Yoshimura T (2001) Phenotypic and functional change of cytokine-activated neutrophils: inflammatory neutrophils are heterogeneous and enhance adaptive immune responses. *J Leukoc Biol* 69(5):698–704
  218. Pham CT (2006) Neutrophil serine proteases: specific regulators of inflammation. *Nat Rev Immunol* 6(7):541–550. doi:10.1038/nri1841
  219. Jamieson T, Clarke M, Steele CW, Samuel MS, Neumann J, Jung A, Huels D, Olson MF, Das S, Nibbs RJ, Sansom OJ (2012) Inhibition of CXCR2 profoundly suppresses inflammation-driven and spontaneous tumorigenesis. *J Clin Invest* 122(9):3127–3144. doi:10.1172/JCI161067
  220. Verbeke H, Struyf S, Berghmans N, Van Coillie E, Opendakker G, Uyttenhove C, Van Snick J, Van Damme J (2011) Isotypic neutralizing antibodies against mouse GCP-2/CXCL6 inhibit melanoma growth and metastasis. *Cancer Lett* 302(1):54–62. doi:10.1016/j.canlet.2010.12.013
  221. Richards H, Williams A, Jones E, Hindley J, Godkin A, Simon AK, Gallimore A (2010) Novel role of regulatory T cells in limiting early neutrophil responses in skin. *Immunology* 131(4):583–592. doi:10.1111/j.1365-2567.2010.03333.x
  222. Lyden D, Hattori K, Dias S, Costa C, Blaikie P, Butros L, Chadburn A, Heissig B, Marks W, Witte L, Wu Y, Hicklin D, Zhu Z, Hackett NR, Crystal RG, Moore MA, Hajjar KA, Manova K, Benezra R, Rafii S (2001) Impaired recruitment of bone-marrow-derived endothelial and hematopoietic precursor cells blocks tumor angiogenesis and growth. *Nat Med* 7(11):1194–1201. doi:10.1038/nm1101-1194
  223. Grunewald M, Avraham I, Dor Y, Bachar-Lustig E, Itin A, Jung S, Chimenti S, Landsman L, Abramovitch R, Keshet E (2006) VEGF-induced adult neovascularization: recruitment, retention, and role of accessory cells. *Cell* 124(1):175–189. doi:10.1016/j.cell.2005.10.036
  224. Kusmartsev S, Gabrilovich DI (2006) Effect of tumor-derived cytokines and growth factors on differentiation and immune suppressive features of myeloid cells in cancer. *Cancer Metastasis Rev* 25(3):323–331. doi:10.1007/s10555-006-9002-6
  225. Ji YN, Wang Q, Li Y, Wang Z (2013) Prognostic value of vascular endothelial growth factor A expression in gastric cancer: a meta-analysis. *Tumour Biol*. doi:10.1007/s13277-013-1371-1
  226. Minardi D, Lucarini G, Santoni M, Mazzucchelli R, Burattini L, Pistelli M, Bianconi M, Di Primio R, Scartozzi M, Montironi R, Cascinu S, Muzzonigro G (2013) VEGF expression and response to sunitinib in patients with metastatic clear cell renal cell carcinoma. *Anticancer Res* 33(11):5017–5022
  227. Ohm JE, Carbone DP (2001) VEGF as a mediator of tumor-associated immunodeficiency. *Immunol Res* 23(2–3):263–272. doi:10.1385/IR:23-2-3:263
  228. Salmaggi A, Gelati M, Pollo B, Frigerio S, Eoli M, Silvani A, Broggi G, Ciusani E, Croci D, Boiardi A, De Rossi M (2004) CXCL12 in malignant glial tumors: a possible role in angiogenesis and cross-talk between endothelial and tumoral cells. *J Neurooncol* 67(3):305–317
  229. Santiago B, Calonge E, Del Rey MJ, Gutierrez-Canas I, Izquierdo E, Usategui A, Galindo M, Alcami J, Pablos JL (2011) CXCL12 gene expression is upregulated by hypoxia and growth arrest but not by inflammatory cytokines in rheumatoid synovial fibroblasts. *Cytokine* 53(2):184–190. doi:10.1016/j.cyto.2010.06.006
  230. Kryczek I, Lange A, Mottram P, Alvarez X, Cheng P, Hogan M, Moons L, Wei S, Zou L, Machelon V, Emilie D, Terrassa M, Lackner A, Curiel TJ, Carmeliet P, Zou W (2005) CXCL12 and vascular endothelial growth factor synergistically induce neoangiogenesis in human ovarian cancers. *Cancer Res* 65(2):465–472
  231. Chung AS, Wu X, Zhuang G, Ngu H, Kasman I, Zhang J, Vernes JM, Jiang Z, Meng YG, Peale FV, Ouyang W, Ferrara N (2013) An interleukin-17-mediated paracrine network promotes tumor resistance to anti-angiogenic therapy. *Nat Med*. doi:10.1038/nm.3291
  232. Martin-Orozco N, Muranski P, Chung Y, Yang XO, Yamazaki T, Lu S, Hwu P, Restifo NP, Overwijk WW, Dong C (2009) T helper 17 cells promote cytotoxic T cell activation in tumor immunity. *Immunity* 31(5):787–798. doi:10.1016/j.immuni.2009.09.014
  233. Wang Z, Liu JQ, Liu Z, Shen R, Zhang G, Xu J, Basu S, Feng Y, Bai XF (2013) Tumor-derived IL-35 promotes tumor growth by enhancing myeloid cell accumulation and angiogenesis. *J Immunol* 190(5):2415–2423. doi:10.4049/jimmunol.1202535
  234. Mantovani A, Sica A (2010) Macrophages, innate immunity and cancer: balance, tolerance, and diversity. *Curr Opin Immunol* 22(2):231–237. doi:10.1016/j.coi.2010.01.009
  235. Bailey C, Negus R, Morris A, Ziprin P, Goldin R, Allavena P, Peck D, Darzi A (2007) Chemokine expression is associated with the accumulation of tumour associated macrophages (TAMs) and progression in human colorectal cancer. *Clin Exp Metastasis* 24(2):121–130. doi:10.1007/s10585-007-9060-3
  236. Saji H, Koike M, Yamori T, Saji S, Seiki M, Matsushima K, Toi M (2001) Significant correlation of monocyte chemoattractant protein-1 expression with neovascularization and progression of breast carcinoma. *Cancer* 92(5):1085–1091. doi:10.1002/1097-0142(20010901)92:5<1085::AID-CNCR1424>3.0.CO;2-K
  237. Tanaka K, Kurebayashi J, Sohda M, Nomura T, Prabhakar U, Yan L, Sonoo H (2009) The expression of monocyte chemoattractant protein-1 in papillary thyroid carcinoma is correlated with lymph node metastasis and tumor recurrence. *Thyroid* 19(1):21–25. doi:10.1089/thy.2008.0237
  238. Yoshidome H, Kohno H, Shida T, Kimura F, Shimizu H, Ohtsuka M, Nakatani Y, Miyazaki M (2009) Significance of monocyte chemoattractant protein-1 in angiogenesis and survival in colorectal liver metastases. *Int J Oncol* 34(4):923–930
  239. Zijlmans HJ, Fleuren GJ, Baelde HJ, Eilers PH, Kenter GG, Gorter A (2006) The absence of CCL2 expression in cervical carcinoma is associated with increased survival and loss of heterozygosity at 17q11.2. *J Pathol* 208(4):507–517. doi:10.1002/path.1918
  240. Qian BZ, Li J, Zhang H, Kitamura T, Zhang J, Campion LR, Kaiser EA, Snyder LA, Pollard JW (2011) CCL2 recruits inflammatory monocytes to facilitate breast-tumour metastasis. *Nature* 475(7355):222–225. doi:10.1038/nature10138. nature10138
  241. Huang B, Lei Z, Zhao J, Gong W, Liu J, Chen Z, Liu Y, Li D, Yuan Y, Zhang GM, Feng ZH (2007) CCL2/CCR2 pathway mediates recruitment of myeloid suppressor cells to cancers. *Cancer Lett* 252(1):86–92. doi:10.1016/j.canlet.2006.12.012
  242. Nakashima E, Mukaida N, Kubota Y, Kuno K, Yasumoto K, Ichimura F, Nakanishi I, Miyasaka M, Matsushima K (1995) Human MCAF gene transfer enhances the metastatic capacity of a mouse cachectic adenocarcinoma cell line in vivo. *Pharm Res* 12(11):1598–1604
  243. Zhang J, Lu Y, Pienta KJ (2010) Multiple roles of chemokine (C-C motif) ligand 2 in promoting prostate cancer growth. *J Natl Cancer Inst* 102(8):522–528. doi:10.1093/jnci/djq044
  244. Loberg RD, Ying C, Craig M, Day LL, Sargent E, Neeley C, Wojno K, Snyder LA, Yan L, Pienta KJ (2007) Targeting CCL2 with systemic delivery of neutralizing antibodies induces prostate cancer tumor regression in vivo. *Cancer Res* 67(19):9417–9424
  245. Yang EJ, Choi E, Ko J, Kim DH, Lee JS, Kim IS (2012) Differential effect of CCL2 on constitutive neutrophil apoptosis between normal



- and asthmatic subjects. *J Cell Physiol* 227(6):2567–2577. doi:10.1002/jcp.22995
246. Bolitho C, Hahn MA, Baxter RC, Marsh DJ (2010) The chemokine CXCL1 induces proliferation in epithelial ovarian cancer cells by transactivation of the epidermal growth factor receptor. *Endocr Relat Cancer* 17(4):929–940. doi:10.1677/ERC-10-0107
  247. Strieter RM, Polverini PJ, Kunkel SL, Arenberg DA, Burdick MD, Kasper J, Dzuiba J, Van Damme J, Walz A, Marriott D et al (1995) The functional role of the ELR motif in CXC chemokine-mediated angiogenesis. *J Biol Chem* 270(45):27348–27357
  248. Kuo PL, Shen KH, Hung SH, Hsu YL (2012) CXCL1/GROalpha increases cell migration and invasion of prostate cancer by decreasing fibulin-1 expression through NF-kappaB/HDAC1 epigenetic regulation. *Carcinogenesis* 33(12):2477–2487. doi:10.1093/carcin/bgs299
  249. Scapini P, Morini M, Tecchio C, Minghelli S, Di Carlo E, Tanghetti E, Albini A, Lowell C, Berton G, Noonan DM, Cassatella MA (2004) CXCL1/macrophage inflammatory protein-2-induced angiogenesis in vivo is mediated by neutrophil-derived vascular endothelial growth factor-A. *J Immunol* 172(8):5034–5040
  250. Acharyya S, Oskarsson T, Vanharanta S, Malladi S, Kim J, Morris PG, Manova-Todorova K, Leversha M, Hogg N, Seshan VE, Norton L, Brogi E, Massague J (2012) A CXCL1 paracrine network links cancer chemoresistance and metastasis. *Cell* 150(1):165–178
  251. Nasser MW, Raghuvanshi SK, Grant DJ, Jala VR, Rajarathnam K, Richardson RM (2009) Differential activation and regulation of CXCR1 and CXCR2 by CXCL8 monomer and dimer. *J Immunol* 183(5):3425–3432. doi:10.4049/jimmunol.0900305
  252. Bekes EM, Schweighofer B, Kupriyanova TA, Zajac E, Ardi VC, Quigley JP, Deryugina EI (2011) Tumor-recruited neutrophils and neutrophil TIMP-free MMP-9 regulate coordinately the levels of tumor angiogenesis and efficiency of malignant cell intravasation. *Am J Pathol* 179(3):1455–1470. doi:10.1016/j.ajpath.2011.05.031
  253. Dong C, Robertson GP (2009) Immunoediting of leukocyte functions within the tumor microenvironment promotes cancer metastasis development. *Biorheology* 46(4):265–279. doi:10.3233/BIR-2009-0545
  254. Slattery MJ, Liang S, Dong C (2005) Distinct role of hydrodynamic shear in leukocyte-facilitated tumor cell extravasation. *Am J Physiol Cell Physiol* 288(4):C831–C839. doi:10.1152/ajpcell.00439.2004
  255. Nakamoto Y, Guidotti LG, Kuhlen CV, Fowler P, Chisari FV (1998) Immune pathogenesis of hepatocellular carcinoma. *J Exp Med* 188(2):341–350
  256. Rogler G (2013) Chronic ulcerative colitis and colorectal cancer. *Cancer Lett*. doi:10.1016/j.canlet.2013.07.032
  257. Haqqani AS, Sandhu JK, Birnboim HC (2000) Expression of interleukin-8 promotes neutrophil infiltration and genetic instability in mutect tumors. *Neoplasia* 2(6):561–568
  258. Knaapen AM, Gungor N, Schins RP, Borm PJ, Van Schooten FJ (2006) Neutrophils and respiratory tract DNA damage and mutagenesis: a review. *Mutagenesis* 21(4):225–236. doi:10.1093/mutage/gel032
  259. Gungör N, Knaapen AM, Munnia A, Peluso M, Haenen GR, Chiu RK, Godschalk RW, van Schooten FJ (2010) Genotoxic effects of neutrophils and hypochlorous acid. *Mutagenesis* 25(2):149–154. doi:10.1093/mutage/gep053
  260. Gomez-Mejiba SE, Zhai Z, Gimenez MS, Ashby MT, Chilakapati J, Kitchin K, Mason RP, Ramirez DC (2010) Myeloperoxidase-induced genomic DNA-centered radicals. *J Biol Chem* 285(26):20062–20071. doi:10.1074/jbc.M109.086579
  261. Shang K, Bai YP, Wang C, Wang Z, Gu HY, Du X, Zhou XY, Zheng CL, Chi YY, Mukaida N, Li YY (2012) Crucial involvement of tumor-associated neutrophils in the regulation of chronic colitis-associated carcinogenesis in mice. *PLoS One* 7(12):e51848. doi:10.1371/journal.pone.0051848
  262. Ginzberg HH, Cherapanov V, Dong Q, Cantin A, McCulloch CA, Shannon PT, Downey GP (2001) Neutrophil-mediated epithelial injury during transmigration: role of elastase. *Am J Physiol Gastrointest Liver Physiol* 281(3):G705–G717
  263. Grosse-Steffen T, Giese T, Giese N, Longrich T, Schirmacher P, Hansch GM, Gaida MM (2012) Epithelial-to-mesenchymal transition in pancreatic ductal adenocarcinoma and pancreatic tumor cell lines: the role of neutrophils and neutrophil-derived elastase. *Clin Dev Immunol* 2012:720768. doi:10.1155/2012/720768
  264. Pekarek LA, Starr BA, Toledano AY, Schreiber H (1995) Inhibition of tumor growth by elimination of granulocytes. *J Exp Med* 181(1):435–440
  265. Tazawa H, Okada F, Kobayashi T, Tada M, Mori Y, Une Y, Sendo F, Kobayashi M, Hosokawa M (2003) Infiltration of neutrophils is required for acquisition of metastatic phenotype of benign murine fibrosarcoma cells: implication of inflammation-associated carcinogenesis and tumor progression. *Am J Pathol* 163(6):2221–2232. doi:10.1016/S0002-9440(10)63580-8
  266. Coussens LM, Tinkle CL, Hanahan D, Werb Z (2000) MMP-9 supplied by bone marrow-derived cells contributes to skin carcinogenesis. *Cell* 103(3):481–490
  267. Balbin M, Fueyo A, Tester AM, Pendas AM, Pitiot AS, Astudillo A, Overall CM, Shapiro SD, Lopez-Otin C (2003) Loss of collagenase-2 confers increased skin tumor susceptibility to male mice. *Nat Genet* 35(3):252–257. doi:10.1038/ng1249
  268. Bodey B, Bodey B Jr, Siegel SE, Luck JV, Kaiser HE (1996) Immunophenotypic characterization of human primary and metastatic melanoma infiltrating leukocytes. *Anticancer Res* 16(6B):3439–3446
  269. Yu J, Ren X, Chen Y, Liu P, Wei X, Li H, Ying G, Chen K, Winkler H, Hao X (2013) Dysfunctional activation of neurotensin/IL-8 pathway in hepatocellular carcinoma is associated with increased inflammatory response in microenvironment, more epithelial mesenchymal transition in cancer and worse prognosis in patients. *PLoS One* 8(2):e56069. doi:10.1371/journal.pone.0056069
  270. Xie K (2001) Interleukin-8 and human cancer biology. *Cytokine Growth Factor Rev* 12(4):375–391
  271. Lippitz BE (2013) Cytokine patterns in patients with cancer: a systematic review. *Lancet Oncol* 14(6):e218–e228. doi:10.1016/S1470-2045(12)70582-X
  272. Waugh DJ, Wilson C (2008) The interleukin-8 pathway in cancer. *Clin Cancer Res* 14(21):6735–6741. doi:10.1158/1078-0432.CCR-07-4843
  273. Anderson IC, Mari SE, Broderick RJ, Mari BP, Shipp MA (2000) The angiogenic factor interleukin 8 is induced in non-small cell lung cancer/pulmonary fibroblast cocultures. *Cancer Res* 60(2):269–272
  274. Chen JJ, Yao PL, Yuan A, Hong TM, Shun CT, Kuo ML, Lee YC, Yang PC (2003) Up-regulation of tumor interleukin-8 expression by infiltrating macrophages: its correlation with tumor angiogenesis and patient survival in non-small cell lung cancer. *Clin Cancer Res* 9(2):729–737
  275. Mueller MM, Herold-Mende CC, Riede D, Lange M, Steiner HH, Fusenig NE (1999) Autocrine growth regulation by granulocyte colony-stimulating factor and granulocyte macrophage colony-stimulating factor in human gliomas with tumor progression. *Am J Pathol* 155(5):1557–1567. doi:10.1016/S0002-9440(10)65472-7
  276. Yamano T, Morii E, Ikeda J, Aozasa K (2007) Granulocyte colony-stimulating factor production and rapid progression of gastric cancer after histological change in the tumor. *Jpn J Clin Oncol* 37(10):793–796. doi:10.1093/jjco/hym094
  277. Shojaei F, Singh M, Thompson JD, Ferrara N (2008) Role of Bv8 in neutrophil-dependent angiogenesis in a transgenic model of cancer progression. *Proc Natl Acad Sci U S A* 105(7):2640–2645. doi:10.1073/pnas.0712185105
  278. Schmielau J, Finn OJ (2001) Activated granulocytes and granulocyte-derived hydrogen peroxide are the underlying



- mechanism of suppression of t-cell function in advanced cancer patients. *Cancer Res* 61(12):4756–4760
279. Okazaki T, Ebihara S, Asada M, Kanda A, Sasaki H, Yamaya M (2006) Granulocyte colony-stimulating factor promotes tumor angiogenesis via increasing circulating endothelial progenitor cells and Gr1+CD11b+ cells in cancer animal models. *Int Immunol* 18(1):1–9. doi:10.1093/intimm/dxh334
  280. Natori T, Sata M, Washida M, Hirata Y, Nagai R, Makuuchi M (2002) G-CSF stimulates angiogenesis and promotes tumor growth: potential contribution of bone marrow-derived endothelial progenitor cells. *Biochem Biophys Res Commun* 297(4):1058–1061
  281. Voloshin T, Gingis-Velitski S, Bril R, Benayoun L, Munster M, Milsom C, Man S, Kerbel RS, Shaked Y (2011) G-CSF supplementation with chemotherapy can promote revascularization and subsequent tumor regrowth: prevention by a CXCR4 antagonist. *Blood* 118(12):3426–3435. doi:10.1182/blood-2010-11-320812
  282. Shojaei F, Ferrara N (2008) Refractoriness to antivascular endothelial growth factor treatment: role of myeloid cells. *Cancer Res* 68(14):5501–5504. doi:10.1158/0008-5472.CAN-08-0925
  283. Shojaei F, Wu X, Zhong C, Yu L, Liang XH, Yao J, Blanchard D, Bais C, Peale FV, van Bruggen N, Ho C, Ross J, Tan M, Carano RA, Meng YG, Ferrara N (2007) Bv8 regulates myeloid-cell-dependent tumour angiogenesis. *Nature* 450(7171):825–831. doi:10.1038/nature06348
  284. Qu X, Zhuang G, Yu L, Meng G, Ferrara N (2012) Induction of Bv8 expression by granulocyte colony-stimulating factor in CD11b+ Gr1+ cells: key role of Stat3 signaling. *J Biol Chem* 287(23):19574–19584. doi:10.1074/jbc.M111.326801
  285. Ohki Y, Heissig B, Sato Y, Akiyama H, Zhu Z, Hicklin DJ, Shimada K, Ogawa H, Daida H, Hattori K, Ohsaka A (2005) Granulocyte colony-stimulating factor promotes neovascularization by releasing vascular endothelial growth factor from neutrophils. *FASEB J* 19(14):2005–2007. doi:10.1096/fj.04-3496fje
  286. Bergers G, Brekken R, McMahon G, Vu TH, Itoh T, Tamaki K, Tanzawa K, Thorpe P, Itohara S, Werb Z, Hanahan D (2000) Matrix metalloproteinase-9 triggers the angiogenic switch during carcinogenesis. *Nat Cell Biol* 2(10):737–744. doi:10.1038/35036374
  287. Nozawa H, Chiu C, Hanahan D (2006) Infiltrating neutrophils mediate the initial angiogenic switch in a mouse model of multi-stage carcinogenesis. *Proc Natl Acad Sci U S A* 103(33):12493–12498. doi:10.1073/pnas.0601807103
  288. Imai Y, Kubota Y, Yamamoto S, Tsuji K, Shimatani M, Shibatani N, Takamido S, Matsushita M, Okazaki K (2005) Neutrophils enhance invasion activity of human cholangiocellular carcinoma and hepatocellular carcinoma cells: an in vitro study. *J Gastroenterol Hepatol* 20(2):287–293. doi:10.1111/j.1440-1746.2004.03575.x
  289. Di Carlo E, Forni G, Musiani P (2003) Neutrophils in the antitumoral immune response. *Chem Immunol Allergy* 83:182–203
  290. Shamamian P, Schwartz JD, Pocock BJ, Monea S, Whiting D, Marcus SG, Mignatti P (2001) Activation of progelatinase A (MMP-2) by neutrophil elastase, cathepsin G, and proteinase-3: a role for inflammatory cells in tumor invasion and angiogenesis. *J Cell Physiol* 189(2):197–206. doi:10.1002/jcp.10014
  291. De Larco JE, Wuertz BR, Furcht LT (2004) The potential role of neutrophils in promoting the metastatic phenotype of tumors releasing interleukin-8. *Clin Cancer Res* 10(15):4895–4900. doi:10.1158/1078-0432.CCR-03-0760
  292. Sun Z, Yang P (2004) Role of imbalance between neutrophil elastase and alpha 1-antitrypsin in cancer development and progression. *Lancet Oncol* 5(3):182–190. doi:10.1016/S1470-2045(04)01414-7
  293. Slattery MJ, Dong C (2003) Neutrophils influence melanoma adhesion and migration under flow conditions. *Int J Cancer* 106(5):713–722. doi:10.1002/ijc.11297
  294. Huh SJ, Liang S, Sharma A, Dong C, Robertson GP (2010) Transiently entrapped circulating tumor cells interact with neutrophils to facilitate lung metastasis development. *Cancer Res* 70(14):6071–6082. doi:10.1158/0008-5472.CAN-09-4442
  295. Drost EM, MacNee W (2002) Potential role of IL-8, platelet-activating factor and TNF-alpha in the sequestration of neutrophils in the lung: effects on neutrophil deformability, adhesion receptor expression, and chemotaxis. *Eur J Immunol* 32(2):393–403. doi:10.1002/1521-4141(200202)32:2<393::AID-IMMU393>3.0.CO;2-5
  296. Wu QD, Wang JH, Condron C, Bouchier-Hayes D, Redmond HP (2001) Human neutrophils facilitate tumor cell transendothelial migration. *Am J Physiol Cell Physiol* 280(4):C814–C822
  297. Jadhav S, Bochner BS, Konstantopoulos K (2001) Hydrodynamic shear regulates the kinetics and receptor specificity of polymorphonuclear leukocyte-colon carcinoma cell adhesive interactions. *J Immunol* 167(10):5986–5993
  298. Jadhav S, Konstantopoulos K (2002) Fluid shear- and time-dependent modulation of molecular interactions between PMNs and colon carcinomas. *Am J Physiol Cell Physiol* 283(4):C1133–C1143. doi:10.1152/ajpcell.00104.2002
  299. Hiratsuka S, Nakamura K, Iwai S, Murakami M, Itoh T, Kijima H, Shipley JM, Senior RM, Shibuya M (2002) MMP9 induction by vascular endothelial growth factor receptor-1 is involved in lung-specific metastasis. *Cancer Cell* 2(4):289–300
  300. Kaplan RN, Riba RD, Zacharoulis S, Bramley AH, Vincent L, Costa C, MacDonald DD, Jin DK, Shido K, Kerns SA, Zhu Z, Hicklin D, Wu Y, Port JL, Altorki N, Port ER, Ruggero D, Shmelkov SV, Jensen KK, Rafii S, Lyden D (2005) VEGFR1-positive haematopoietic bone marrow progenitors initiate the premetastatic niche. *Nature* 438(7069):820–827
  301. Hiratsuka S, Watanabe A, Aburatani H, Maru Y (2006) Tumour-mediated upregulation of chemoattractants and recruitment of myeloid cells predetermines lung metastasis. *Nat Cell Biol* 8(12):1369–1375
  302. Yan HH, Pickup M, Pang Y, Gorska AE, Li Z, Chytil A, Geng Y, Gray JW, Moses HL, Yang L (2010) Gr-1+CD11b+ myeloid cells tip the balance of immune protection to tumor promotion in the premetastatic lung. *Cancer Res* 70(15):6139–6149. doi:10.1158/0008-5472.CAN-10-0706
  303. Gao D, Joshi N, Choi H, Ryu S, Hahn M, Catena R, Sadik H, Argani P, Wagner P, Vahdat LT, Port JL, Stiles B, Sukumar S, Altorki NK, Rafii S, Mittal V (2012) Myeloid progenitor cells in the premetastatic lung promote metastases by inducing mesenchymal to epithelial transition. *Cancer Res* 72(6):1384–1394
  304. Sceneay J, Chow MT, Chen A, Halse HM, Wong CS, Andrews DM, Sloan EK, Parker BS, Bowtell DD, Smyth MJ, Moller A (2012) Primary tumor hypoxia recruits CD11b+Ly6Cmed/Ly6G+ immune suppressor cells and compromises NK cell cytotoxicity in the premetastatic niche. *Cancer Res* 72(16):3906–3911. doi:10.1158/0008-5472.CAN-11-3873
  305. Erler JT, Bennewith KL, Cox TR, Lang G, Bird D, Koong A, Le QT, Giaccia AJ (2009) Hypoxia-induced lysyl oxidase is a critical mediator of bone marrow cell recruitment to form the premetastatic niche. *Cancer Cell* 15(1):35–44. doi:10.1016/j.ccr.2008.11.012
  306. Cools-Lartigue J, Spicer J, McDonald B, Gowing S, Chow S, Giannias B, Bourdeau F, Kubes P, Ferri L (2013) Neutrophil extracellular traps sequester circulating tumor cells and promote metastasis. *J Clin Invest*. doi:10.1172/JCI67484
  307. Spicer JD, McDonald B, Cools-Lartigue JJ, Chow SC, Giannias B, Kubes P, Ferri LE (2012) Neutrophils promote liver metastasis via Mac-1-mediated interactions with circulating tumor cells. *Cancer Res* 72(16):3919–3927. doi:10.1158/0008-5472.CAN-11-2393
  308. Bubenik J, Perlmann P, Helmstein K, Moberger G (1970) Cellular and humoral immune responses to human urinary bladder carcinomas. *Int J Cancer* 5(3):310–319
  309. Shau H (1988) Cytostatic and tumoricidal activities of tumor necrosis factor-treated neutrophils. *Immunol Lett* 17(1):47–51

310. Shau H (1988) Characteristics and mechanism of neutrophil-mediated cytostasis induced by tumor necrosis factor. *J Immunol* 141(1):234–240
311. Miyake Y, Ajitsu S, Yamashita T, Sendo F (1988) Enhancement by recombinant interferon-gamma of spontaneous tumor cytostasis by human neutrophils. *Mol Biother* 1(1):37–42
312. Yamashita T, Uchida T, Araki A, Sendo F (1997) Nitric oxide is an effector molecule in inhibition of tumor cell growth by rIFN-gamma-activated rat neutrophils. *Int J Cancer* 71(2):223–230. doi:10.1002/(SICI)1097-0215(19970410)71:2<223::AID-IJC17>3.0.CO;2-I
313. Uchida T, Yamashita T, Araki A, Watanabe H, Sendo F (1997) rIFN-gamma-activated rat neutrophils induce tumor cell apoptosis by nitric oxide. *Int J Cancer* 71(2):231–236. doi:10.1002/(SICI)1097-0215(19970410)71:2<231::AID-IJC18>3.0.CO;2-K
314. Slivka A, LoBuglio AF, Weiss SJ (1980) A potential role for hypochlorous acid in granulocyte-mediated tumor cell cytotoxicity. *Blood* 55(2):347–350
315. Clark RA, Klebanoff SJ (1979) Role of the myeloperoxidase-H<sub>2</sub>O<sub>2</sub>-halide system in concanavalin A-induced tumor cell killing by human neutrophils. *J Immunol* 122(6):2605–2610
316. Weiss SJ, Slivka A (1982) Monocyte and granulocyte-mediated tumor cell destruction. A role for the hydrogen peroxide-myeloperoxidase-chloride system. *J Clin Invest* 69(2):255–262
317. Dallegri F, Frumento G, Patrone F (1983) Mechanisms of tumour cell destruction by PMA-activated human neutrophils. *Immunology* 48(2):273–279
318. Abrams SI, Brahmī Z (1984) Compared mechanisms of tumor cytotoxicity by human natural killer cells and activated polymorphonuclear leukocytes. *J Immunol* 132(6):3192–3196
319. Bru A, Albertos S, Lopez Garcia-Asenjo JA, Bru I (2004) Pinning of tumoral growth by enhancement of the immune response. *Phys Rev Lett* 92(23):238101
320. Su YB, Vickers AJ, Zelefsky MJ, Kraus DH, Shaha AR, Shah JP, Serio AM, Harrison LB, Bosl GJ, Pfister DG (2006) Double-blind, placebo-controlled, randomized trial of granulocyte-colony stimulating factor during postoperative radiotherapy for squamous head and neck cancer. *Cancer J* 12(3):182–188
321. Rini BI, Fong L, Weinberg V, Kavanaugh B, Small EJ (2006) Clinical and immunological characteristics of patients with serologic progression of prostate cancer achieving long-term disease control with granulocyte-macrophage colony-stimulating factor. *J Urol* 175(6):2087–2091. doi:10.1016/S0022-5347(06)00261-8
322. Bottoni U, Bonaccorsi P, Devirgiliis V, Panasiti V, Borroni RG, Trasimeni G, Clerico R, Calvieri S (2005) Complete remission of brain metastases in three patients with stage IV melanoma treated with BOLD and G-CSF. *Jpn J Clin Oncol* 35(9):507–513. doi:10.1093/jco/hyi141
323. Borregaard N (2010) Neutrophils, from marrow to microbes. *Immunity* 33(5):657–670. doi:10.1016/j.immuni.2010.11.011
324. Kolaczowska E, Kubes P (2013) Neutrophil recruitment and function in health and inflammation. *Nat Rev Immunol* 13(3):159–175. doi:10.1038/nri3399
325. Dissemmond J, Weimann TK, Schneider LA, Schneeberger A, Scharffetter-Kochanek K, Goos M, Wagner SN (2003) Activated neutrophils exert antitumor activity against human melanoma cells: reactive oxygen species-induced mechanisms and their modulation by granulocyte-macrophage-colony-stimulating factor. *J Invest Dermatol* 121(4):936–938. doi:10.1046/j.1523-1747.2003.12475.x
326. Zivkovic M, Poljak-Blazi M, Egger G, Sunjic SB, Schaur RJ, Zarkovic N (2005) Oxidative burst and anticancer activities of rat neutrophils. *Biofactors* 24(1–4):305–312
327. Clark RA, Klebanoff SJ (1975) Neutrophil-mediated tumor cell cytotoxicity: role of the peroxidase system. *J Exp Med* 141(6):1442–1447
328. Lichtenstein A, Kahle J (1985) Anti-tumor effect of inflammatory neutrophils: characteristics of in vivo generation and in vitro tumor cell lysis. *Int J Cancer* 35(1):121–127
329. Caruso RA, Speciale G, Inferrera C (1994) Neutrophil interaction with tumour cells in small early gastric cancer: ultrastructural observations. *Histol Histopathol* 9(2):295–303
330. Saito H, Fukumura D, Kurose I, Suematsu M, Tada S, Kagawa T, Miura S, Morizane T, Tsuchiya M (1992) Visualization of oxidative processes at the cellular level during neutrophil-mediated cytotoxicity against a human hepatoma cell line, HCC-M. *Int J Cancer* 51(1):124–129
331. Di Carlo E, Forni G, Lollini P, Colombo MP, Modesti A, Musiani P (2001) The intriguing role of polymorphonuclear neutrophils in antitumor reactions. *Blood* 97(2):339–345
332. Yamazaki M, Ikenami M, Sugiyama T (1988) Cytolytic heterogeneity of polymorphonuclear leukocytes in killing of murine tumor cells. *Jpn J Cancer Res* 79(11):1217–1223
333. Koga Y, Matsuzaki A, Suminoe A, Hattori H, Hara T (2004) Neutrophil-derived TNF-related apoptosis-inducing ligand (TRAIL): a novel mechanism of antitumor effect by neutrophils. *Cancer Res* 64(3):1037–1043
334. Al-Benna S, Shai Y, Jacobsen F, Steinstraesser L (2011) Oncolytic activities of host defense peptides. *Int J Mol Sci* 12(11):8027–8051. doi:10.3390/ijms12118027
335. Lichtenstein A, Ganz T, Selsted ME, Lehrer RI (1986) In vitro tumor cell cytotoxicity mediated by peptide defensins of human and rabbit granulocytes. *Blood* 68(6):1407–1410
336. Kushner BH, Cheung NK (1992) Absolute requirement of CD11/CD18 adhesion molecules, FcRII and the phosphatidylinositol-linked FcRIII for monoclonal antibody-mediated neutrophil antitumor cytotoxicity. *Blood* 79(6):1484–1490
337. Iliopoulos D, Ernst C, Stepkowski Z, Jambrosic JA, Rodeck U, Herlyn M, Clark WH Jr, Koprowski H, Herlyn D (1989) Inhibition of metastases of a human melanoma xenograft by monoclonal antibody to the GD2/GD3 gangliosides. *J Natl Cancer Inst* 81(6):440–444
338. Valerius T, Repp R, de Wit TP, Berthold S, Platzer E, Kalden JR, Gramatzki M, van de Winkel JG (1993) Involvement of the high-affinity receptor for IgG (Fc gamma RI; CD64) in enhanced tumor cell cytotoxicity of neutrophils during granulocyte colony-stimulating factor therapy. *Blood* 82(3):931–939
339. Lieschke GJ, Burgess AW (1992) Granulocyte colony-stimulating factor and granulocyte-macrophage colony-stimulating factor (2). *N Engl J Med* 327(2):99–106. doi:10.1056/NEJM199207093270207
340. Clynes RA, Towers TL, Presta LG, Ravetch JV (2000) Inhibitory Fc receptors modulate in vivo cytotoxicity against tumor targets. *Nat Med* 6(4):443–446. doi:10.1038/74704
341. Hernandez-Ilizaliturri FJ, Jupudy V, Ostberg J, Oflazoglu E, Huberman A, Repasky E, Czuczman MS (2003) Neutrophils contribute to the biological antitumor activity of rituximab in a non-Hodgkin's lymphoma severe combined immunodeficiency mouse model. *Clin Cancer Res* 9(16 Pt 1):5866–5873
342. Hubert P, Heitzmann A, Viel S, Nicolas A, Sastre-Garau X, Oppezzo P, Pritsch O, Osinaga E, Amigorena S (2011) Antibody-dependent cell cytotoxicity synapses form in mice during tumor-specific antibody immunotherapy. *Cancer Res* 71(15):5134–5143. doi:10.1158/0008-5472.CAN-10-4222
343. Guettinger Y, Barbin K, Peipp M, Bruenke J, Dechant M, Horner H, Thierschmidt D, Valerius T, Repp R, Fey GH, Stockmeyer B (2010) A recombinant bispecific single-chain fragment variable specific for HLA class II and Fc alpha RI (CD89) recruits polymorphonuclear neutrophils for efficient lysis of malignant B lymphoid cells. *J Immunol* 184(3):1210–1217. doi:10.4049/jimmunol.0902033

344. Stockmeyer B, Dechant M, van Egmond M, Tutt AL, Sundarapandiyam K, Graziano RF, Repp R, Kalden JR, Gramatzki M, Glennie MJ, van de Winkel JG, Valerius T (2000) Triggering Fc alpha-receptor I (CD89) recruits neutrophils as effector cells for CD20-directed antibody therapy. *J Immunol* 165(10):5954–5961
345. Otten MA, Rudolph E, Dechant M, Tuk CW, Reijmers RM, Beelen RH, van de Winkel JG, van Egmond M (2005) Immature neutrophils mediate tumor cell killing via IgA but not IgG Fc receptors. *J Immunol* 174(9):5472–5480
346. Cheung IY, Hsu K, Cheung NK (2012) Activation of peripheral-blood granulocytes is strongly correlated with patient outcome after immunotherapy with anti-GD2 monoclonal antibody and granulocyte-macrophage colony-stimulating factor. *J Clin Oncol* 30(4):426–432. doi:10.1200/JCO.2011.37.6236
347. Mocsai A (2013) Diverse novel functions of neutrophils in immunity, inflammation, and beyond. *J Exp Med* 210(7):1283–1299. doi:10.1084/jem.20122220
348. Tanaka E, Sendo F (1993) Abrogation of tumor-inhibitory MRC-OX8+ (CD8+) effector T-cell generation in rats by selective depletion of neutrophils in vivo using a monoclonal antibody. *Int J Cancer* 54(1):131–136
349. Berger-Achituv S, Brinkmann V, Abed UA, Kuhn LI, Ben-Ezra J, Elhasid R, Zychlinsky A (2013) A proposed role for neutrophil extracellular traps in cancer immunoediting. *Front Immunol* 4:48. doi:10.3389/fimmu.2013.00048
350. Tillack K, Breiden P, Martin R, Sospedra M (2012) T lymphocyte priming by neutrophil extracellular traps links innate and adaptive immune responses. *J Immunol* 188(7):3150–3159. doi:10.4049/jimmunol.1103414
351. Jackaman C, Lew AM, Zhan Y, Allan JE, Koloska B, Graham PT, Robinson BW, Nelson DJ (2008) Deliberately provoking local inflammation drives tumors to become their own protective vaccine site. *Int Immunol* 20(11):1467–1479. doi:10.1093/intimm/dxn104
352. Takami K, Miura K, Takeuchi H, Egawa S, Moriya T, Nakamura Y, Tanabe A, Sugita J, Karasawa H, Unno M, Sasaki I (2008) Granulocyte colony stimulating factor-producing pancreatic cancer: report of a case. *Surg Today* 38(5):453–457. doi:10.1007/s00595-007-3636-z
353. Fujiwara Y, Yamazaki O, Takatsuka S, Kaizaki R, Inoue T (2011) Granulocyte colony-stimulating factor-producing ascending colon cancer as indicated by histopathological findings: report of a case. *Osaka City Med J* 57(2):79–84
354. Kaira K, Ishizuka T, Tanaka H, Tanaka Y, Yanagitani N, Sunaga N, Hisada T, Mori M (2008) Lung cancer producing granulocyte colony-stimulating factor and rapid spreading to peritoneal cavity. *J Thorac Oncol* 3(9):1054–1055. doi:10.1097/JTO.0b013e3181834f7b
355. Matsuyama S, Shimonishi T, Yoshimura H, Higaki K, Nasu K, Toyooka M, Aoki S, Watanabe K, Sugihara H (2008) An autopsy case of granulocyte-colony-stimulating-factor-producing extrahepatic bile duct carcinoma. *World J Gastroenterol* 14(18):2924–2927
356. Kawaguchi M, Asada Y, Terada T, Takehara A, Munemoto Y, Fujisawa K, Mitsui T, Iida Y, Miura S, Sudo Y (2010) Aggressive recurrence of gastric cancer as a granulocyte-colony-stimulating factor-producing tumor. *Int J Clin Oncol* 15(2):191–195. doi:10.1007/s10147-010-0023-3
357. Ito S, Iwai Y, Fujii T, Yoshida N, Hayashi S (1999) Two cases of bladder tumor producing granulocyte colony stimulating factor. *Hinyokika Kyo* 45(1):57–60
358. Nasu K, Inoue C, Takai N, Kashima K, Miyakawa I (2004) Squamous cell carcinoma of the cervix producing granulocyte colony-stimulating factor. *Obstet Gynecol* 104(5 Pt 1):1086–1088. doi:10.1097/01.AOG.0000141552.87313.c8
359. Horii A, Shimamura K, Honjo Y, Mitani K, Miki T, Takashima S, Yoshida J (1997) Granulocyte colony stimulating factor-producing tongue carcinoma. *Head Neck* 19(4):351–356. doi:10.1002/(SICI)1097-0347(199707)19:4<351::AID-HED15>3.0.CO;2-C
360. Sakamoto A, Yamamoto H, Tanaka K, Matsuda S, Harimaya K, Oda Y, Tsuneyoshi M, Iwamoto Y (2006) Dedifferentiated chondrosarcoma with leukocytosis and elevation of serum G-CSF. A case report. *World J Surg Oncol* 4:37
361. Sakamoto A, Matono H, Yoshida T, Tanaka K, Matsuda S, Oda Y, Iwamoto Y (2007) Dedifferentiated liposarcoma with leukocytosis. A case report of G-CSF-producing soft-tissue tumors, possible association with undifferentiated liposarcoma lineage. *World J Surg Oncol* 5:131
362. Droeser RA, Hirt C, Eppenberger-Castori S, Zlobec I, Viehl CT, Frey DM, Nebiker CA, Rosso R, Zuber M, Amicarella F, Iezzi G, Sconocchia G, Heberer M, Lugli A, Tornillo L, Oertli D, Terracciano L, Spagnoli GC (2013) High myeloperoxidase positive cell infiltration in colorectal cancer is an independent favorable prognostic factor. *PLoS One* 8(5):e64814. doi:10.1371/journal.pone.0064814
363. Hirt C, Eppenberger-Castori S, Sconocchia G, Iezzi G, Tornillo L, Terracciano L, Spagnoli GC, Droeser RA (2013) Colorectal carcinoma infiltration by myeloperoxidase-expressing neutrophil granulocytes is associated with favorable prognosis. *Oncotarget* 2(10):e25990. doi:10.4161/onci.25990
364. Sconocchia G, Zlobec I, Lugli A, Calabrese D, Iezzi G, Karamitopoulou E, Patsouris ES, Peros G, Horcic M, Tornillo L, Zuber M, Droeser R, Muraro MG, Mengus C, Oertli D, Ferrone S, Terracciano L, Spagnoli GC (2011) Tumor infiltration by FcgammaRIII (CD16)+ myeloid cells is associated with improved survival in patients with colorectal carcinoma. *Int J Cancer* 128(11):2663–2672. doi:10.1002/ijc.25609
365. Rao HL, Chen JW, Li M, Xiao YB, Fu J, Zeng YX, Cai MY, Xie D (2012) Increased intratumoral neutrophil in colorectal carcinomas correlates closely with malignant phenotype and predicts patients' adverse prognosis. *PLoS One* 7(1):e30806. doi:10.1371/journal.pone.0030806
366. Riesco A (1970) Five-year cancer cure: relation to total amount of peripheral lymphocytes and neutrophils. *Cancer* 25(1):135–140
367. Donskov F (2013) Immunomonitoring and prognostic relevance of neutrophils in clinical trials. *Semin Cancer Biol* 23(3):200–207. doi:10.1016/j.semcancer.2013.02.001
368. Teramukai S, Kitano T, Kishida Y, Kawahara M, Kubota K, Komuta K, Minato K, Mio T, Fujita Y, Yonei T, Nakano K, Tsuboi M, Shibata K, Furuse K, Fukushima M (2009) Pretreatment neutrophil count as an independent prognostic factor in advanced non-small-cell lung cancer: an analysis of Japan Multinational Trial Organisation LC00-03. *Eur J Cancer* 45(11):1950–1958. doi:10.1016/j.ejca.2009.01.023
369. Ubukata H, Motohashi G, Tabuchi T, Nagata H, Konishi S (2010) Evaluations of interferon-gamma/interleukin-4 ratio and neutrophil/lymphocyte ratio as prognostic indicators in gastric cancer patients. *J Surg Oncol* 102(7):742–747. doi:10.1002/jso.21725
370. Donskov F, von der Maase H (2006) Impact of immune parameters on long-term survival in metastatic renal cell carcinoma. *J Clin Oncol* 24(13):1997–2005. doi:10.1200/JCO.2005.03.9594
371. Wang GY, Yang Y, Zhang Q, Li H, Chen GZ, Yi SH, Xu C, Wang GS, Zhang J, Yi HM, Jiang N, Fu BS, Zhao H, Li MR, Chen YH, Cai CJ, Lu MQ, Chen GH (2011) Preoperative neutrophil-lymphocyte ratio as a prognostic predictor after liver transplantation for hepatocellular carcinoma. *Zhonghua Yi Xue Za Zhi* 91(22):1519–1522
372. Hung HY, Chen JS, Yeh CY, Changchien CR, Tang R, Hsieh PS, Tasi WS, You JF, You YT, Fan CW, Wang JY, Chiang JM (2011) Effect of preoperative neutrophil-lymphocyte ratio on the surgical outcomes of stage II colon cancer patients who do not receive adjuvant chemotherapy. *Int J Colorectal Dis* 26(8):1059–1065. doi:10.1007/s00384-011-1192-x



373. Liu H, Ubukata H, Tabuchi T, Takemura A, Motohashi G, Nishimura M, Satani T, Hong J, Katano M, Nakada I, Saniabadi AR (2009) It is possible that tumour-infiltrating granulocytes promote tumour progression. *Oncol Rep* 22(1):29–33
374. Trellakis S, Farjah H, Bruderek K, Dumitru CA, Hoffmann TK, Lang S, Brandau S (2011) Peripheral blood neutrophil granulocytes from patients with head and neck squamous cell carcinoma functionally differ from their counterparts in healthy donors. *Int J Immunopathol Pharmacol* 24(3):683–693
375. Dumitru CA, Fechner MK, Hoffmann TK, Lang S, Brandau S (2012) A novel p38-MAPK signaling axis modulates neutrophil biology in head and neck cancer. *J Leukoc Biol* 91(4):591–598. doi:10.1189/jlb.0411193
376. Dumitru CA, Gholaman H, Trellakis S, Bruderek K, Dominas N, Gu X, Bankfalvi A, Whiteside TL, Lang S, Brandau S (2011) Tumor-derived macrophage migration inhibitory factor modulates the biology of head and neck cancer cells via neutrophil activation. *Int J Cancer* 129(4):859–869. doi:10.1002/ijc.25991
377. Xu X, Wang B, Ye C, Yao C, Lin Y, Huang X, Zhang Y, Wang S (2008) Overexpression of macrophage migration inhibitory factor induces angiogenesis in human breast cancer. *Cancer Lett* 261(2):147–157. doi:10.1016/j.canlet.2007.11.028
378. Ren Y, Tsui HT, Poon RT, Ng IO, Li Z, Chen Y, Jiang G, Lau C, Yu WC, Bacher M, Fan ST (2003) Macrophage migration inhibitory factor: roles in regulating tumor cell migration and expression of angiogenic factors in hepatocellular carcinoma. *Int J Cancer* 107(1):22–29. doi:10.1002/ijc.11287
379. Ren Y, Law S, Huang X, Lee PY, Bacher M, Srivastava G, Wong J (2005) Macrophage migration inhibitory factor stimulates angiogenic factor expression and correlates with differentiation and lymph node status in patients with esophageal squamous cell carcinoma. *Ann Surg* 242(1):55–63
380. Rendon BE, Willer SS, Zundel W, Mitchell RA (2009) Mechanisms of macrophage migration inhibitory factor (MIF)-dependent tumor microenvironmental adaptation. *Exp Mol Pathol* 86(3):180–185. doi:10.1016/j.yexmp.2009.01.001
381. Zhou SL, Dai Z, Zhou ZJ, Wang XY, Yang GH, Wang Z, Huang XW, Fan J, Zhou J (2012) Overexpression of CXCL5 mediates neutrophil infiltration and indicates poor prognosis for hepatocellular carcinoma. *Hepatology* 56(6):2242–2254. doi:10.1002/hep.25907
382. Ietomi K (1990) A study on the role of granulocytes in carcinoma-bearing hosts—G/L ratio as a new host indicator. *Nihon Gan Chiryō Gakkai Shi* 25(3):662–671
383. Terasawa M, Nagata K, Kobayashi Y (2008) Neutrophils and monocytes transport tumor cell antigens from the peritoneal cavity to secondary lymphoid tissues. *Biochem Biophys Res Commun* 377(2):589–594. doi:10.1016/j.bbrc.2008.10.011
384. Kusmartsev S, Nefedova Y, Yoder D, Gabrilovich DI (2004) Antigen-specific inhibition of CD8<sup>+</sup> T cell response by immature myeloid cells in cancer is mediated by reactive oxygen species. *J Immunol* 172(2):989–999
385. Patel KD, Zimmerman GA, Prescott SM, McEver RP, McIntyre TM (1991) Oxygen radicals induce human endothelial cells to express GMP-140 and bind neutrophils. *J Cell Biol* 112(4):749–759
386. Cherdyntseva NV, Bogdashin IV, Nebera SA, Vasil'ev NV (1989) Neutrophil granulocyte function in patients with malignant neoplasms. *Vopr Onkol* 35(4):429–433
387. Li YW, Qiu SJ, Fan J, Zhou J, Gao Q, Xiao YS, Xu YF (2011) Intratumoral neutrophils: a poor prognostic factor for hepatocellular carcinoma following resection. *J Hepatol* 54(3):497–505. doi:10.1016/j.jhep.2010.07.044
388. Tsuda Y, Fukui H, Asai A, Fukunishi S, Miyaji K, Fujiwara S, Teramura K, Fukuda A, Higuchi K (2012) An immunosuppressive subtype of neutrophils identified in patients with hepatocellular carcinoma. *J Clin Biochem Nutr* 51(3):204–212. doi:10.3164/jcbn.12-32
389. Jensen TO, Schmidt H, Moller HJ, Donskov F, Hoyer M, Sjoegren P, Christensen IJ, Steiniche T (2012) Intratumoral neutrophils and plasmacytoid dendritic cells indicate poor prognosis and are associated with pSTAT3 expression in AJCC stage I/II melanoma. *Cancer* 118(9):2476–2485. doi:10.1002/ncr.26511
390. Jensen HK, Donskov F, Marcussen N, Nordmark M, Lundbeck F, von der Maase H (2009) Presence of intratumoral neutrophils is an independent prognostic factor in localized renal cell carcinoma. *J Clin Oncol* 27(28):4709–4717. doi:10.1200/JCO.2008.18.9498
391. Wu Y, Zhao Q, Peng C, Sun L, Li XF, Kuang DM (2011) Neutrophils promote motility of cancer cells via a hyaluronan-mediated TLR4/PI3K activation loop. *J Pathol* 225(3):438–447. doi:10.1002/path.2947
392. Chee DO, Townsend CM Jr, Galbraith MA, Eilber FR, Morton DL (1978) Selective reduction of human tumor cell populations by human granulocytes in vitro. *Cancer Res* 38(12):4534–4539
393. Halbrecht I, Komlos L (1974) Cytotoxic effects of leukocytes and plasma on primary cultures of ovarian carcinoma. *Obstet Gynecol* 43(2):268–275
394. Korec S, Herberman RB, Dean JH, Cannon GB (1980) Cytostasis of tumor cell lines by human granulocytes. *Cell Immunol* 53(1):104–115
395. Gerrard TL, Cohen DJ, Kaplan AM (1981) Human neutrophil-mediated cytotoxicity to tumor cells. *J Natl Cancer Inst* 66(3):483–488
396. Cameron DJ (1983) A comparison of the cytotoxic potential in polymorphonuclear leukocytes obtained from normal donors and cancer patients. *Clin Immunol Immunopathol* 28(1):115–124
397. Cameron DJ (1986) Lidocaine inhibits macrophage mediated cytotoxicity and enhances neutrophil mediated cytotoxicity. *Jpn J Exp Med* 56(6):265–269
398. Cameron DJ (1985) Inhibition of neutrophil-mediated cytotoxicity by exogenous adenosine 5'-triphosphate. *Clin Immunol Immunopathol* 37(2):230–235
399. Cameron DJ, Majeski JA (1988) Inhibition of macrophage- and neutrophil-mediated cytotoxicity by verapamil. *J Surg Oncol* 37(1):5–9
400. Cameron DJ (1986) Role of hydrocortisone, chloroquine and prednisolone in neutrophil mediated cytotoxicity. *Jpn J Exp Med* 56(5):207–212
401. Sauri H, Kim AT, Shau H (1995) Characterization of an oxidation-resistant tumor cell line and its sensitivity to immune response and chemotherapy. *J Surg Res* 58(5):526–535. doi:10.1006/jsre.1995.1083
402. Heidecke H, Eckert K, Schulze-Forster K, Maurer HR (1997) Prothymosin alpha 1 effects in vitro on chemotaxis, cytotoxicity and oxidative response of neutrophils from melanoma, colorectal and breast tumor patients. *Int J Immunopharmacol* 19(8):413–420
403. Godleski JJ, Lee RE, Leighton J (1970) Studies on the role of polymorphonuclear leukocytes in neoplastic disease with the chick embryo and Walker carcinosarcoma 256 in vivo and in vitro. *Cancer Res* 30(7):1986–1993
404. Pickaver AH, Ratcliffe NA, Williams AE, Smith H (1972) Cytotoxic effects of peritoneal neutrophils on a syngeneic rat tumour. *Nat New Biol* 235(58):186–187
405. Fisher B, Saffer EA (1978) Tumor cells cytotoxicity by granulocytes from peripheral blood of tumor-bearing mice. *J Natl Cancer Inst* 60(3):687–691
406. Nathan CF, Brukner LH, Silverstein SC, Cohn ZA (1979) Extracellular cytolysis by activated macrophages and granulocytes. I. Pharmacologic triggering of effector cells and the release of hydrogen peroxide. *J Exp Med* 149(1):84–99
407. Becker S, Haskill S (1980) Non-T-cell-mediated cytotoxicity in mice with tumors induced by Moloney murine sarcoma virus (M-MuSV). II. Granulocyte-mediated cytotoxicity against autochthonous target cells isolated from M-MuSV-induced tumors. *J Natl Cancer Inst* 65(2):469–475



408. Morikawa K, Kamegaya S, Yamazaki M, Mizuno D (1985) Hydrogen peroxide as a tumoricidal mediator of murine polymorphonuclear leukocytes induced by a linear beta-1,3-D-glucan and some other immunomodulators. *Cancer Res* 45(8):3482–3486
409. Morikawa K, Takeda R, Yamazaki M, Mizuno D (1985) Induction of tumoricidal activity of polymorphonuclear leukocytes by a linear beta-1,3-D-glucan and other immunomodulators in murine cells. *Cancer Res* 45(4):1496–1501
410. Lichtenstein AK, Kahle J, Berek J, Zigelboim J (1984) Successful immunotherapy with intraperitoneal *Corynebacterium parvum* in a murine ovarian cancer model is associated with the recruitment of tumor-lytic neutrophils into the peritoneal cavity. *J Immunol* 133(1): 519–526
411. Hirose K, Hakozaki M, Nyunoya Y, Kobayashi Y, Matsushita K, Takenouchi T, Mikata A, Mukaida N, Matsushima K (1995) Chemokine gene transfection into tumour cells reduced tumorigenicity in nude mice in association with neutrophilic infiltration. *Br J Cancer* 72(3):708–714
412. Chen RL, Reynolds CP, Seeger RC (2000) Neutrophils are cytotoxic and growth-inhibiting for neuroblastoma cells with an anti-GD2 antibody but, without cytotoxicity, can be growth-stimulating. *Cancer Immunol Immunother* 48(11):603–612
413. Lozupone F, Luciani F, Venditti M, Rivoltini L, Pupa S, Parmiani G, Belardelli F, Fais S (2000) Murine granulocytes control human tumor growth in SCID mice. *Int J Cancer* 87(4):569–573
414. Jaganjac M, Poljak-Blazi M, Zarkovic K, Schaur RJ, Zarkovic N (2008) The involvement of granulocytes in spontaneous regression of Walker 256 carcinoma. *Cancer Lett* 260(1–2):180–186. doi:10.1016/j.canlet.2007.10.039
415. Jaganjac M, Poljak-Blazi M, Kirac I, Borovic S, Joerg Schaur R, Zarkovic N (2010) Granulocytes as effective anticancer agent in experimental solid tumor models. *Immunobiology* 215(12):1015–1020. doi:10.1016/j.imbio.2010.01.002
416. Jaganjac M, Poljak-Blazi M, Schaur RJ, Zarkovic K, Borovic S, Cipak A, Cindric M, Uchida K, Waeg G, Zarkovic N (2012) Elevated neutrophil elastase and acrolein-protein adducts are associated with W256 regression. *Clin Exp Immunol* 170(2):178–185. doi:10.1111/j.1365-2249.2012.04639.x
417. Ibata M, Takahashi T, Shimizu T, Inoue Y, Maeda S, Tashiro-Yamaji J, Okada M, Ueda K, Kubota T, Yoshida R (2011) Spontaneous rejection of intradermally transplanted non-engineered tumor cells by neutrophils and macrophages from syngeneic strains of mice. *Microbiol Immunol* 55(10):726–735. doi:10.1111/j.1348-0421.2011.00369.x
418. Edelson PJ, Cohn ZA (1973) Peroxidase-mediated mammalian cell cytotoxicity. *J Exp Med* 138(1):318–323
419. Clark RA, Klebanoff SJ, Einstein AB, Fefer A (1975) Peroxidase-H<sub>2</sub>O<sub>2</sub>-halide system: cytotoxic effect on mammalian tumor cells. *Blood* 45(2):161–170
420. Clark RA, Olsson I, Klebanoff SJ (1976) Cytotoxicity for tumor cells of cationic proteins from human neutrophil granules. *J Cell Biol* 70(3):719–723
421. Ikenami M, Yamazaki M (1985) Participation of polymorphonuclear leukocyte-derived factor in murine tumour cell killing. *Br J Cancer* 52(4):575–581
422. Mikami M, Yamazaki M, Yui S (1998) Kinetic analysis of tumor cell death-inducing mechanism by polymorphonuclear leukocyte-derived calprotectin: involvement of protein synthesis and generation of reactive oxygen species in target cells. *Microbiol Immunol* 42(3):211–221
423. Noh H, Eomm M, Han A (2013) Usefulness of pretreatment neutrophil to lymphocyte ratio in predicting disease-specific survival in breast cancer patients. *J Breast Cancer* 16(1):55–59. doi:10.4048/jbc.2013.16.1.55
424. Azab B, Bhatt VR, Phookan J, Murukutla S, Kohn N, Terjanian T, Widmann WD (2012) Usefulness of the neutrophil-to-lymphocyte ratio in predicting short- and long-term mortality in breast cancer patients. *Ann Surg Oncol* 19(1):217–224. doi:10.1245/s10434-011-1814-0
425. Oh BS, Jang JW, Kwon JH, You CR, Chung KW, Kay CS, Jung HS, Lee S (2013) Prognostic value of C-reactive protein and neutrophil-to-lymphocyte ratio in patients with hepatocellular carcinoma. *BMC Cancer* 13:78. doi:10.1186/1471-2407-13-78
426. Tibaldi C, Vasile E, Bernardini I, Orlandini C, Andreuccetti M, Falcone A (2008) Baseline elevated leukocyte count in peripheral blood is associated with poor survival in patients with advanced non-small cell lung cancer: a prognostic model. *J Cancer Res Clin Oncol* 134(10):1143–1149. doi:10.1007/s00432-008-0378-2
427. Satomi A, Murakami S, Ishida K, Mastuki M, Hashimoto T, Sonoda M (1995) Significance of increased neutrophils in patients with advanced colorectal cancer. *Acta Oncol* 34(1):69–73
428. Shibutani M, Maeda K, Nagahara H, Noda E, Ohtani H, Nishiguchi Y, Hirakawa K (2013) A high preoperative neutrophil-to-lymphocyte ratio is associated with poor survival in patients with colorectal cancer. *Anticancer Res* 33(8):3291–3294
429. Chua W, Charles KA, Baracos VE, Clarke SJ (2011) Neutrophil/lymphocyte ratio predicts chemotherapy outcomes in patients with advanced colorectal cancer. *Br J Cancer* 104(8):1288–1295. doi:10.1038/bjc.2011.100
430. He W, Yin C, Guo G, Jiang C, Wang F, Qiu H, Chen X, Rong R, Zhang B, Xia L (2013) Initial neutrophil lymphocyte ratio is superior to platelet lymphocyte ratio as an adverse prognostic and predictive factor in metastatic colorectal cancer. *Med Oncol* 30(1): 439. doi:10.1007/s12032-012-0439-x
431. Ohno Y, Nakashima J, Ohori M, Hatano T, Tachibana M (2010) Pretreatment neutrophil-to-lymphocyte ratio as an independent predictor of recurrence in patients with nonmetastatic renal cell carcinoma. *J Urol* 184(3):873–878. doi:10.1016/j.juro.2010.05.028
432. Atzpodien J, Royston P, Wandert N, Reitz M (2003) Metastatic renal carcinoma comprehensive prognostic system. *Br J Cancer* 88(3): 348–353. doi:10.1038/sj.bjc.6600768
433. Can C, Baseskioglu B, Yilmaz M, Colak E, Ozen A, Yenilmez A (2012) Pretreatment parameters obtained from peripheral blood sample predicts invasiveness of bladder carcinoma. *Urol Int* 89(4): 468–472. doi:10.1159/000343278
434. Perisanidis C, Kornek G, Poschl PW, Holzinger D, Pirklbauer K, Schopper C, Ewers R (2013) High neutrophil-to-lymphocyte ratio is an independent marker of poor disease-specific survival in patients with oral cancer. *Med Oncol* 30(1):334. doi:10.1007/s12032-012-0334-5
435. Sharaiha RK, Halazun KJ, Mirza F, Port JL, Lee PC, Neugut AI, Altorki NK, Abrams JA (2011) Elevated preoperative neutrophil: lymphocyte ratio as a predictor of postoperative disease recurrence in esophageal cancer. *Ann Surg Oncol* 18(12):3362–3369. doi:10.1245/s10434-011-1754-8
436. Sato H, Tsubosa Y, Kawano T (2012) Correlation between the pretherapeutic neutrophil to lymphocyte ratio and the pathologic response to neoadjuvant chemotherapy in patients with advanced esophageal cancer. *World J Surg* 36(3):617–622. doi:10.1007/s00268-011-1411-1
437. Rashid F, Waraich N, Bhatti I, Saha S, Khan RN, Ahmed J, Leeder PC, Larvin M, Iftikhar SY (2010) A pre-operative elevated neutrophil: lymphocyte ratio does not predict survival from oesophageal cancer resection. *World J Surg Oncol* 8:1. doi:10.1186/1477-7819-8-1
438. Wang L, Lin Y, Long H, Liu H, Rao H, He Y, Rong T, Liang Y (2013) Esophageal carcinosarcoma: a unique entity with better prognosis. *Ann Surg Oncol* 20(3):997–1004. doi:10.1245/s10434-012-2658-y
439. Kim YH, Choi WJ (2012) The effectiveness of postoperative neutrophils to lymphocytes ratio in predicting long-term recurrence after stomach cancer surgery. *J Korean Surg Soc* 83(6):352–359. doi:10.4174/jkss.2012.83.6.352
440. Kao SC, Pavlakis N, Harvie R, Vardy JL, Boyer MJ, van Zandwijk N, Clarke SJ (2010) High blood neutrophil-to-lymphocyte ratio is an indicator of poor prognosis in malignant mesothelioma patients

- undergoing systemic therapy. *Clin Cancer Res* 16(23):5805–5813. doi:[10.1158/1078-0432.CCR-10-2245](https://doi.org/10.1158/1078-0432.CCR-10-2245)
441. Kao SC, Klebe S, Henderson DW, Reid G, Chatfield M, Armstrong NJ, Yan TD, Vardy J, Clarke S, van Zandwijk N, McCaughan B (2011) Low calretinin expression and high neutrophil-to-lymphocyte ratio are poor prognostic factors in patients with malignant mesothelioma undergoing extrapleural pneumonectomy. *J Thorac Oncol* 6(11):1923–1929. doi:[10.1097/JTO.0b013e31822a3740](https://doi.org/10.1097/JTO.0b013e31822a3740)
442. Pinato DJ, Mauri FA, Ramakrishnan R, Wahab L, Lloyd T, Sharma R (2012) Inflammation-based prognostic indices in malignant pleural mesothelioma. *J Thorac Oncol* 7(3):587–594. doi:[10.1097/JTO.0b013e31823f45c1](https://doi.org/10.1097/JTO.0b013e31823f45c1)
443. Wang S, Zhang Z, Fang F, Gao X, Sun W, Liu H (2011) The neutrophil/lymphocyte ratio is an independent prognostic indicator in patients with bone metastasis. *Oncol Lett* 2(4):735–740. doi:[10.3892/ol.2011.304](https://doi.org/10.3892/ol.2011.304)