

## OPINION

# The seven deadly sins of DNA barcoding

R. A. COLLINS\* and R. H. CRUICKSHANK†

\*Bio-Protection Research Centre, Lincoln University, PO Box 84, Lincoln, 7647, Canterbury, New Zealand, †Department of Ecology, Faculty of Agriculture and Life Sciences, Lincoln University, Lincoln, 7647, Canterbury, New Zealand

## Abstract

Despite the broad benefits that DNA barcoding can bring to a diverse range of biological disciplines, a number of shortcomings still exist in terms of the experimental design of studies incorporating this approach. One underlying reason for this lies in the confusion that often exists between species discovery and specimen identification, and this is reflected in the way that hypotheses are generated and tested. Although these aims can be associated, they are quite distinct and require different methodological approaches, but their conflation has led to the frequently inappropriate use of commonly used analytical methods such as neighbour-joining trees, bootstrap resampling and fixed distance thresholds. Furthermore, the misidentification of voucher specimens can also have serious implications for end users of reference libraries such as the Barcode of Life Data Systems, and in this regard we advocate increased diligence in the a priori identification of specimens to be used for this purpose. This commentary provides an assessment of seven deficiencies that we identify as common in the DNA barcoding literature, and outline some potential improvements for its adaptation and adoption towards more reliable and accurate outcomes.

**Keywords:** bootstrap, DNA barcoding, hypothesis testing, neighbour joining, reference library, species identification

Received 5 June 2012; revision received 27 August 2012; accepted 9 November 2012

## Introduction

DNA barcoding is now a well established method, and has been shown to bring substantial benefits to applications such as food product regulation (Becker *et al.* 2011), conservation (Francis *et al.* 2010), and investigating species interactions (Valentini *et al.* 2009). Nevertheless, we feel that a number of misconceptions have pervaded the DNA barcoding literature, and these are manifested in shortcomings of experimental design and analytical procedure. We have loosely identified seven such problems, and here we highlight each of them and suggest how improvements could be made. Most of these concerns have been raised previously in the literature, but as DNA barcoding approaches its tenth year, a timely synthesis is perhaps required, especially when DNA barcoding may in future provide admissible evidence in wildlife crime cases (Alacs *et al.* 2010; Linacre & Tobe 2011).

Our main concern is over the goal of DNA barcoding (Moritz & Cicero 2004; De Salle 2006; Rubinoff *et al.* 2006; Goldstein & De Salle 2011; Taylor & Harris 2012). We acknowledge that DNA barcoding (*sensu* Hebert *et al.*

2003) can comprise two distinct aims: specimen identification and species discovery (Schindel & Miller 2005). Specimen identification involves assigning taxonomic names to unknown specimens using a DNA reference library of morphologically preidentified vouchers. Here, DNA barcoding is particularly useful for applications such as biosecurity (Collins *et al.* 2012a) or marketplace substitution (Lowenstein *et al.* 2010). Species discovery with DNA barcodes, on the other hand, is best thought of as a quick and dirty 'molecular parataxonomy' process, analogous to physically sorting specimens into morpho-species (Brower 2006). Due to the relative speed at which COI sequence data can be generated and analysed, DNA barcoding therefore represents a powerful triage tool for biodiversity assessment, quickly sorting collections into species-like units (Schindel & Miller 2005). We use the term 'species-like units' to acknowledge that although short-length single-locus markers such as COI are often effective proxies for species, they are frequently not representative of full phylogenetic history (i.e. the species tree), due to their idiosyncratic behaviour (Brower *et al.* 1996; Rubinoff *et al.* 2006; Dasmahapatra *et al.* 2010; Dupuis *et al.* 2012; Fujita *et al.* 2012). Of course, subsequent to an initial DNA barcode triage, COI data can then be incorporated into more sophisticated species delimitation systems using multiple loci (Dupuis *et al.* 2012; Fujita

Correspondence: R. A. Collins, Fax: +64 3 325 3864; E-mail: rupertcollins@gmail.com

*et al.* 2012), and ultimately into an integrative (Will *et al.* 2005; Padial *et al.* 2010; Goldstein & De Salle 2011) or DNA-based taxonomy framework (Tautz *et al.* 2003; Vogler & Monaghan 2007). We specifically make the point of distinguishing relatively crude single-locus methods such as DNA barcoding as ‘species discovery’, and multilocus/integrative methods as ‘species delimitation’ (Sites & Marshall 2004).

We feel that both of the outlined aims of DNA barcoding (specimen identification and species discovery) are uncontroversial, provided that they are clearly defined. However, several authors have raised repeated concerns regarding the blurring of these boundaries (De Salle *et al.* 2005; De Salle 2006; Vogler & Monaghan 2007; Meier 2008; Goldstein & De Salle 2011), and it seems impossible to separate these objectives in many examples from the DNA barcoding literature. This provides the basis for many of the criticisms herein.

### **Sin number one: failure to test clear hypotheses**

Perhaps one of the gravest sins in many barcoding studies is the lack of clearly stated, objective hypotheses. A ‘typical’ barcoding study—‘DNA barcoding the [insert taxon] of [insert geographic region]’—aims to: (i) assemble a DNA reference library from specimens identified to species using morphological characters; (ii) test how effective this DNA library is for future identification purposes and then (iii) explore previously unrecognized diversity apparent in the DNA barcodes. However, it is in regard to these three steps that we feel there is often confusion in how hypotheses are generated and tested. Too frequently, objectives (ii) and (iii) are conflated, and methodological approaches do not appear to reflect these different goals (Meier 2008; Goldstein & De Salle 2011). Analytical techniques presented in many studies do not explicitly set out to test identification success (objective ii) by simulating a quantified identification scenario. Rather, they tend to employ the same method (usually a neighbour-joining tree) to test both objectives (ii) and (iii), and usually present a descriptive summary rather than a hypothesis-based test of the data. If the data collected are intended to be used as an identification tool, then they should be tested as such. Conversely, if a study aims to test the suitability of DNA barcoding as a biodiversity assessment tool (species discovery), then hypotheses of species richness should be estimated independently of the taxonomic names, and then compared *a posteriori*.

In light of these distinct goals, we recommend defining each objective more clearly in the methods section of the work, and explicitly separating the experimental procedures used to achieve each aim.

### **Sin number two: inadequate a priori identification of specimens**

A serious limitation to the utility of DNA barcoding as a practical resource for regulation and molecular diagnostics is human error and uncertainty in creating and curating reference libraries. Becker *et al.* (2011) identify this as the primary source of error in FISH-BOL barcode data. Conflicting identifications can be made when multiple labs are working on the same taxa, and in the process of their morphological identifications they ascribe different taxonomic names to the same species.

As an example of the potential severity of the problem, we investigated each BIN cluster (Barcode Index Number) in the ‘RCYY’ project of ornamental cyprinid fishes (Collins *et al.* 2012a) on BOLD, the Barcode of Life Data System (Ratnasingham & Hebert 2007, URL: <http://v3.boldsystems.org/>). A total of 54 BINs in the project contained data from external projects (as of February 13, 2012). Of the 54, 19 BINs (35%) contained more than one species name, meaning that BOLD was unable at that time to offer an unambiguous species-level identification for these taxa. Even more worryingly, the same analysis was repeated some months later (June 26, 2012); the number of BINs found to contain data from external projects increased to 70, and the proportion of these BINs comprising taxonomic conflict (more than one species name) increased to 53% (=37).

A crucial aspect of DNA barcoding is the maintenance of records, supporting information and voucher specimens; this is what sets BOLD apart from GenBank (Ratnasingham & Hebert 2007). However, there are currently few safeguards against misidentifying a specimen, and once a name has been added to a database, it may be difficult for a third party to convince data managers that it should be changed. A new feature of BOLD v3.0 is a wiki-like framework for community-based annotation of barcode data (<http://v3.boldsystems.org/>), but preemptive solutions are perhaps preferable. To this effect, a system of identification confidence has been proposed, which rates identifications according to the degree of expertise used and effort made (Steinke & Hanner 2011). This should encourage increasing diligence over how identifications are generated and justified.

The importance of accurate identification is obvious (Bortolus 2008), and providing a bibliography of reference material and morphological characters used for identification should be mandatory for publication (Vink *et al.* 2012). These additional metadata may be extremely valuable for correcting mistakes without recourse to the effort of loaning and reexamining voucher material. To quantify the extent to which identifications are unjustified, all articles published in *Molecular Ecology Resources*

from 2010 to 2012 (up to August 16), with either 'DNA barcoding' or 'DNA barcodes' in the title were downloaded (total 61 articles). Of these, 46 presented empirical DNA barcode data, but of those only 16 (35%) cited literature to support their identifications. Only three presented morphological characters to directly support each taxon identification.

### **Sin number three: the use of the term 'species identification'**

The term 'species identification' is ubiquitous in the DNA barcoding literature, but in our view this terminology is misleading, and reflects a long-standing confusion between the two subdisciplines of DNA barcoding (specimen identification vs. species discovery; see above). We interpret 'species identification' here as shorthand for: identification of biological material—a specimen—to the level of species. However, we are concerned that it can be seen as species discovery or species delimitation (as used in Ferguson 2002). One way to minimize this confusion and to clarify the distinct role of each of the two separate objectives is to use the terms 'specimen identification' or 'species discovery' in place of 'species identification', as appropriate. This more objectively states what hypotheses are being tested, and better ensures that identification of individuals is not confused with discovery of groups. Both of these aims fall within the purview of DNA barcoding, but they should be clearly distinguished as they require different methodological and analytical approaches.

### **Sin number four: inappropriate use of neighbour-joining trees**

Almost every DNA barcoding study presents a neighbour-joining (NJ) tree as part of the standard analytical procedure (Casiraghi *et al.* 2010). NJ trees were initially presented in the DNA barcoding literature by Hebert *et al.* (2003), although the method had been used prior to that for similar purposes (e.g. Hsieh *et al.* 2001). NJ trees use a distance matrix of sequence similarity to produce a hierarchical clustering phenogram (Little & Stevenson 2007). They are fast and easy to compute for large data sets, and are always fully resolved (i.e. do not produce ambiguous polytomies). However, several authors have expressed concerns as to whether this kind of phylogenetic approach to DNA barcoding is appropriate (Will & Rubinoff 2004; Meier *et al.* 2008; Goldstein & De Salle 2011).

For specimen identification purposes, it has been well documented both empirically and theoretically that NJ trees perform poorly (Meier *et al.* 2006; Virgilio *et al.* 2010; Little 2011; Zhang *et al.* 2012). They can be mislead-

ing to interpret, especially in conjunction with an incompletely sampled reference library. Unless nested directly within a cluster, the tree alone presents no way to discern if an unknown belongs to the closest topological species or not; i.e. it is unable to provide a 'no identification' result when an exact match is not present (Will *et al.* 2005; Collins *et al.* 2012a). It is important to note at this point that problems with NJ trees are not resolved by using any other tree inference method such as maximum likelihood or parsimony. The problem is with relying on tree topologies and monophyly (in a topological rather than cladistic sense) as an identification criterion. In situations of incomplete lineage sorting and species-level paraphyly, tree-based identification methods will result in ambiguous or incorrect identifications (Lowenstein *et al.* 2010). Few species concepts require reciprocal monophyly (Meier 2008), and in any case, monophyly of mtDNA lineages can be an unrealistic scenario in many closely related groups (Funk & Omland 2003; Zhang *et al.* 2012).

Despite the popularity and intuitiveness of NJ trees, identification success generally improves when using more accurate tree-free techniques, which are usually based directly on the genetic distance matrix. The 'best close match' method (Meier *et al.* 2006) has been shown to be reliable, predictable, and computationally tractable (Virgilio *et al.* 2010; Collins *et al.* 2012a). Intra-/interspecific variation is discriminated by a user-set threshold (but see sin number six). The method is easy to interpret, being one of the most simple available, but is able to make identifications even in the presence of paraphyly. Alternatively, many other criteria are also available for measuring identification success (see Casiraghi *et al.* 2010), and comparisons of performance between some of these have already been made (Meier *et al.* 2006; Little & Stevenson 2007; Ross *et al.* 2008; Austerlitz *et al.* 2009; Virgilio *et al.* 2010; Zhang *et al.* 2012).

It is important to note, however, that a quantification of monophyly still remains a useful description of the data, and should still be used in conjunction with other methods. Problems mostly occur when NJ trees are the only analytical method presented, and identification success rates are not quantified (Little & Stevenson 2007). Likewise, for the purpose of graphically summarizing the data, NJ trees can be considered appropriate. It is also worth mentioning that the ubiquitous use of the Kimura two-parameter (K2P) model for constructing initial distance matrices is also questionable; uncorrected *p* distances should be used instead (Collins *et al.* 2012b; Srivathsan & Meier 2012).

In some scenarios, character-based methods using diagnostic nucleotide combinations may be preferable (Davis & Nixon 1992; Brower 1999; De Salle 2007), and this is particularly the case for small groups of closely

related taxa where similarity methods perform poorly (Lowenstein *et al.* 2009). However, character-based approaches such as those implemented in the CAOS software (Sarkar *et al.* 2008), have yet to be fully characterized in terms of their sensitivity to taxon sampling and homoplasy, and are therefore at present perhaps limited to restricted applications (Kerr *et al.* 2009). The use of discrete characters could be seen in terms of 'DNA barcoding 2.0', potentially offering additional benefits after sampling is extended beyond simply collecting baseline data.

Ultimately, phenetic (similarity) methods using genetic distances may be regarded as something of a stop-gap solution. In the near future, the problem of accurately assigning identifications is likely to be addressed by either likelihood-based information-theoretic approaches, or machine learning and statistical tools such as supervised classification and pattern recognition (e.g. Zhang *et al.* 2008; Austerlitz *et al.* 2009). Bayesian coalescent methods also promise statistical advantages, but may be too computationally inefficient in their current incarnations (Zhang *et al.* 2012).

When evaluating DNA barcoding as a species discovery tool, a method is required that can approximate the number of species in a sample directly from the DNA sequence data, and independently from the preassigned taxonomic names (i.e. the data set used to subsequently measure congruence between the two approaches). NJ trees are a poor choice in this respect, as on their own, they are unable to make a quantitative assessment. Techniques such as the Automatic Barcode Gap Discovery method (Puillandre *et al.* 2012), the general mixed Yule coalescent model (Monaghan *et al.* 2009), or the BOLD v3.0 BIN system are able to use genetic information to generate an estimate of the number of species-like units, and should therefore be used in preference to NJ trees for this kind of study.

#### **Sin number five: inappropriate use of bootstrap resampling**

The use of bootstrap resampling in DNA barcoding studies typifies the confusion between species discovery and specimen identification. The bootstrap, along with reciprocal monophyly, is one method among many that can be used to test whether a species-like cluster is well supported, and can be used as part of a species discovery/delimitation process (i.e. a test of genetic distinctiveness for a potentially new taxon). Bootstrapping in this situation also helps address problems with NJ trees such as taxon-order bias and tied trees (Meier 2008; Lowenstein *et al.* 2009).

However, the use of bootstrapping for specimen identification is somewhat perplexing. The aim here is to

maximize congruence with a priori defined species, *viz.* the taxonomic names from a morphological identification process. A species with low bootstrap support does not falsify a species hypothesis when this assessment was based on independent data (i.e. morphology from the original description). In many cases, recently diverged sister species on short branches will have low support and therefore fail to be identified, even if they are morphologically distinct and diagnosable by unique mutations (Lowenstein *et al.* 2009). Thus, using a bootstrap value as a cut-off for correct identification severely compromises the efficacy of a reference library (Collins *et al.* 2012a; Zhang *et al.* 2012), and exacerbates the previously outlined weaknesses of using tree-based methods in general. On top of this, bootstrap resampling does not make an assessment of the uncertainty in identification; an unknown can group with a reference specimen at 100% bootstrap support, and yet be an entirely different species. Perhaps an improved way to measure uncertainty in identification is to better understand interspecific threshold distances (see below), calculate group membership probabilities (Zhang *et al.* 2012) and make explicit 'caveats in relation to the breath [sic] of sampling' (Moritz & Cicero 2004).

#### **Sin number six: inappropriate use of fixed distance thresholds**

The use of distance thresholds has been extensively debated (see references in Puillandre *et al.* 2012; Virgilio *et al.* 2012; Zhang *et al.* 2012), so our aim is not to cover old ground, but to reemphasize points already made. A threshold is essential when identifying specimens using genetic distance data; in the absence of complete sampling, distance thresholds aim to minimize misidentifications of unknowns that do not have conspecifics represented in the reference library (Virgilio *et al.* 2012). However, there is no a priori reason to assume that a universal threshold is applicable, as coalescent depths among species will vary considerably due to differences in population size, rate of mutation and time since speciation (Monaghan *et al.* 2009; Fujita *et al.* 2012).

A generic threshold such as 1% is perhaps not an unreasonable heuristic in some cases, but it can be considered arbitrary, and is likely to suffer from varying rates of false-positive and false-negative error, depending on the data. Rather than relying on prescribed cut-offs, optimized thresholds should be generated directly from the data (Meyer & Paulay 2005; Virgilio *et al.* 2012). Software and protocols are now available to calculate these optimized thresholds, and for species discovery, can even be generated in the absence of taxonomic names (Brown *et al.* 2012; Puillandre *et al.* 2012; Virgilio *et al.* 2012). However, a newly developed

fuzzy-set-theory technique (Zhang *et al.* 2012) perhaps obviates the requirement for distance thresholds, offering a more sophisticated group membership parameter that provides additional information lacking in threshold-based implementations.

### Sin number seven: incorrectly interpreting the barcoding gap

The barcoding gap as proposed by Meyer & Paulay (2005) can represent two distinct scenarios: one for specimen identification, with an individual being closer to a member of its own species than a different species (i.e. a 'local' barcoding gap); and one for species discovery, a distance that equates to a threshold applicable to all species (i.e. a 'global' barcoding gap). Although this has been previously discussed (Wiemers & Fiedler 2007), the two scenarios are frequently confused, and this again demonstrates conflation of the two objectives of DNA barcoding.

Many DNA barcoding studies present histograms showing frequency distributions of both intra- and interspecific divergences for all pooled species analysed in a

study. Overlap between the two distributions can be interpreted as a failure of DNA barcoding, but the only failure demonstrated in this case is that of defining a universal cut-off value. In this regard, and as stated previously, it is widely acknowledged that coalescent depths vary among species, and substantial overlap between intra- and interspecific distances may be the rule, rather than the exception (Wiemers & Fiedler 2007; Virgilio *et al.* 2010). Therefore, for specimen identification purposes this type of presentation is wholly uninformative, as intraspecific distances for one species can exceed interspecific distances for other species in the analysis, but without necessarily compromising identification success (the local gap).

A better display of distance data for specimen identification is a dotplot in which, for each individual in the data set, the distance to the furthest conspecific is plotted against the distance to the nearest nonconspecific, with a 1:1 slope representing the point at which the difference between the two is zero (i.e. no local barcoding gap). An example of this method is illustrated in Fig. 4 of Robinson *et al.* (2009). It is also important to note that these statistics should be generated using the smallest

**Table 1** Outline of potential problems, consequences and solutions for the 'seven deadly sins' of DNA barcoding, as presented here

Problem	Consequence	Solution
Failure to test clear hypotheses	Choice of inappropriate or suboptimal analytical method due to confusion as to the objectives of the study	Explicitly state each hypothesis, and for each distinct aspect of the study present separate headings in methods and results sections
Inadequate a priori identification of specimens	Conflicting identifications made by different labs can compromise the effectiveness of reference libraries that are ultimately used as a resource for scientific or regulatory purposes	Present a bibliography of references, as well as the distinguishing morphological characters used in the identification process. Follow recommendations outlined by Steinke & Hanner (2011)
The use of the term 'species identification'	Confusion between identification of individuals, and delimitation/discovery of species	To clarify objectives, use the term 'specimen identification' or 'species discovery' where appropriate
Inappropriate use of neighbour-joining trees	(a) Relying on strict monophyly for identification can reduce the apparent effectiveness of DNA barcoding as an identification tool. This can be due to either mtDNA paralogy or misidentification of specimens. (b) For biodiversity assessment and species discovery, NJ trees cannot estimate the number of species independently with respect to the taxonomic names	(a) Alternative criteria such as 'best close match' are readily available, and have higher rates of identification success. This method can be implemented using the free software packages TaxonDNA (Meier <i>et al.</i> 2006) or Spider (Brown <i>et al.</i> 2012). (b) Estimate species richness using ABGD (Puillandre <i>et al.</i> 2012), GMYC (Monaghan <i>et al.</i> 2009) or BOLD's BIN system ( <a href="http://v3.boldsystems.org">http://v3.boldsystems.org</a> )
Inappropriate use of bootstrap resampling	For specimen identification purposes, bootstrap resampling can further reduce the already low identification success rates associated with NJ trees	Only use bootstrapping where appropriate: e.g. as part of a species delimitation process on preestimated groups
Inappropriate use of fixed distance thresholds	For specimen identification purposes, a generic threshold which is set too low or high can reduce or bias identification error rates	Thresholds can now be optimized for specific data sets using the method of Virgilio <i>et al.</i> (2012), or with software such as ABGD (Puillandre <i>et al.</i> 2012) and Spider (Brown <i>et al.</i> 2012)
Incorrectly interpreting the barcoding gap	Overlapping distributions of intra-/interspecific distances do not necessarily mean that barcodes perform poorly for identification	For specimen identification studies, dotplots of intra-/interspecific distances are a better way to illustrate the barcoding gap (e.g. Robinson <i>et al.</i> 2009)

rather than the mean interspecific distance (Meier *et al.* 2008).

## Summary

We hope that future barcoding studies will push forward improvements in data analysis, and make more use of alternative methods. One possible cause of the limited uptake of some new approaches may be the lack of a common platform for carrying out analyses (Sarkar & Trizna 2011). Exploring different analytical methods is important, but many software applications have quite different input and output formats, making comparison between them difficult. Fortunately, it is now increasingly possible to conduct a variety of analyses in a universal, open-source environment such as R (R Development Core Team 2011), which could potentially supersede the inflexible and piecemeal software applications that are currently available (Freckleton 2009). This is one solution that could ultimately encourage improved dissemination, sharing and benchmarking of new techniques among labs.

In conclusion, we feel that more care should be taken in setting clear hypotheses for DNA barcoding studies, and choosing appropriate methods for testing each distinct hypothesis. Given the opportunities DNA barcoding can offer to both biodiversity research and regulatory science, it would be unfortunate to compromise its potential due to misconceptions or inadequacies in study design. Table 1 provides an outline of possible solutions to some of the problems outlined herein.

## Acknowledgements

This work was supported by funding from the Ministry for Primary Industries Biosecurity New Zealand and the Bio-Protection Research Centre for RAC. The authors thank Karen Armstrong, Rudolf Meier, Laura Boykin and two reviewers for suggesting improvements to earlier versions of the manuscript.

## References

- Alacs EA, Georges A, FitzSimmons NN, Robertson J (2010) DNA detective: a review of molecular approaches to wildlife forensics. *Forensic Science, Medicine and Pathology*, **6**, 180–194.
- Austerlitz F, David O, Schaeffer B (2009) DNA barcode analysis: a comparison of phylogenetic and statistical classification methods. *BMC Bioinformatics*, **10**, S10.
- Becker S, Hanner R, Steinke D (2011) Five years of FISH-BOL: brief status report. *Mitochondrial DNA*, **22**(Suppl. 1), 3–9.
- Bortolus A (2008) Error cascades in the biological sciences: the unwanted consequences of using bad taxonomy in ecology. *Ambio*, **37**, 114–118.
- Brower AVZ (1999) Delimitation of phylogenetic species with DNA sequences: a critique of Davis and Nixon's population aggregation analysis. *Systematic Biology*, **48**, 199–213.
- Brower AVZ (2006) Problems with DNA barcodes for species delimitation: 'ten species' of *Astraptus fulgerator* reassessed (Lepidoptera: Hesperidae). *Systematics and Biodiversity*, **4**, 127–132.
- Brower AVZ, De Salle R, Vogler A (1996) Gene trees, species trees, and systematics: a cladistic perspective. *Annual Review of Ecology and Systematics*, **27**, 423–450.
- Brown SDJ, Collins RA, Boyer S (2012) Spider: an R package for the analysis of species identity and evolution, with particular reference to DNA barcoding. *Molecular Ecology Resources*, **12**, 562–565.
- Casiraghi M, Labra M, Ferri E, Galimberti A, De Mattia F (2010) DNA barcoding: a six-question tour to improve users' awareness about the method. *Briefing in Bioinformatics*, **11**, 440–453.
- Collins RA, Armstrong KF, Meier R (2012a) Barcoding and border biosecurity: identifying cyprinid fishes in the aquarium trade. *PLoS ONE*, **7**, e28381.
- Collins RA, Boykin LM, Cruickshank RH, Armstrong KF (2012b) Barcoding's next top model: an evaluation of nucleotide substitution models for specimen identification. *Methods in Ecology and Evolution*, **3**, 457–465.
- Dasmahapatra KK, Elias M, Hill RI, Hoffman JI, Mallet J (2010) Mitochondrial DNA barcoding detects some species that are real, and some that are not. *Molecular Ecology Resources*, **10**, 264–273.
- Davis JJ, Nixon KC (1992) Populations, genetic variation, and the delimitation of phylogenetic species. *Systematic Biology*, **41**, 421–435.
- De Salle R (2006) Species discovery versus species identification in DNA barcoding efforts: response to Rubinoff. *Conservation Biology*, **20**, 1545–1547.
- De Salle R (2007) Phenetic and DNA taxonomy; a comment on Waugh. *BioEssays*, **29**, 1289–1290.
- De Salle R, Egan MG, Siddall M (2005) The unholy trinity: taxonomy, species delimitation and DNA barcoding. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **360**, 1905–1916.
- Dupuis JR, Roe AD, Sperling FAH (2012) Multi-locus species delimitation in closely related animals and fungi: one marker is not enough. *Molecular Ecology*, **21**, 4422–4436.
- Ferguson JWH (2002) On the use of genetic divergence for identifying species. *Biological Journal of the Linnean Society*, **75**, 509–516.
- Francis CM, Borisenko AV, Ivanova NV (2010) The role of DNA barcodes in understanding and conservation of mammal diversity in Southeast Asia. *PLoS ONE*, **5**, e12575.
- Freckleton RP (2009) The seven deadly sins of comparative analysis. *Journal of Evolutionary Biology*, **22**, 1367–1375.
- Fujita MK, Leaché AD, Burbrink FT, McGuire JA, Moritz C (2012) Coalescent-based species delimitation in an integrative taxonomy. *Trends in Ecology and Evolution*, **27**, 480–488.
- Funk DJ, Omland KE (2003) Species-level paraphyly and polyphyly: frequency, causes, and consequences, with insights from animal mitochondrial DNA. *Annual Review of Ecology, Evolution, and Systematics*, **34**, 397–423.
- Goldstein PZ, De Salle R (2011) Integrating DNA barcode data and taxonomic practice: determination, discovery and description. *BioEssays*, **33**, 135–147.
- Hebert PDN, Cywinska A, Ball SL, de Waard JR (2003) Biological identifications through DNA barcodes. *Proceedings of the Royal Society B: Biological Sciences*, **270**, 313–321.
- Hsieh HM, Chiang HL, Tsai LC (2001) Cytochrome *b* gene for species identification of the conservation animals. *Forensic Science International*, **122**, 7–18.
- Kerr KCR, Birks SM, Kalyakin MV, Red'kin YA, Koblik EA, Hebert PDN (2009) Filling the gap - COI barcode resolution in eastern Palearctic birds. *Frontiers in Zoology*, **6**, 1–13.
- Linacre A, Tobe SS (2011) An overview to the investigative approach to species testing in wildlife forensic science. *Investigative Genetics*, **2**, 1–9.
- Little DP (2011) DNA barcode sequence identification incorporating taxonomic hierarchy and within taxon variability. *PLoS ONE*, **6**, e20552.
- Little DP, Stevenson DW (2007) A comparison of algorithms for the identification of specimens using DNA barcodes: examples from gymnosperms. *Cladistics*, **23**, 1–21.

- Lowenstein JH, Amato G, Kolokotronis SO (2009) The real *maccoyii*: identifying tuna sushi with DNA barcodes - contrasting characteristic attributes and genetic distances. *PLoS ONE*, **4**, e7866.
- Lowenstein JH, Burger J, Jeitner CW, Amato G, Kolokotronis SO, Gochfeld M (2010) DNA barcodes reveal species-specific mercury levels in tuna sushi that pose a health risk to consumers. *Biology Letters*, **6**, 692–695.
- Meier R (2008) DNA sequences in taxonomy: opportunities and challenges. In *The New Taxonomy* (ed. Wheeler QD), Chap. 7, pp. 95–127. CRC Press, New York.
- Meier R, Shiyang K, Vaidya G, Ng PKL (2006) DNA barcoding and taxonomy in Diptera: a tale of high intraspecific variability and low identification success. *Systematic Biology*, **55**, 715–728.
- Meier R, Zhang G, Ali F (2008) The use of mean instead of smallest interspecific distances exaggerates the size of the 'barcoding gap' and leads to misidentification. *Systematic Biology*, **57**, 809–813.
- Meyer CP, Paulay G (2005) DNA barcoding: error rates based on comprehensive sampling. *PLoS Biology*, **3**, 2229–2238.
- Monaghan MT, Wild R, Elliot M (2009) Accelerated species inventory on Madagascar using coalescent-based models of species delineation. *Systematic Biology*, **58**, 298–311.
- Moritz C, Cicero C (2004) DNA barcoding: promise and pitfalls. *PLoS Biology*, **2**, e354.
- Padial JM, Miralles A, De la Riva I, Vences M (2010) The integrative future of taxonomy. *Frontiers in Zoology*, **7**, 1–14.
- Puillandre N, Lambert A, Brouillet S, Achaz G (2012) ABGD, Automatic Barcode Gap Discovery for primary species delimitation. *Molecular Ecology*, **21**, 1864–1877.
- R Development Core Team (2011) *R: A language and environment for statistical computing*. R Development Core Team, Vienna, Austria. Available from <http://www.r-project.org/>.
- Ratnasingham S, Hebert PDN (2007) BOLD: the barcode of life data system. *Molecular Ecology Notes*, **7**, 355–364. URL: [www.barcodinglife.org](http://www.barcodinglife.org)
- Robinson EA, Blagoev GA, Hebert PDN, Adamowicz SJ (2009) Prospects for using DNA barcoding to identify spiders in species-rich genera. *ZooKeys*, **46**, 27–46.
- Ross HA, Murugan S, Li WLS (2008) Testing the reliability of genetic methods of species identification via simulation. *Systematic Biology*, **57**, 216–230.
- Rubinoff D, Cameron S, Will K (2006) A genomic perspective on the shortcomings of mitochondrial DNA for 'barcoding' identification. *Journal of Heredity*, **97**, 581–594.
- Sarkar IN, Trizna M (2011) The Barcode of Life Data Portal: bridging the biodiversity informatics divide for DNA barcoding. *PLoS ONE*, **6**, e14689.
- Sarkar IN, Planet PJ, De Salle R (2008) CAOS software for use in character-based DNA barcoding. *Molecular Ecology Resources*, **8**, 1256–1259.
- Schindel DE, Miller SE (2005) DNA barcoding a useful tool for taxonomists. *Nature*, **435**, 17.
- Sites JWJ, Marshall JC (2004) Operational criteria for delimiting species. *Annual Review of Ecology, Evolution and Systematics*, **35**, 199–227.
- Srivathsan A, Meier R (2012) On the inappropriate use of Kimura-2-parameter (K2P) divergences in the DNA-barcoding literature. *Cladistics*, **28**, 190–194.
- Steinke D, Hanner R (2011) The FISH-BOL collaborators' protocol. *Mitochondrial DNA*, **22**(Suppl. 1), 10–14.
- Tautz D, Arctander P, Minelli A, Thomas RH, Vogler AP (2003) A plea for DNA taxonomy. *Trends in Ecology and Evolution*, **18**, 70–74.
- Taylor HR, Harris WE (2012) An emergent science on the brink of irrelevance: a review of the past 8 years of DNA barcoding. *Molecular Ecology Resources*, **12**, 377–388.
- Valentini A, Pompanon F, Taberlet P (2009) DNA barcoding for ecologists. *Trends in Ecology and Evolution*, **24**, 110–117.
- Vink CJ, Paquin P, Cruickshank RH (2012) Taxonomy and irreproducible biological science. *BioScience*, **62**, 451–452.
- Virgilio M, Backeljau T, Nevado B, De Meyer M (2010) Comparative performances of DNA barcoding across insect orders. *BMC Bioinformatics*, **11**, 206.
- Virgilio M, Jordaens K, Breman FC, Backeljau T, De Meyer M (2012) Identifying insects with incomplete DNA barcode libraries, African fruit flies (Diptera: tephritidae) as a test case. *PLoS ONE*, **7**, e31581.
- Vogler AP, Monaghan MT (2007) Recent advances in DNA taxonomy. *Journal of Zoological Systematics and Evolutionary Research*, **45**, 1–10.
- Wiemers M, Fiedler K (2007) Does the DNA barcoding gap exist? A case study in blue butterflies (Lepidoptera: lycaenidae). *Frontiers in Zoology*, **4**, 8.
- Will KW, Rubinoff D (2004) Myth of the molecule: DNA barcodes for species cannot replace morphology for identification and classification. *Cladistics*, **20**, 47–55.
- Will KW, Mishler BD, Wheeler QD (2005) The perils of DNA barcoding and the need for integrative taxonomy. *Systematic Biology*, **54**, 844–851.
- Zhang AB, Sikes DS, Muster C, Li SQ (2008) Inferring species membership using DNA sequences with back-propagation neural networks. *Systematic Biology*, **57**, 202–215.
- Zhang AB, Muster C, Liang HB (2012) A fuzzy-set-theory-based approach to analyse species membership in DNA barcoding. *Molecular Ecology*, **21**, 1848–1863.